

Supplementary material

Long term impacts of *Phytophthora cinnamomi* infestation on heathy woodland in the Great Otway National Park in south-eastern Australia

B. A. Wilson^{A,C}, *K. Annett*^A, *W. S. Laidlaw*^A, *D. M. Cahill*^A, *M. J. Garkaklis*^B
and *L. Zhuang-Griffin*^A

^ASchool of Life and Environmental Sciences, Deakin University,
Geelong, Vic. 3217, Australia.

^BState of the Environment Pty Ltd, PO Box 14, Aireys Inlet, Vic. 3231, Australia.

^CCorresponding author. Email: barbara.wilson@deakin.edu.au.

Table S1. Species percentage contribution (of the cumulative 90% contribution) from similarity percentages routine analysis in non-infected, active and post disease *Phytophthora* disease quadrats surveyed in 2015

Species	Average abundance	Percentage contribution	Cumulative percentage
Non-infected			
<i>Xanthorrhoea australis</i>	1.79	10.53	10.53
Litter cover	1.55	9.07	19.6
<i>Leptospermum continentale</i>	1.56	8.78	28.38
<i>Dillwynia glaberrima</i>	1.32	7.22	35.6
Bare ground	1.21	6.42	42.02
<i>Gahnia radula</i>	1.22	6.1	48.12
<i>Leptospermum myrsinoides</i>	1.29	5.69	53.81
<i>Monotoca scoparia</i>	1.18	5.06	58.86
Lichen	0.92	4.25	63.11
Moss	1	4.16	67.27
<i>Banksia marginata</i>	1.01	4.14	71.41
<i>Eucalyptus willisii</i>	1.07	3.99	75.4
<i>Epacris impressa</i>	0.84	2.66	78.06
<i>Lepidosperma filiforme</i>	0.74	2.52	80.59
<i>Cassytha glabella</i>	0.6	2.07	82.66
<i>Drosera whittakeri</i>	0.7	2.07	84.73
<i>Aotus ericoides</i>	0.74	2.06	86.79
<i>Schoenus tenuissimus</i>	0.71	1.82	88.61
<i>Drosera peltata peltata</i>	0.58	1.61	90.22
Active			
<i>Leptospermum continentale</i>	1.68	12.15	12.15
<i>Gahnia radula</i>	1.44	10.03	22.18
<i>Xanthorrhoea australis</i>	1.54	9.97	32.15
Litter cover	1.36	8.31	40.46
Lichen	1.03	6.87	47.33
<i>Leptospermum myrsinoides</i>	1.26	6.24	53.57
Bare ground	1.09	6.21	59.78
<i>Lepidosperma filiforme</i>	1.06	5.2	64.98
<i>Cassytha glabella</i>	0.92	4.91	69.89
Moss	0.88	4.88	74.76
<i>Banksia marginata</i>	0.94	4.56	79.32
<i>Dillwynia glaberrima</i>	0.85	4.37	83.69
<i>Monotoca scoparia</i>	0.8	3.24	86.93
<i>Eucalyptus willisii</i>	0.75	2.57	89.51
<i>Aotus ericoides</i>	0.5	1.23	90.74
Post-disease			
<i>Leptospermum continentale</i>	2.05	15.05	15.02
<i>Gahnia radula</i>	1.84	13.59	28.61
Litter cover	1.54	11.12	39.73
Bare ground	1.38	9.34	49.07
Lichen	1.13	6.68	55.76
<i>Leptospermum myrsinoides</i>	1.02	4.28	60.03
Moss	0.84	3.98	64.01
<i>Dillwynia glaberrima</i>	0.9	3.8	67.81
<i>Lindsaea linearis</i>	0.78	3.44	71.25
<i>Hypolaena fastigiata</i>	0.83	3.07	74.32
<i>Banksia marginata</i>	0.78	2.86	77.18
<i>Lepidosperma filiforme</i>	0.85	2.82	80
<i>Cassytha glabella</i>	0.68	2.69	82.69
<i>Schoenus tenuissimus</i>	0.82	2.62	85.31
<i>Eucalyptus willisii</i>	0.76	2.21	87.52
<i>Burchardia umbellata</i>	0.56	2.09	89.61
<i>Acacia suaveolens</i>	0.67	1.91	91.52