

[10.1071/BT22006](https://doi.org/10.1071/BT22006)

*Australian Journal of Botany*

### Supplementary Material

**Evaluating conservation strategies for the endangered daisy *Schoenia filifolia* subsp. *subulifolia* (Asteraceae): fitness consequences of genetic rescue and hybridisation with a widespread subspecies**

*Leonie Monks<sup>A,B,\*</sup>, David Coates<sup>A</sup>, Shelley McArthur<sup>A</sup>, and Rachel J. Standish<sup>B</sup>*

<sup>A</sup>Biodiversity Conservation Science, Department of Biodiversity, Conservation and Attractions, Locked Bag 104, Bentley Delivery Centre, Bentley, WA 6983, Australia.

<sup>B</sup>Harry Butler Institute, Murdoch University, 90 South Street, Murdoch, WA 6150, Australia.

\*Correspondence to: Leonie Monks Biodiversity Conservation Science, Department of Biodiversity, Conservation and Attractions, Locked Bag 104, Bentley Delivery Centre, Bentley, WA 6983, Australia  
Email: [leonie.monks@dbca.wa.gov.au](mailto:leonie.monks@dbca.wa.gov.au)

**Table S1. Description of 7 microsatellite loci in *Schoenia filifolia*. For individuals sampled from six populations, information is given for locus name, primer sequence, GenBank accession number, repeat motif, and the size range of observed alleles in base pairs.**

Locus	Primer Sequence	GenBank Accession No.	Repeat Motif	Allele size range (bp)
SCF01	F:ACGGGCAGATAGTATAAAGTACA R:TACAGGTACAGGCGTGATGA	MT520980	(AC) <sub>9</sub>	85-104
SCF04	F:CGTCAAGAGTTGAAGTTAGCGAT R:AAATTAGGGCTTCTAGTTCTGCTCT	MT520981	(AAC) <sub>8</sub>	112-130
SCF08	F:AGCGCCAAGATGACATCACA R:TGGTTGGTTTGAATATGGAGTT	MT520982	(AC) <sub>7</sub>	118-132
SCF16	F:ACCACCTTTCATTACATCACC R:ACGAGTTTGTTGCTAAGATCTGC	MT520983	(ATC) <sub>8</sub>	126-164
SCF20	F:GGTGCAGTACGGTTAGTGGT R:AGAAGAACTTCTCCAATTACGCT	MT520984	(AAT) <sub>6</sub>	143-173
SCF23	F:ACCAGGTTGCCCGTGATG R:TTCATCCACCTGGTGTCCG	MT520985	(AAC) <sub>6</sub>	216-225
SCF35	F:TGGTCATGCATGGGTCCAC R:TGCCATAGTTCAAATTCAGTCATTCT	MT520986	(AAG) <sub>6</sub>	217-236

**Table S2. Number of plants of *Schoenia filifolia* used for each cross type. Each plant donated pollen to and received pollen from another plant within the same cross type.**

		Lockier	Mingenew	Tardun	Morawa	Black Rd	Kondinin
<i>S. filifolia</i> subsp.	Lockier	20	20	20	20	10	0
<i>subulifolia</i>	Mingenew		20	20	20	10	0
<i>S. filifolia</i> subsp.	Tardun			20	20	20	10
<i>filifolia</i>	Morawa				20	20	16
	Black Rd					16	16
	Kondinin						8