

# Particle aggregation, pH changes and metal behaviour during estuarine mixing: review and integration

Luke M. Mosley<sup>A,C</sup> and Peter S. Liss<sup>B</sup>

<sup>A</sup>School of Biological Sciences, University of Adelaide Waite Campus, PMB 1, Glen Osmond, SA 5064, Australia.

<sup>B</sup>Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK.

<sup>C</sup>Corresponding author. Email: [luke.mosley@adelaide.edu.au](mailto:luke.mosley@adelaide.edu.au)

**Abstract.** Estuaries are dynamic mixing zones where river water interacts with seawater, resulting in large and complex geochemical changes. How two key factors, particle aggregation and pH, affect metal behaviour in estuaries is reviewed and integrated in this paper. Riverine particles are coated with organic matter and electrostatic repulsive forces restrict aggregation. In estuaries, increased concentrations of divalent cations reduce the repulsive forces between particles at low salinities, resulting in their rapid coagulation and removal of particulate-associated metals (e.g. Fe and Pb). However, truly dissolved metals may mix conservatively, and metals associated more with colloidal and dissolved organic material (e.g. Cu and Zn) can show variable behaviour. In many field studies and modelling of river inputs with different compositions, pH decreases slightly at low salinity. Geochemical model simulations of dissolved metal speciation indicated that Zn would be desorbed from iron oxide binding surfaces due to these pH and cation concentration changes, with Cu also showing less binding to dissolved organic matter (DOM). DOM, pH and particle surfaces can influence individual metal behaviour at various spatial and temporal scales. Further integrated field and laboratory research in estuaries where key geochemical processes affecting metal concentrations are measured and modelled is needed.

Received 26 May 2019, accepted 30 June 2019, published online 4 October 2019

## Introduction

Estuaries are complex geochemical reactors that can greatly modify the delivery of dissolved and suspended material from rivers to the ocean. Understanding how pollutants behave in estuarine environments is important because they have very high environmental and socioeconomic value (e.g. culturally, aesthetically, fisheries and recreation). Although there have been many field and laboratory studies on estuaries over the past 40–50 years, there are gaps remaining in our understanding of geochemical processes and how they interact. The aim of this paper is to review and integrate how two fundamental processes, particle aggregation and pH changes, influence metal dynamics in estuaries.

The degree to which metals sorb or desorb to particles in estuaries, and how those particles behave, is a very important driver of their fate and transport in estuaries (Burton and Liss 1976; Benoit *et al.* 1994; Dai and Martin 1995; Waeles *et al.* 2008; Illuminati *et al.* 2019). Metals adsorbed to particles greater than ~1 µm in size can be removed from the water column by sedimentation. However, at low salinity, particles <1 µm (colloids) will not settle out of the water column even over long periods of time because their gravitational settling rate is less than their random displacement by the Brownian motion of the surrounding water molecules. In estuaries, the destabilising influence of the seawater cations can lead to aggregation of

colloidal material (Sholkovitz 1976, 1978; Boyle *et al.* 1977; Mosley *et al.* 2003). This also provides a mechanism for the removal of other metals that have a tendency to be adsorbed to colloid surfaces (e.g. Pb, Zn). In estuaries affected by pollution, significant amounts of anthropogenic metal inputs may end up in the sediment (Liu *et al.* 2019). Following deposition, mixing and redistribution of sediment phases and associated metals can occur (Reese *et al.* 2019). Flux of metals to the water column from the sediment arises due to diffusion or resuspension and oxidation of sulfide phases (Richards *et al.* 2018; Duan *et al.* 2019).

Binding of some metals (e.g. Cu) to colloidal and dissolved organic matter (DOM) is also very important in estuaries because a quite different geochemistry can be observed for these metals compared with the predominantly particle-associated metals (Santschi *et al.* 1997; Tang *et al.* 2002; Illuminati *et al.* 2019). The binding behaviour of individual metals to DOM, and the functional composition of the DOM itself, can also vary with salinity in estuaries (Zhou *et al.* 2016; Ksionzek *et al.* 2018). In addition, pH and inorganic carbon speciation may vary markedly in an estuary as river water, with typically lower pH and alkalinity (except in some calcareous drainage basins), mixes with seawater. The partial pressure of carbon dioxide ( $p\text{CO}_2$ ) is often higher in rivers due to system respiration, driven by microbial degradation of organic matter, exceeding net primary production (Cole and Caraco 2001).

Typically, pH increases towards seawater values in a non-linear way in estuaries, and decreasing pH has also been observed at low salinity (Mook and Koene 1975; Mosley *et al.* 2004, 2010). pH is a ‘master variable’ controlling metal sorption to particles surfaces (Dzombak and Morel 1990), so it could be expected that pH changes in estuaries will also greatly affect metal behaviour. The increased ionic strength, and changing inorganic and organic ligand concentration, in estuaries could also result in altered dissolved metal speciation and binding compared with river water.

Processes affecting trace metal behaviour in an estuary can be inferred from field studies, but it is often difficult to separate the effects of physicochemical processes occurring in the water column from the effects of biological activity or input of material from the underlying sediment. Controlled laboratory experiments in which river water and seawater are mixed in varying amounts to simulate the likely salinity gradient in an estuary have been widely used to provide information on processes occurring in the water column generally on short time scales (Hunter 1983). By comparing the laboratory results to the results from field studies for the same estuary, it may be possible to identify other processes that affect the behaviour of trace metals (Sholkovitz *et al.* 1978; Shiller and Boyle 1991). The speciation and partitioning of trace metals in estuaries has also been calculated using a knowledge of the equilibrium constants for the interaction of metal ions with various ligands and particle surfaces (Mantoura *et al.* 1978; Bourg 1987; Tipping *et al.* 1998; Hamilton-Taylor *et al.* 2002; Pearson *et al.* 2018). However, to the best of our knowledge, all the potential factors affecting metal behaviour (pH, ionic strength, metal oxide surfaces, inorganic and organic complexation) have not been integrated.

To achieve the aim of this review, the relevant international peer-reviewed literature in this area was synthesised. Geochemical model simulations were also performed to help elucidate why individual metal behaviour may vary in estuaries.

### Particle aggregation due to increasing salinity in estuaries

River-borne particles entering estuaries can be very diverse in terms of their origin, mineralogy, hydrous metal (Fe, Mn) oxide concentration, elemental composition and size distribution. However, the surface properties of particles are usually rendered quite uniform by the ubiquitous surface film of natural organic matter that envelops the particles (Hunter and Liss 1979, 1982). Exceptions to this general rule may occur in rivers that are very rich in Fe particulates and relatively low in organic matter (Newton and Liss 1987). Almost all particles are found to be negatively charged, which is attributed to the presence of carboxyl acid groups ( $pK_a \approx 4.5$ ) on the organic matter dominating the surface chemistry (Hunter 1980). The negative charge on the particle–organic matter surface creates an electrostatic ‘double layer’ that extends from the particle surface. The first layer in the double layer is the charged surface that originates from the dissociated functional groups (e.g.  $\text{COO}^-$ ) and tightly absorbed or substituted ions. The second layer corresponds to the diffuse layer, which contains the neutralising charge consisting of accumulated counter ions (cations in the case of negatively charged surfaces in natural waters) and depleted co-ions (i.e. different concentration compared with the bulk

solution). Atomic force microscopy (AFM) measurements have shown that the double layer distance controls the range of interparticle repulsive forces and is sufficient to prevent particle aggregation in fresh water (Mosley *et al.* 2003; Sander *et al.* 2004). However, coagulation may occur in cation-rich rivers before they reach their estuaries.

The flocculation of riverine colloids in estuaries is driven by the increase in the amount of seawater cations such as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  (Eckert and Sholkovitz 1976; Boyle *et al.* 1977; Nowostawska *et al.* 2008). The multivalent ions  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  are more effective than singly charged cations ( $\text{Na}^+$ ,  $\text{K}^+$ ) at screening the electrostatic repulsive force in the diffuse double layer, which then compresses closer to the surface of the particles (Fig. 1). The distance the double layer extends is called the Debye length ( $\delta_D$ ; nm) and can be approximated in estuaries by the following relationship (Stumm and Morgan 1996):

$$\delta_D = 0.28 \div I^{0.5}$$

where  $I$  is ionic strength. In fresh water with a low ionic strength ( $I \approx 0.0001$  M),  $\delta_D$  extends  $\sim 30$  nm, which prevents particles from aggregating. By a salinity of  $\sim 1$  ( $I \approx 0.02$  M), the electric double layer is compressed to  $\sim 2$  nm, which is close enough for attractive short-range van der Waals forces to occur (Fig. 1; Mosley *et al.* 2003). These forces are insensitive to salinity, and aggregate and stick together particles that come into close contact. Interparticle forces in river–estuarine water have been measured directly and predicted quantitatively (Zhou *et al.* 1994; Mosley *et al.* 2003; Sander *et al.* 2004) using classical DLVO theory (Verwey and Overbeek 1948). The organic matter on the surface can introduce additional steric forces but, based on AFM measurements, these do not appear to provide a barrier to aggregation except at short interparticle separations (Mosley *et al.* 2003; Sander *et al.* 2004). Conformational changes may also occur on the natural organic matter due to increased salinity that assist in coiling it or compressing it closer to the particle surface (Kruger *et al.* 2011). Organic matter coatings and divalent cations may create additional bridging mechanisms that influence particle aggregation and separation forces (Mosley *et al.* 2003). Different organic matter amounts and sources relative to the particulate load may result in different behaviour in this regard (Furukawa *et al.* 2014). The role of transparent exopolymeric particles (TEP), released by phytoplankton and other microorganisms, has also been shown to enhance aggregation in estuaries, particularly at mid to high salinity ranges (Wetz *et al.* 2009; Mari *et al.* 2012).

The removal of colloidal particles by coagulation in estuaries was elucidated by the early laboratory studies of Sholkovitz (1976), who used a batch mixing approach (a portion of river water mixed with a portion of seawater to create a particular salinity). Continuous mixing approaches may more accurately simulate the mixing processes in an estuary, where the mixing occurs gradually and in a continuous and sequential manner (Hunter *et al.* 1997). Using such approaches, the kinetics of the particle coagulation process have been found to be fast, with a significant amount of aggregation occurring within seconds to minutes of salinity increasing in estuaries (Hunter and Leonard 1988; Nowostawska *et al.* 2008; Gigault *et al.* 2018). The aggregation rate has been found to be dependent on salinity




Fig. 1. Schematic illustration of key processes driving particle aggregation in estuaries.

but independent of the mixing rate and temperature (Hunter *et al.* 1997). The presence of larger ( $>1\ \mu\text{m}$ ) particles has been found to enhance the aggregation process in laboratory experiments due to increasing interparticle collisions (Hunter and Leonard 1988). However, it is important to note that particle dynamics in real estuaries are much more complex than in laboratory experimental settings (e.g. Wang *et al.* 2013). For example, redistribution of particles can occur very dynamically in response to riverine, tidal and wind-driven currents and, along with salinity stratification, create what is known as the estuarine turbidity maxima (McLachlan *et al.* 2017). Hence, it can be difficult to understand and separate the different particle aggregation and dynamic processes occurring in estuaries, although there has been some success in simulating these (Trento and Vinzon 2014; Ramírez-Mendoza *et al.* 2014).

### Metal behaviour during estuarine mixing

Metals in estuarine water are predominantly derived from riverine inputs, atmospheric deposition and fluxes from the sediment, all of which can have natural and anthropogenic sources. Metal behaviour in different estuaries is often interpreted by comparing concentrations that would be expected if conservative mixing (linear interpolation between freshwater and seawater end member metal concentrations and salinity) v. non-conservative mixing (removal or addition of metals above conservative mixing line) occurred (Liss 1976). Table 1 shows a collation of metal behaviour results from different studies for selected different metals that are commonly studied (Fe, Pb, Cu, Zn) and have different propensities for partitioning to

particulate, organic matter or truly dissolved solution phases. As shown below, these differences can explain why these metals have quite different estuarine mixing behaviour. However, it should be noted that there are potential pitfalls of the conservative mixing line approach because it assumes it is possible to identify a unique freshwater end-member composition when the river water may come from several sources, which themselves are time variant or may enter within the estuarine mixing zone. External sources of metals can be significant in contaminated estuaries (Liu *et al.* 2019). Fixing the seawater end-member composition is often more straightforward due to its relative homogeneity. Other problems include the collection of sufficient samples to cover the whole salinity range, the effect of high suspended loads making pre-analysis filtration difficult and analytical difficulties arising from the wide salinity range of the samples. Further information and a discussion of issues arising in the interpretation of results can be found in Liss (1976).

Fe is found mainly in particulate (including colloidal)  $\text{Fe}^{\text{III}}$  oxide form in rivers, which is its most stable form in air-saturated waters at near neutral pH values (e.g. Boyle *et al.* 1977; Hunter 1983; Dai and Martin 1995). River inputs of Fe to estuaries can show large seasonal variation (Fu *et al.* 2013). Fe has exhibited non-conservative removal behaviour in estuaries across a wide range of different studies (see Table 1). The aggregation of colloids by the mechanisms described above is the accepted reason for the observed removal of Fe in estuaries and, indeed, Fe concentration has been used as a proxy measure for colloid concentration in laboratory experiments (Hunter and Leonard 1988). However, with regard to the truly dissolved Fe

**Table 1. Estuarine mixing behaviour of metals (Fe, Pb, Zn, Cu) in selected studies globally**

Conservative v. non-conservative patterns reported is for the  $\sim <0.4\text{-}\mu\text{m}$  filtered fraction, although truly dissolved metals may behave differently to this fraction as noted in the text

Metal	Estuarine behaviour	Reference and estuary studied
Fe	Non-conservative mixing (removal)	Boyle <i>et al.</i> (1974), Merrimack, USA Boyle <i>et al.</i> (1977), multiple estuaries in the US Holliday and Liss (1976), Beaulieu, UK Sholkovitz (1976, 1978), Scotland Hunter (1983), Taieri, New Zealand Yan <i>et al.</i> (1991), New Jersey, USA Dai and Martin (1995), Ob and Yenisey, Russia Powell (1996), Ochlockonee, USA Fu <i>et al.</i> (2013), Wanquan and Wenchang, Wenjiao, China Mulholland <i>et al.</i> (2015), Amazon, Brazil
Pb	Non-conservative mixing (removal)	Chiffolleau <i>et al.</i> (1994), Seine, France Dai and Martin (1995), Ob and Yenisey, Russia Wen <i>et al.</i> (1999), Galveston, USA Waeles <i>et al.</i> (2008), Penzé, France Tanguy <i>et al.</i> (2011), Penzé, France Fu <i>et al.</i> (2013), Wanquan and Wenchang, Wenjiao, China Illuminati <i>et al.</i> (2019), Po River plume, Adriatic Sea
Cu	Conservative mixing	Boyle <i>et al.</i> (1982), Amazon, Brazil Windom <i>et al.</i> (1983), multiple estuaries, south-eastern USA Shiller and Boyle (1991), Mississippi, USA Guieu and Martin (2002), Danube Sulina Branch Dai <i>et al.</i> (1995), Rhône, France Dai and Martin (1995), Ob and Yenisey, Russia
	Non-conservative mixing (removal)	Benoit <i>et al.</i> (1994), Texas, USA Guieu and Martin (2002), Danube Chilia Branch, Ukraine Waeles <i>et al.</i> (2005, 2008), Penzé, France Illuminati <i>et al.</i> (2019), Po River plume, Adriatic Sea
	Non-conservative mixing (addition)	Ackroyd <i>et al.</i> (1986), Tamar, UK Hunter (1983), Taieri, New Zealand van den Berg <i>et al.</i> (1987), Scheldt, UK Chiffolleau <i>et al.</i> (1994), Seine, France Guieu <i>et al.</i> (1996), Lena, Russia Waeles <i>et al.</i> (2005), Penzé, France
Zn	Conservative mixing	Holliday and Liss (1976), Beaulieu, UK Shiller and Boyle (1991), Mississippi, USA Benoit <i>et al.</i> (1994), Texas, USA Guieu <i>et al.</i> (1996), Lena, Russia
	Non-conservative mixing (addition)	Ackroyd <i>et al.</i> (1986), Tamar, UK van den Berg <i>et al.</i> (1987), Scheldt, UK Yan <i>et al.</i> (1991), New Jersey, USA Chiffolleau <i>et al.</i> (1994), Seine, France Benoit <i>et al.</i> (1994), Texas, USA Guieu and Martin (2002), Danube, Ukraine

fraction (typically a much lower concentration than the solid Fe phase), Dai and Martin (1995) found it behaved conservatively in the Ob and Yenisey estuaries. Some dissolved and colloidal

Fe-binding constituents of aquatic humic substances also appear not to be subject to coagulation along the estuarine salinity gradient (Merschel *et al.* 2017; Muller 2018). Hence, although it seems likely much particulate Fe is deposited to estuarine sediments, a portion can still be transported through estuaries to the coastal ocean (Sanders *et al.* 2015). The application of advanced isotope techniques has been shown to be useful to trace riverine sediment–metal redistribution in estuaries (Reese *et al.* 2019). The redox state of the water column and sediment is also an important control of Fe speciation and flux. The formation of iron (hydr)oxides is effective for the removal of trace metals under oxidising conditions, whereas the formation of sulfides can be conducive to the removal of trace metals under reducing conditions (Duan *et al.* 2019). Resuspension and oxidation of solid FeS phases can also release dissolved Fe and other metals (Richards *et al.* 2018).

Pb has also been found predominantly in the particulate (including colloidal) fraction of rivers and estuaries (Erel *et al.* 1991; Benoit *et al.* 1994; Benoit 1995; Tanguy *et al.* 2011). Pb shows strong binding to iron oxide phases at typical estuarine, circumneutral, pH values (Dzombak and Morel 1990). Consistent with this partitioning, the behaviour of Pb has generally showed non-conservative removal in the low salinity ( $<5$ ) zone across multiple studies (see studies in Table 1). Rare earth elements, which also strongly partition to iron oxides, also show strong removal behaviour in estuaries (Hoyle *et al.* 1984; Sholkovitz 1995; Merschel *et al.* 2017). This indicates the estuarine mixing profile of particle-reactive elements is dominantly controlled by the extent of partitioning to colloidal material (e.g. iron oxides) and the aggregation of the colloids typically removes these metals from solution. However, like Fe, the truly dissolved Pb fraction, which is much lower in concentration, may show conservative mixing (Dai and Martin 1995; Tanguy *et al.* 2011) and increase to form a large component of the total dissolved Pb approaching seawater values (Illuminati *et al.* 2019). The dynamics of Pb in estuaries can be disturbed by anthropogenic inputs and the activity of benthic organisms (Duan *et al.* 2019).

Measurements of Cu speciation have indicated that most ( $>99\%$ ) of the Cu in rivers (Sunda and Hanson 1979) and estuaries (van den Berg *et al.* 1987; Paulson *et al.* 1994) is strongly bound by an unidentified organic ligand or group of ligands present at low concentrations. In terms of organic matter size fraction partitioning, some studies have found that Cu was predominantly found in the truly dissolved ( $<10$  kDa) size fraction (Benoit *et al.* 1994; Paulson *et al.* 1994; Greenamoyer and Moran 1997), whereas others have found that the majority of the Cu was in the colloidal fraction (Dai and Martin 1995; Wen *et al.* 1996). During estuarine mixing, the behaviour of Cu has also been observed to be variable, with differing partitioning behaviour (conservative, addition, removal) found between different studies (Table 1). Laboratory mixing experiments have also shown a mixed behaviour with, for example, Sholkovitz (1978) finding a 40% removal of Cu during estuarine mixing (Water of Luce), whereas Shiller and Boyle (1991) found that Cu mixed conservatively (Mississippi River). This variable behaviour has been postulated to be dependent on whether Cu is bound to colloidal organic matter fractions (typically high molecular weight humic acids) that can flocculate out of



solution and remove Cu or to DOM, which may behave conservatively and hence also lead to the conservative behaviour of Cu (Sholkovitz 1978; Dai and Martin 1995). Laboratory experiments have shown that the binding of Cu to organic ligands may be unaffected by increases in salinity in estuaries (Hamilton-Taylor *et al.* 2002). However, biological ligand production and metal uptake may affect Cu concentrations (Guieu and Martin 2002; Illuminati *et al.* 2019). Waeles *et al.* (2008) found that seasonal variation in Cu complexation, with the production of strongly complexing hydrophobic organic ligands by phytoplankton in summer, increased organic complexation. This also highlights a further limitation of the conservative mixing line approach, which assumes organic and other ligands that bind metals are not produced internally during estuarine mixing. In general, the observed variable behaviour of Cu is probably to be expected because it is too simplistic to treat all rivers as having the same properties (e.g. pH, organic matter amount and type, as well as the amount and type of suspended solids).

Zn has also been found to be strongly organically complexed (but to a lesser extent than Cu) to ligands present at low concentration (van den Berg *et al.* 1987; Bruland 1989). In terms of partitioning, Wen *et al.* (1999) found that on average 91% of Zn was present in the 1-kDa or 0.4- $\mu\text{m}$  (colloidal) fraction, which is similar to the partitioning also found in Texan estuaries (Benoit *et al.* 1994). Similar to Cu, Zn has also exhibited variable behaviour in estuaries, with addition above the conservative mixing line in some studies and conservative behaviour in others (Table 1). Guieu and Martin (2002) postulated that Zn was behaving more similarly to inorganic solid or colloidal phases in estuaries, which could be expected if predominantly colloid partitioning occurred. Many studies have highlighted addition of Zn occurring at low salinity (Table 1), which would not be expected if the binding to colloid surfaces was maintained during the early stages of estuarine mixing. Church (1986) found Zn was desorbed when unfiltered river water was used in a mixing experiment, whereas conservative mixing was observed when filtered water was used. The potential role of pH changes in estuaries leading to desorption of Zn from particles (non-conservative addition) is discussed and modelled further below. There can also be dynamic production of different-sized colloids, sediment : water fluxes, resuspension and sorption-desorption reactions that can affect the concentration of Zn and other metals (Benoit *et al.* 1994; Wen *et al.* 1999; Richards *et al.* 2018). Duan *et al.* (2019) found that the dynamics of Cu and Zn were primarily controlled by Fe and Mn diagenesis, and that sediment fluxes were a significant source of trace metals in the water column.

### Changes in pH in estuaries


pH measurements in estuaries were first made in the 1970s using the conventional hydrogen electrode (Mook and Koene 1975; Morris 1978). Subsequent studies found that there were serious difficulties in the accurate and reproducible measurement using this method because of the need to calibrate the electrode for different ionic strengths (Millero 1986), as well as variation in electrode responses with time and between different electrodes (Whitfield *et al.* 1985). Cai and Wang (1998) measured pH in several US estuaries but had only limited success with

calibration of the pH electrodes using a series of Tris buffers of different salinities. Despite these earlier findings, glass electrode measurements have continued to be made in estuaries, but some results may be questionable because ionic strength-specific calibration has often not been reported.

Increasingly spectrophotometric methods are being used for pH measurement in seawater and were first applied to estuaries by Mosley *et al.* (2004). In these methods, a sulfonephthalein indicator dye (e.g. *m*-cresol purple or thymol blue) is added to the sample and, based on knowledge of the dye dissociation constant (salinity and temperature dependent) and molar absorptivity, pH can be calculated following spectrophotometric measurement of the absorption maxima of the acid and base forms of the dye. Recently the *m*-cresol purple indicator dye characterisation has been reassessed to enable accurate spectrophotometric pH measurements for the entire estuarine salinity range (Douglas and Byrne 2017).

Based on spectrophotometric pH measurements in three New Zealand estuaries, pH was observed to decrease slightly at low salinities (salinity <2), increase rapidly at intermediate salinities (salinity 2–15) and then stabilise at higher salinities (salinity >15; Mosley *et al.* 2010). Similar trends in estuarine pH measured using glass electrodes have been reported previously, including slight decreases in pH at low salinities (Mook and Koene 1975; Morris 1978; Cai and Wang 1998; Guo *et al.* 2008). Using measurements and modelling, Mosley *et al.* (2010) explained the drivers of the general pH patterns during mixing of a circumneutral river with seawater as follows: (1) bicarbonate is the dominant species in circumneutral river water and at all salinities in estuaries, and exhibits a generally linear increase in concentration with salinity; (2) due to the large increases in ionic strength at low salinities (<5), the inorganic carbon dissociation constants change markedly, which can result in a decline in pH in this zone; and (3) following the initial pH decline, pH increases rapidly and then stabilises as salinities rise above 15 and the relative effects of changes in ionic strength on the inorganic carbon system equilibrium constants becomes smaller.

pH can vary widely in rivers, ranging from high pH ( $\geq 8$ ) in calcium carbonate-dominated systems (e.g. Edwards 1973) to low pH ( $\leq 4$ ) in organic or mineral acid-affected systems (e.g. Raymond and Oh 2009). The pH and inorganic carbon concentrations in rivers can also vary seasonally and annually depending on discharge patterns and longer-term climatic and anthropogenic drivers (Edwards 1973; Raymond and Oh 2009; Biswas and Mosley 2019). Given the uncertainty in pH changes in many estuaries due to the complications noted above, we used the end-member mixing model approach validated by Mosley *et al.* (2010) to explore how pH may vary with four different river water input compositions (Fig. 2): Case 1, average river water alkalinity (500  $\mu\text{mol kg}^{-1}$ ) with near atmospheric  $p\text{CO}_2$  (400 ppm); Case 2, average river water alkalinity with elevated  $p\text{CO}_2$  (4000 ppm); Case 3, higher alkalinity river water (2000  $\mu\text{mol kg}^{-1}$ ) and near atmospheric  $p\text{CO}_2$ ; and Case 4, higher alkalinity river water with elevated  $p\text{CO}_2$ . The salinity of the freshwater and seawater end-members in the model was 0.05 and 35 respectively. The model assumed equilibrium and used the salinity, alkalinity and  $p\text{CO}_2$  values in the four cases to calculate the total dissolved inorganic carbon (DIC) in the



**Fig. 2.** Modelled pH changes in estuaries based on end-member mixing of river water and seawater (for methods, see Mosley *et al.* 2010). Four cases were modelled with the seawater composition the same in all cases (based on global average seawater; alkalinity =  $2300 \mu\text{mol kg}^{-1}$ ,  $p\text{CO}_2 = 400 \text{ ppm}$ ). For the river water end-member, the four modelled cases were as follows: Case 1: alkalinity =  $500 \mu\text{mol kg}^{-1}$ ,  $p\text{CO}_2 = 400 \text{ ppm}$ ; Case 2: alkalinity =  $500 \mu\text{mol kg}^{-1}$ ,  $p\text{CO}_2 = 4000 \text{ ppm}$ ; Case 3: alkalinity =  $2000 \mu\text{mol kg}^{-1}$ ,  $p\text{CO}_2 = 400 \text{ ppm}$ ; and Case 4: alkalinity =  $2000 \mu\text{mol kg}^{-1}$ ,  $p\text{CO}_2 = 4000 \text{ ppm}$ .

end-members. The alkalinity and DIC were then mixed conservatively between the end-members in the model and the pH calculated for different salinities between 0.05 and 35.

All cases showed a slight decrease in pH at low salinities before being relatively stable at values  $>15$  (Fig. 2), which was generally consistent with the field and modelling studies described above. The average alkalinity and elevated  $p\text{CO}_2$  river water model (Case 2) exhibited the largest changes in pH throughout the estuarine salinity range. This case is likely most representative of a large number of rivers globally that have elevated  $p\text{CO}_2$  due to respiration exceeding net primary production (Cole and Caraco 2001). The higher alkalinity with atmospheric  $p\text{CO}_2$  model (Case 3) showed only a decrease and stabilisation rather than an increase in pH at salinities between 3 and 15, as in the other cases. This is likely because the alkalinity is similar to seawater, so the main pH changes are due to variation in inorganic carbon system equilibrium constants at low salinity. It should be noted that the assumption made in this model was that that dissolved organic carbon (DOC) and alkalinity behaved conservatively between the end-members. This is likely to be a valid assumption for estuaries with short residence times, but additions of DIC and alkalinity have been observed in estuaries that have been suggested to result from air–sea exchange of  $\text{CO}_2$  (for DIC because alkalinity is unaffected by this) and processes occurring in the underlying sediments such as denitrification, sulfate reduction, carbonate dissolution and the diffusion of organic acids (Cai and Wang 1998; Cai *et al.* 1998; Mosley *et al.* 2010; Shen *et al.* 2019). DIC isotope measurements have been found to be useful in tracing internal alkalinity sources (Atekwana *et al.* 2003), with calcium and strontium isotopes also useful for tracing carbonate precipitation and dissolution (Shao *et al.*


2018). Eutrophication, anoxia and elevated  $p\text{CO}_2$  due to climate change have been implicated in observations of increased acidification in some estuaries (Cai *et al.* 2017; Shen *et al.* 2019), as has resuspension of volatile sulfide phases (Richards *et al.* 2018). Spatial and temporal changes in pH in estuaries could greatly affect the sorption of trace metals and ligands to particle surfaces (Dzombak and Morel 1990), which is assessed further below.

### Insights and integration using geochemical modelling

A geochemical model was developed in the Visual MINTEQ program (G. P. Gustafsson, see <https://vminteq.lwr.kth.se>, accessed 22 August 2019) to help integrate and understand the key processes affecting the estuarine behaviour of trace metals (Cu, Zn and Pb in this case). The model included competitive metal binding to dissolved inorganic ligands, hydrous ferric oxide (HFO) particle surfaces (Dzombak and Morel 1990) and DOM by the Non-Ideal Competitive Adsorption (NICA)–Donnan model (Kinniburgh *et al.* 1999; Mosley *et al.* 2015). The pH variation in the initial model was based on the average alkalinity, high  $p\text{CO}_2$  scenario above (Case 2; Fig. 2). Metal concentrations in the river water end-member were Zn  $1 \mu\text{M}$ , Cu  $1 \mu\text{M}$  and Pb  $0.1 \mu\text{M}$ , compared with seawater end-member concentrations of 1, 1 and  $0.1 \text{ nM}$  for Zn, Cu and Pb respectively. The HFO concentration was maintained at  $0.1 \text{ g L}^{-1}$  throughout the estuarine salinity range. Further details of the model setup and input parameters are described in the ‘Modelling Description and Input File Parameters’ section of the Supplementary material.

The model results for Zn, Cu and Pb are shown in Fig. 3. More Pb and Zn is predicted to be bound to the Fe oxide phase than Cu, which is generally consistent with previous measurements of metal partitioning (see Table 1 and the above discussion). For Zn, the model results show an addition of dissolved Zn above the conservative mixing line at low to moderate salinities (0–10; Fig. 3). This can be explained by desorption from HFO surfaces due to decreasing pH (Fig. 2), increased competitive binding of Ca and Mg to these surfaces and to DOM for Mg (see Fig. S1). The model predictions of desorption of Zn at low salinities appear consistent with those found in many previous studies (Holliday and Liss 1976; Ackroyd *et al.* 1986; Yan *et al.* 1991; Chiffoleau *et al.* 1994). With regard to the pH-driven desorption, the sorption isotherm for Zn binding to HFO is in the range of pH 7–8, which is in the range of pH changes, at least in the seaward part, of most estuaries (Dzombak and Morel 1990). However, the desorbed Zn may become readsorbed as pH increases towards mid to high salinities according to the model results (Fig. 3). Standing *et al.* (2002) found, by isotope labelling experiments, that Zn–sediment interactions could be broadly defined as a two-step process, namely a short period of relatively fast surface sorption followed by a longer period of redistribution between different binding sites within the sediment until equilibrium is attained. Sediment characteristics, such as cation exchange capacity and the amount of organic material present, were proposed to affect these interactions, as would the prior distribution of Zn in suspended sediment in river water entering an estuary.

With regard to cation interactions, the model results show Mg binding to DOM increasing at low salinities, whereas Ca



**Fig. 3.** Geochemical model results exploring competitive metal binding for (a) Zn, (b) Cu and (c) Pb to dissolved inorganic ligands, hydrous ferric oxide particle surfaces and dissolved organic matter (DOM) in a typical estuary. The pH variation in the base case model was based on the average alkalinity, high  $p\text{CO}_2$  scenario (Case 2, Fig. 2).


binding decreases. This suggests Mg is competitively replacing Ca initially, and also likely Cu, from DOM. The model suggests released Cu then binds to HFO (Fig. 3). Ksionzek *et al.* (2018) measured a reduction in Cu–DOM binding with increasing salinity, which is consistent with our modelling results. Hamilton-Taylor *et al.* (2002) also predicted that Cu–humic binding decreased substantially with increasing ionic strength under typical estuarine conditions, but they did not postulate Ca and Mg binding as a significant factor. Waeles *et al.* (2005) found desorption of Cu occurred at low salinities for a large part of the year in the Penzé Estuary. Chiffoleau *et al.* (1994) also

found additions of Cu to the dissolved phase during estuarine mixing, but suggested this was desorption from particles, whereas van den Berg *et al.* (1987) suggested Cu and Zn will remain bound to the organic ligands throughout the salinity range of an estuary. The NICA–Donnan model suggests metal binding to DOM will change in response to increasing divalent cation concentrations (Kinniburgh *et al.* 1999). More integrated laboratory and field experimental research and modelling on competitive binding interactions would be useful to try and better determine the nature of Cu and Zn complexation during estuarine mixing. This should be coupled with spectrophotometric pH measurements in order to accurately quantify the pH changes. In contrast with Cu and Zn, Pb was predicted to remain strongly bound to HFO in the river water and throughout the estuarine salinity range (Fig. 3), which is generally consistent with field observations (Table 1).

To help explore how different river–estuarine water compositions may affect metal behaviour, the HFO, DOC and pH were varied in the geochemical model (Fig. 4). As the HFO increased, more Zn and Cu became bound to this phase, with the dissolved inorganic Zn and dissolved organic Cu fractions decreasing respectively. A similar general pattern was seen for Pb, but Pb was predominantly bound to HFO, except at the lowest concentration assessed ( $1 \text{ mg L}^{-1}$ ). Varying the DOC had little effect on Zn, but Cu binding to the dissolved organic fraction increased greatly as DOC increased. In the NICA–Donnan model, Cu is predicted to be a stronger competitor for organic binding sites than Zn and Pb when at equimolar concentrations. For Pb, the proportion bound to DOM increased only at very high DOC concentrations ( $>100 \text{ mg L}^{-1}$ ), with it remaining strongly bound to the HFO phase. Decreasing the pH markedly reduced Zn binding to HFO, with little effect on Cu and Pb binding. Again, this is consistent with pH decreases in the early stages of estuarine mixing. Taken as a whole, these model sensitivity results help explain why metal behaviour could vary between different estuaries (e.g. particle- or Fe-rich v. dissolved organic-rich riverine inputs), as well as at different temporal and spatial scales in the same estuary. Over the longer term, hydrological alterations to rivers or climate change may also alter the flux of metals, pH and other key geochemical processes (e.g. anoxia) affecting metal behaviour in estuaries.

## Conclusion

The results integrate and provide insight into the complex geochemical processes affecting metals in estuaries. The behaviour of a particular trace metal was found to be at least partially dependent on its partitioning in the river water end-member. The metals found in the colloidal fraction (e.g. Fe and Pb) in the river water end-member were removed non-conservatively during the early stages of estuarine mixing as a result of the aggregation of colloidal material. Modelling indicated Pb would remain bound to iron oxide surfaces during the large ionic strength increases and minor pH decreases during estuarine mixing. Hence, metals strongly bound to particles such as Pb should be rapidly deposited to the sediments and their distributions dominated by particle-phase dynamics. In contrast, other metals, such as Cu and Zn, that are less strongly bound to particle surfaces or predominantly associated with colloidal and DOM can behave quite differently, with variable behaviour



**Fig. 4.** Geochemical model results exploring the effects of varying (a) hydrous ferric oxide (HFO) concentrations (1–1000 mg L<sup>-1</sup>), (b) dissolved organic carbon (DOC) and (c) pH (6.5–7.8). The total metal concentration in the simulation was 0.9  $\mu$ M for Cu and Zn and 0.09  $\mu$ M for Pb. Salinity was 3 with the pH fixed at 6.88 in the varying HFO and DOC simulations (as per Case 2 results in Fig. 2 and 3).

between different estuaries and even seasonally within the same estuary. During the early stages of estuarine mixing, pH decreases occur and can result in, at least temporarily, desorption of metals such as Zn for particle surfaces. In contrast, Cu may remain strongly bound to dissolved organic ligands, although cation exchange may potentially result in less binding on DOM. Based on these findings, the distribution of metals such as Cu and Zn is observed and predicted to be much more complex and variable in estuaries, influenced dynamically by ionic strength, pH, sediment and metal oxide concentration, as well as the amount and nature of organic ligands. Integrated studies linking metal partitioning and speciation, particle aggregation and transport dynamics, biological uptake, sediment flux and pH changes would be beneficial. An increasing

use of state-of-the-art isotope tracing and DOM and metal speciation techniques, as well as geochemical models incorporating multi-assemblage metal binding mechanisms, should help improve our understanding of the drivers of variable metal behaviour in different estuaries.

### Conflicts of interest

The authors declare that they have no conflicts of interest.

### Declaration of funding

This research did not receive any specific funding.

### Acknowledgments

This review is dedicated to the memory of the late Professor Keith Hunter. Keith, together with his students and collaborators, made several seminal contributions to estuarine chemistry that are incorporated in this review. This includes his early research as a Ph.D. student that organic matter coatings on particles in natural waters are ubiquitous and determine their surface properties and stability, his research on the kinetics of Fe colloid aggregation and metal behaviour in estuaries using laboratory and field measurements, the use of atomic force microscopy for the first direct measurement of inter-particle forces in seawater and estuarine water, and the development of new methods for the measurement and modelling of pH in estuaries. Keith was a passionate and highly talented scientist who was widely respected internationally. He was a strong advocate for the environment, in particular the effects of climate change on aquatic systems. He is sorely missed but leaves an enduring influence in his high-quality body of published works and the hearts and mind of the people who were fortunate enough to have worked with him.

### References

- Ackroyd, D. R., Bale, A. J., Howland, R. J. M., Knox, S., Millward, G. E., and Morris, A. W. (1986). Distributions and behaviour of dissolved Cu, Zn and Mn in the Tamar estuary. *Estuarine, Coastal and Shelf Science* **23**, 621–640. doi:10.1016/0272-7714(86)90103-4
- Atekwana, E. A., Tedesco, L. P., and Jackson, L. R. (2003). Dissolved inorganic carbon (DIC) and hydrologic mixing in a subtropical riverine estuary, southwest Florida, USA. *Estuaries* **26**, 1391–1400. doi:10.1007/BF02803648
- Benoit, G. (1995). Evidence of the particle concentration effect for lead and other metals in freshwaters based on ultraclean technique analysis. *Geochimica et Cosmochimica Acta* **59**, 2677–2687. doi:10.1016/0016-7037(95)00164-U
- Benoit, G., Oktay-Marshall, S. D., Cantu, A., Hood, E. M., Coleman, C. H., Corapcioglu, M. O., and Santschi, P. H. (1994). Partitioning of Cu, Pb, Ag, Zn, Al, and Mn between filter retained particles, colloids, and solution in six Texas estuaries. *Marine Chemistry* **45**, 307–336. doi:10.1016/0304-4203(94)90076-0
- Biswas, T. K., and Mosley, L. M. (2019). From mountain ranges to sweeping plains, in droughts and flooding rains; River Murray water quality over the last four decades. *Water Resources Management* **33**, 1087–1101. doi:10.1007/S11269-018-2168-1
- Bourg, A. C. M. (1987). Trace metal adsorption modelling and particle-water interactions in estuarine environments. *Continental Shelf Research* **7**, 1319–1332. doi:10.1016/0278-4343(87)90036-7
- Boyle, E., Collier, R., Dengler, A. T., Edmond, J. M., Ng, A. C., and Stallard, R. F. (1974). On the chemical mass-balance in estuaries. *Geochimica et Cosmochimica Acta* **38**, 1719–1728. doi:10.1016/0016-7037(74)90188-4
- Boyle, E. A., Edmond, J. M., and Sholkovitz, E. R. (1977). The mechanism of iron removal in estuaries. *Geochimica et Cosmochimica Acta* **41**, 1313–1324. doi:10.1016/0016-7037(77)90075-8



- Boyle, E. A., Husted, S. S., and Grant, B. (1982). The chemical mass balance of the Amazon plume – II. Copper, nickel, and cadmium. *Deep-Sea Research* **29**, 1355–1364. doi:10.1016/0198-0149(82)90013-9
- Bruland, K. W. (1989). Complexation of zinc by natural organic ligands in the central North Pacific. *Limnology and Oceanography* **34**, 269–285. doi:10.4319/LO.1989.34.2.0269
- Burton, J. D., and Liss, P. S. (1976). 'Estuarine Chemistry.' (Academic Press: London, UK.)
- Cai, W.-J., and Wang, Y. (1998). The chemistry, fluxes, and sources of CO<sub>2</sub> in the estuarine waters of the Satilla and Altamaha rivers, Georgia. *Limnology and Oceanography* **43**, 657–668. doi:10.4319/LO.1998.43.4.0657
- Cai, W.-J., Wang, Y., and Hodson, R. E. (1998). Acid-base properties of dissolved organic matter in the estuarine waters of Georgia, USA. *Geochimica et Cosmochimica Acta* **62**, 473–483. doi:10.1016/S0016-7037(97)00363-3
- Cai, W. J., Huang, W. J., Luther, G. W., Pierrot, D., Li, M., Testa, J., Xue, M., Joesoef, A., Mann, R., Brodeur, J., Xu, Y. Y., Chen, B., Hussain, N., Waldbusser, G. G., Cornwell, J., and Kemp, W. M. (2017). Redox reactions and weak buffering capacity lead to acidification in the Chesapeake Bay. *Nature Communications* **8**, 369. doi:10.1038/S41467-017-00417-7
- Chiffolleau, J.-F., Cossa, D., Auger, D., and Truquet, I. (1994). Trace metal distribution, partition and fluxes in the Seine estuary (France) in low discharge regime. *Marine Chemistry* **47**, 145–158. doi:10.1016/0304-4203(94)90105-8
- Church, T. M. (1986). Biogeochemical factors influencing the residence time of microconstituents in a large tidal estuary, Delaware Bay. *Marine Chemistry* **18**, 393–406. doi:10.1016/0304-4203(86)90020-4
- Cole, J. J., and Caraco, N. F. (2001). Carbon in catchments: connecting terrestrial carbon losses with aquatic metabolism. *Marine and Freshwater Research* **52**, 101–110. doi:10.1071/MF00084
- Dai, M.-H., and Martin, J.-M. (1995). First data on trace metal level and behaviour in two major Arctic river–estuarine systems (Ob and Yenisey) and in the adjacent Kara Sea, Russia. *Earth and Planetary Science Letters* **131**, 127–141. doi:10.1016/0012-821X(95)00021-4
- Dai, M., Martin, J.-M., and Cauwet, G. (1995). The significant role of colloids in the transport and transformation of organic carbon and associated trace metals (Cd, Cu and Ni) in the Rhône delta (France). *Marine Chemistry* **51**, 159–175. doi:10.1016/0304-4203(95)00051-R
- Douglas, N. K., and Byrne, R. H. (2017). Spectrophotometric pH measurements from river to sea: Calibration of mCP for 0 ≤ S ≤ 40 and 278.15 ≤ T ≤ 308.15K. *Marine Chemistry* **197**, 64–69. doi:10.1016/J.MARCHEM.2017.10.001
- Duan, L., Song, J., Liang, X., Yin, M., Yuan, H., Li, X., Ren, C., Zhou, B., Kang, X., and Yin, X. (2019). Dynamics and diagenesis of trace metals in sediments of the Changjiang Estuary. *The Science of the Total Environment* **675**, 247–259. doi:10.1016/J.SCITOTENV.2019.04.190
- Dzombak, D. A., and Morel, F. M. M. (1990). 'Surface Complexation Modeling Hydrous Ferric Oxide.' (Wiley: New York, NY, USA.)
- Eckert, J. M., and Sholkovitz, E. R. (1976). The flocculation of iron, aluminium and humates from river water by electrolytes. *Geochimica et Cosmochimica Acta* **40**, 847–848. doi:10.1016/0016-7037(76)90036-3
- Edwards, A. M. C. (1973). Dissolved load and tentative solute budgets of some Norfolk catchments. *Journal of Hydrology* **18**, 201–217. doi:10.1016/0022-1694(73)90048-6
- Erel, Y., Morgan, J. J., and Paterson, C. C. (1991). Natural levels of lead and cadmium in a remote mountain stream. *Geochimica et Cosmochimica Acta* **55**, 707–719. doi:10.1016/0016-7037(91)90335-3
- Fu, J., Tang, X.-L., Zhang, J., and Balzer, W. (2013). Estuarine modification of dissolved and particulate trace metals in major rivers of East-Hainan, China. *Continental Shelf Research* **57**, 59–72. doi:10.1016/J.CSR.2012.06.015
- Furukawa, Y., Reed, A. H., and Zhang, G. (2014). Effect of organic matter on estuarine flocculation: a laboratory study using montmorillonite, humic acid, xanthan gum, guar gum and natural estuarine flocs. *Geochemical Transactions* **15**, 1. doi:10.1186/1467-4866-15-1
- Gigault, J., Balaresque, M., and Tabuteau, H. (2018). Estuary-on-a-chip: unexpected results for the fate and transport of nanoparticles. *Environmental Science. Nano* **5**, 1231–1236. doi:10.1039/C8EN00184G
- Greenamoyer, J. M., and Moran, S. B. (1997). Investigation of Cd, Cu, Ni and <sup>234</sup>Th in the colloidal size range in the Gulf of Maine. *Marine Chemistry* **57**, 217–226. doi:10.1016/S0304-4203(97)00015-7
- Guieu, C., and Martin, J. M. (2002). The level and fate of metals in the Danube River plume. *Estuarine, Coastal and Shelf Science* **54**, 501–512. doi:10.1006/ECSS.2000.0660
- Guieu, C., Huang, W. W., Martin, J.-M., and Yong, Y. Y. (1996). Outflow of trace metals into the Laptev Sea by the Lena River. *Marine Chemistry* **53**, 255–267. doi:10.1016/0304-4203(95)00093-3
- Guo, X., Cai, W.-J., Zhai, W., Dai, M., Wang, Y., and Chen, B. (2008). Seasonal variations in the inorganic carbon system in the Pearl River (Zhujiang) estuary. *Continental Shelf Research* **28**, 1424–1434. doi:10.1016/J.CSR.2007.07.011
- Hamilton-Taylor, J., Postill, A. S., Tipping, E., and Harper, M. P. (2002). Laboratory measurements and modeling of metal–humic interactions under estuarine conditions. *Geochimica et Cosmochimica Acta* **66**, 403–415. doi:10.1016/S0016-7037(01)00777-3
- Holliday, L. M., and Liss, P. S. (1976). The behaviour of dissolved iron, manganese and zinc in the Beaulieu Estuary, S. England. *Estuarine and Coastal Marine Science* **4**, 349–353. doi:10.1016/0302-3524(76)-90066-9
- Hoyle, J., Elderfield, H., Gledhill, A., and Greaves, M. (1984). The behaviour of the rare earth elements during mixing of river and sea waters. *Geochimica et Cosmochimica Acta* **48**, 143–149. doi:10.1016/0016-7037(84)90356-9
- Hunter, K. A. (1980). Microelectrophoretic properties of natural surface-active organic matter in coastal seawater. *Limnology and Oceanography* **25**, 807–822. doi:10.4319/LO.1980.25.5.0807
- Hunter, K. A. (1983). On the estuarine mixing of dissolved substances in relation to colloid stability and surface properties. *Geochimica et Cosmochimica Acta* **47**, 467–473. doi:10.1016/0016-7037(83)90269-7
- Hunter, K. A., and Leonard, M. W. (1988). Colloid stability and aggregation in estuaries: 1. Aggregation kinetics of riverine dissolved iron after mixing with seawater. *Geochimica et Cosmochimica Acta* **52**, 1123–1130. doi:10.1016/0016-7037(88)90266-9
- Hunter, K. A., and Liss, P. S. (1979). The surface charge of suspended particles in estuarine and coastal waters. *Nature* **282**, 823–825. doi:10.1038/282823A0
- Hunter, K. A., and Liss, P. S. (1982). Organic matter and the surface charge of suspended particles in estuarine waters. *Limnology and Oceanography* **27**, 322–335. doi:10.4319/LO.1982.27.2.0322
- Hunter, K. A., Leonard, M. R., Carpenter, P. D., and Smith, J. D. (1997). Aggregation of iron colloids in estuaries: a heterogeneous kinetics study using continuous mixing of river and sea waters. *Colloids and Surfaces – A. Physicochemical and Engineering Aspects* **120**, 111–121. doi:10.1016/S0927-7757(96)03719-3
- Illuminati, S., Annibaldi, A., Truzzi, C., Tercier-Waeber, M.-L., Nöl, S., Braungardt, C. B., Achterberg, E. P., Howell, K. A., Turner, D., Marini, M., Romagnoli, T., Totti, C., Confalonieri, F., Graziottin, F., Buffle, J., and Scarponi, G. (2019). *In situ* trace metal (Cd, Pb, Cu) speciation along the Po River plume (Northern Adriatic Sea) using submersible systems. *Marine Chemistry* **212**, 47–63. doi:10.1016/J.MARCHEM.2019.04.001
- Kinniburgh, D. G., van Riemsdijk, W. H., Koopal, L. K., Borkovec, M., Benedetti, M. F., and Avena, M. (1999). Ion binding to natural organic matter: competition, heterogeneity, stoichiometry and thermodynamic consistency. *Colloids and Surfaces – A. Physicochemical and Engineering Aspects* **151**, 147–166. doi:10.1016/S0927-7757(98)00637-2
- Kruger, B. R., Dalzell, B. J., and Austin-Minor, E. C. (2011). Effect of organic matter source and salinity on dissolved organic matter isolation

- via ultrafiltration and solid phase extraction. *Aquatic Sciences* **73**, 405–417. doi:10.1007/S00027-011-0189-4
- Ksionzek, K. B., Zhang, J., Ludwiczowski, K.-U., Wilhelms-Dick, D., Trimborn, S., Jendrosseck, T., Kattner, G., and Koch, B. P. (2018). Stoichiometry, polarity, and organometallics in solid-phase extracted dissolved organic matter of the Elbe–Weser estuary. *PLoS One* **13**(9), e0203260. doi:10.1371/JOURNAL.PONE.0203260
- Liss, P. S. (1976). Conservative and non-conservative behaviour of dissolved constituents during estuarine mixing. In 'Estuarine Chemistry'. (Eds J. D. Burton and P. S. Liss.) pp. 93–130. (Academic Press: London, UK.)
- Liu, Q., Jia, Z., Li, S., and Hu, J. (2019). Assessment of heavy metal pollution, distribution and quantitative source apportionment in surface sediments along a partially mixed estuary (Modaomen, China). *Chemosphere* **225**, 829–838. doi:10.1016/J.CHEMOSPHERE.2019.03.063
- Mantoura, R. F. C., Dickson, A., and Riley, J. (1978). The complexation of metals with humic materials in natural waters. *Estuarine and Coastal Marine Science* **6**, 387–408. doi:10.1016/0302-3524(78)90130-5
- Mari, X., Torr  ton, J.-P., Bich-Thuy Trinh, C., Bouvier, T., Van Thuoc, C., Lefebvre, J.-P., and Ouillon, S. (2012). Aggregation dynamics along a salinity gradient in the Bach Dang estuary, North Vietnam. *Estuarine, Coastal and Shelf Science* **96**, 151–158. doi:10.1016/J.ECSS.2011.10.028
- McLachlan, R. L., Ogston, A. S., and Allison, M. A. (2017). Implications of tidally varying bed stress and intermittent estuarine stratification on fine-sediment dynamics through the Mekong's tidal river to estuarine reach. *Continental Shelf Research* **147**, 27–37. doi:10.1016/J.CSR.2017.07.014
- Merschel, G., Bau, M., and Dantas, E. L. (2017). Contrasting impact of organic and inorganic nanoparticles and colloids on the behavior of particle-reactive elements in tropical estuaries: an experimental study. *Geochimica et Cosmochimica Acta* **197**, 1–13. doi:10.1016/J.GCA.2016.09.041
- Millero, F. J. (1986). The pH of estuarine waters. *Limnology and Oceanography* **31**, 839–847. doi:10.4319/LO.1986.31.4.0839
- Mook, W. G., and Koene, B. K. S. (1975). Chemistry of dissolved inorganic carbon in estuarine and coastal brackish waters. *Estuarine and Coastal Marine Science* **3**, 325–336. doi:10.1016/0302-3524(75)90032-8
- Morris, A. W. (1978). Chemical processes in estuaries: the importance of pH and its variability. In 'The Aquatic Environment'. (Ed. W. E. Krumbein.) pp. 179–187. (Ann Arbor Science: Ann Arbor, MI, USA.)
- Mosley, L. M., Hunter, K. A., and Ducker, W. A. (2003). Forces between particles in natural waters. *Environmental Science & Technology* **37**, 3303–3308. doi:10.1021/ES026216D
- Mosley, L., Hunter, K., and Husheer, S. (2004). Spectrophotometric pH measurement in estuaries using thymol blue and m-cresol purple. *Marine Chemistry* **91**, 175–186. doi:10.1016/J.MARCH.2004.06.008
- Mosley, L. M., Peake, B. M., and Hunter, K. A. (2010). Modelling of pH and inorganic carbon speciation in estuaries using the composition of the river and seawater end members. *Environmental Modelling & Software* **25**, 1658–1663. doi:10.1016/J.ENVSOFT.2010.06.014
- Mosley, L. M., Daly, R., Palmer, D., Yeates, P., Dallimore, C., Biswas, T., and Simpson, S. L. (2015). Predictive modelling of pH and dissolved metal concentrations and speciation following mixing of acid drainage with river water. *Applied Geochemistry* **59**, 1–10. doi:10.1016/J.APGEOCHEM.2015.03.006
- Mulholland, D. S., Poitras, F., Boaventura, G. R., Allard, T., Vieira, L. C., Santos, R. V., Mancini, L., and Seyler, P. (2015). Insights into iron sources and pathways in the Amazon River provided by isotopic and spectroscopic studies. *Geochimica et Cosmochimica Acta* **150**, 142–159. doi:10.1016/J.GCA.2014.12.004
- Muller, F. L. L. (2018). Exploring the potential role of terrestrially derived humic substances in the marine biogeochemistry of iron. *Frontiers of Earth Science* **6**, 159–169. doi:10.3389/FEART.2018.00159
- Newton, P. P., and Liss, P. S. (1987). Positively charged suspended particles: studies in an iron-rich river and its estuary. *Limnology and Oceanography* **32**, 1267–1276. doi:10.4319/LO.1987.32.6.1267
- Nowostawska, U., Kim, J. P., and Hunter, K. A. (2008). Aggregation of riverine colloidal iron in estuaries: a new kinetic study using stopped-flow mixing. *Marine Chemistry* **110**, 205–210. doi:10.1016/J.MARCH.2008.03.001
- Paulson, A. J., Curl, H. C. Jr., and Gendron, J. F. (1994). Partitioning of Cu in estuarine waters, 1. Partitioning in a poisoned system. *Marine Chemistry* **45**, 67–80. doi:10.1016/0304-4203(94)90092-2
- Pearson, H., Comber, S. D. W., Braungardt, C. B., Worsfold, P., Stockdale, A., and Locks, S. (2018). Determination and prediction of zinc speciation in estuaries. *Environmental Science & Technology* **52**(24), 14245–14255. doi:10.1021/ACS.EST.8B04372
- Powell, R. T. (1996). Colloidal trace metals, organic carbon and nitrogen in a southeastern U.S. estuary. *Marine Chemistry* **55**, 165–176. doi:10.1016/S0304-4203(96)00054-0
- Ram  rez-Mendoza, R., Souza, A. J., and Amoudry, L. O. (2014). Modeling flocculation in a hypertidal estuary. *Ocean Dynamics* **64**, 301–313. doi:10.1007/S10236-013-0675-4
- Raymond, P. A., and Oh, N. H. (2009). Long term changes of chemical weathering products in rivers heavily impacted from acid mine drainage: insights on the impact of coal mining on regional and global carbon and sulfur budgets. *Earth and Planetary Science Letters* **284**, 50–56. doi:10.1016/J.EPSL.2009.04.006
- Reese, A., Zimmermann, T., Pr  frock, D., and Irrgeher, J. (2019). Extreme spatial variation of Sr, Nd and Pb isotopic signatures and 48 element mass fractions in surface sediment of the Elbe River Estuary – suitable tracers for processes in dynamic environments? *The Science of the Total Environment* **668**, 512–523. doi:10.1016/J.SCI.TOTENV.2019.02.401
- Richards, C. M., van Puffelen, J. L., and Pallud, C. (2018). Effects of sediment resuspension on the oxidation of acid-volatile sulfides and release of metals (iron, manganese, zinc) in Pescadero Estuary (CA, USA). *Environmental Toxicology and Chemistry* **37**, 993–1006. doi:10.1002/ETC.4047
- Sander, S., Mosley, L., and Hunter, K. (2004). Investigations of inter-particle forces in natural waters using AFM: effects of adsorbed humic acids on iron oxide and alumina surface properties. *Environmental Science & Technology* **38**, 4791–4796. doi:10.1021/ES049602Z
- Sanders, C. J., Santos, I. R., Maher, D. T., Sadat-Noori, M., Schnetger, B., and Brumsack, H. J. (2015). Dissolved iron exports from an estuary surrounded by coastal wetlands: can small estuaries be a significant source of Fe to the ocean? *Marine Chemistry* **176**, 75–82. doi:10.1016/J.MARCH.2015.07.009
- Santschi, P. H., Lenhart, J. J., and Honeyman, B. D. (1997). Heterogeneous processes affecting trace contaminant distribution in estuaries: the role of natural organic matter. *Marine Chemistry* **58**, 99–125. doi:10.1016/S0304-4203(97)00029-7
- Shao, Y., Farkas, J., Holmden, C., Mosley, L., Kell-Duivesteyn, I., Izzo, C., Reis-Santos, P., Tyler, J., Torber, P., Fryda, J., Taylor, H., Haynes, D., Tibby, J., and Gillanders, B. M. (2018). Calcium and strontium isotope systematics in the lagoon estuarine environments of South Australia: implications for water source mixing, carbonate fluxes and fish migration. *Geochimica et Cosmochimica Acta* **239**, 90–108. doi:10.1016/J.GCA.2018.07.036
- Shen, C., Testa, J. M., Li, M., Cai, W.-J., Waldbusser, G. G., Ni, W., Kemp, W. M., Cornwell, J., Chen, B., Brodeur, J., and Su, J. (2019). Controls on carbonate system dynamics in a coastal plain estuary: a modelling study. *Journal of Geophysical Research – Biogeosciences* **124**, 61–78. doi:10.1029/2018JG004802
- Shiller, A. M., and Boyle, E. A. (1991). Trace elements in the Mississippi River Delta outflow region; behaviour at high discharge. *Geochimica et Cosmochimica Acta* **55**, 3241–3251. doi:10.1016/0016-7037(91)90486-O
- Sholkovitz, E. R. (1976). Flocculation of dissolved organic and inorganic matter during the mixing of river water and seawater. *Geochimica et Cosmochimica Acta* **40**, 831–845. doi:10.1016/0016-7037(76)90035-1

- Sholkovitz, E. R. (1978). The flocculation of dissolved Fe, Mn, Al, Cu, Ni, Co, and Cd during estuarine mixing. *Earth and Science Planetary Letters* **41**, 77–86. doi:10.1016/0012-821X(78)90043-2
- Sholkovitz, E. R. (1995). The aquatic chemistry of rare earth elements in rivers and estuaries. *Aquatic Geochemistry* **1**, 1–34. doi:10.1007/BF01025229
- Sholkovitz, E. R., Boyle, E. A., and Price, N. B. (1978). The removal of dissolved humic acids and iron during estuarine mixing. *Earth and Science Planetary Letters* **40**, 130–136. doi:10.1016/0012-821X(78)90082-1
- Standing, W. J. F., Oughton, D. H., and Salbu, B. (2002). Remobilisation of <sup>109</sup>Cd, <sup>65</sup>Zn and <sup>54</sup>Mn from freshwater-labelled river sediments when mixed with seawater. *Environment International* **28**, 185–195. doi:10.1016/S0160-4120(02)00028-4
- Stumm, W., and Morgan, J. (1996). 'Aquatic Chemistry: Chemical Equilibria and Rates in Natural Waters', 3rd edn. (Wiley Interscience: New York, NY, USA.)
- Sunda, W. G., and Hanson, P. J. (1979). Chemical speciation of copper in river water. In 'Chemical Modeling in Aqueous Systems'. (Ed. E.A. Jenne.) pp. 147–180. (American Chemical Society: Washington, DC, USA.)
- Tang, D., Warnken, K. W., and Santschi, P. H. (2002). Distribution and partitioning of trace metals (Cd, Cu, Ni, Pb, Zn) in Galveston Bay waters. *Marine Chemistry* **78**, 29–45. doi:10.1016/S0304-4203(02)00007-5
- Tanguy, V., Waeles, M., Gigault, J., Cabon, J.-Y., Quentel, F., and Riso, R. D. (2011). The removal of colloidal lead during estuarine mixing: seasonal variations and importance of iron oxides and humic substances. *Marine and Freshwater Research* **62**, 329–341. doi:10.1071/MF10220
- Tipping, E., Lofts, S., and Lawlor, A. (1998). Modelling the chemical speciation of trace metals in the surface waters of the Humber system. *The Science of the Total Environment* **210–211**, 63–77. doi:10.1016/S0048-9697(98)00045-X
- Trento, A., and Vinzon, S. (2014). Experimental modelling of flocculation processes – the case of Paraíba do Sul Estuary. *International Journal of Sediment Research* **29**, 378–390. doi:10.1016/S1001-6279(14)60052-4
- van den Berg, C. M. G., Merks, A. G. A., and Duursma, E. K. (1987). Organic complexation and its control of the dissolved concentrations of copper and zinc in the Scheldt Estuary. *Estuarine, Coastal and Shelf Science* **24**, 785–797. doi:10.1016/0272-7714(87)90152-1
- Verwey, E. J. W., and Overbeek, T. G. (1948). 'Theory of the Stability of Lyophobic Colloids.' (Elsevier: Amsterdam, Netherlands.)
- Waeles, M., Riso, R. D., and Corre, P. L. (2005). Seasonal variations of dissolved and particulate copper species in estuarine waters. *Estuarine, Coastal and Shelf Science* **62**, 313–323. doi:10.1016/J.ECSS.2004.09.019
- Waeles, M., Tanguy, V., Lespes, G., and Riso, R. D. (2008). Behaviour of colloidal trace metals (Cu, Pb and Cd) in estuarine waters: an approach using frontal ultrafiltration (UF) and stripping chronopotentiometric methods (SCP). *Estuarine, Coastal and Shelf Science* **80**, 538–544. doi:10.1016/J.ECSS.2008.09.010
- Wang, Y. P., Voulgaris, G., Li, Y., Yang, Y., Gao, J., Chen, J., and Gao, S. (2013). Sediment resuspension, flocculation, and settling in a macrotidal estuary. *Journal of Geophysical Research. Oceans* **118**, 5591–5608. doi:10.1002/JGRC.20340
- Wen, L.-S., Stordal, M. C., Tang, D., Gill, G. A., and Santschi, P. H. (1996). An ultraclean cross-flow ultrafiltration technique for the study of trace metal phase speciation in seawater. *Marine Chemistry* **55**, 129–152. doi:10.1016/S0304-4203(96)00052-7
- Wen, L.-S., Santschi, P., Gill, G., and Paternostro, C. (1999). Estuarine trace metal distributions in Galveston Bay: importance of colloidal forms in the speciation of the dissolved phase. *Marine Chemistry* **63**, 185–212. doi:10.1016/S0304-4203(98)00062-0
- Wetz, M. S., Robbins, M. C., and Paerl, H. W. (2009). Transparent exopolymer particles (TEP) in a river-dominated estuary: spatial-temporal distributions and an assessment of controls upon TEP formation. *Estuaries and Coasts* **32**, 447–455. doi:10.1007/S12237-009-9143-2
- Whitfield, M., Butler, R. A., and Covington, A. K. (1985). The determination of pH in estuarine waters I. Definition of pH scales and the selection of buffers. *Oceanologica Acta* **8**, 423–432.
- Windom, H., Wallace, G., Smith, R., Dudek, N., Maeda, M., Dulmage, R., and Storti, F. (1983). Behaviour of copper in southeastern United States estuaries. *Marine Chemistry* **12**, 183–193. doi:10.1016/0304-4203(83)90079-8
- Yan, L., Stallard, R. F., Key, R. M., and Crerar, D. A. (1991). Trace metals and dissolved organic carbon in estuaries and offshore waters of New Jersey, USA. *Geochimica et Cosmochimica Acta* **55**, 3647–3656. doi:10.1016/0016-7037(91)90062-A
- Zhou, J. L., Rowland, S., Fauzi, R., Mantoura, C., and Braven, J. (1994). The formation of humic coatings on mineral particles under simulated estuarine conditions – a mechanistic study. *Water Research* **28**, 571–579. doi:10.1016/0043-1354(94)90008-6
- Zhou, Z., Stolpe, B., Guo, L., and Shiller, A. M. (2016). Colloidal size spectra, composition and estuarine mixing behavior of DOM in river and estuarine waters of the northern Gulf of Mexico. *Geochimica et Cosmochimica Acta* **181**, 1–17. doi:10.1016/J.GCA.2016.02.032

Handling Editor: Michael Ellwood