

flew against the chart table on the bridge, but escaped apparently uninjured.

## REFERENCES

- Alexander, W. B., 1928. *Birds of the Ocean*. Putnam.  
 Arnold, L. W., 1948. 'Observations in populations of North Pacific pelagic birds.' *Auk*, 65, 553-558.  
 Austin, O. L., 1949. 'The Status of Steller's Albatross.' *Pacific Science*, 3, 283-295.  
 Baker, R. H., 1951. 'The Avifauna of Micronesia.' *Univ. Kansas Pubs. Nat. Hist.* 3, 1-359.  
 Cleland, J. B., 1938. 'Birds noted between Adelaide and England via Suez, and on Return across the Atlantic and Pacific Oceans.' *South Aust. Orn.*, 14, 207-217.  
 Dixon, C. C., 1933. 'Some Observations on the Albatrosses and other birds of the Southern Oceans.' *Trans. Canad. Inst. Toronto*, 19, 117-139.  
 Falla, R. A., 1942. 'Review of the Smaller Pacific Forms of *Pterodroma* and *Cookilaria*.' *Emu*, 42, 111-118.  
 Fleming, C. A., 1950. 'Some South Pacific Sea-bird Logs.' *Emu*, 49, 169-188.  
 Jespersen, P., 1930. 'Distribution of some oceanic birds in the waters east of New Zealand.' *Nature*, London, 126, 280-282.  
 Jespersen, P., 1933. 'Observations on the oceanic birds of the Pacific and adjoining waters.' *Vidensk. Medd. naturh. Foren.*, Copenhagen, 94, 187-221.  
 Kenyon, K. W., 1950. 'Distribution of Albatrosses in the North Pacific and adjacent waters.' *Condor*, 52, 97-103.  
 Macdonald, J. D., 1952. 'A recent sight record of Steller's Albatross.' *Ibis*, 94, 536.  
 Murphy, R. C., 1936. *Oceanic Birds of South America*. Amer. Mus. Nat. Hist., N.Y., 2 vols.  
 Oliver, W. R. B., 1930. *Birds of New Zealand*.  
 Serventy, D. L., 1939. 'Migration records at sea.' *Condor*, 41, 257-258.  
 Starrett, W. C., and Dixon, K. L., 1946. 'The scarcity of the Black-footed Albatross in parts of its known range.' *Condor*, 48, 268-271.  
 Wynne-Edwards, V. C., 1935. 'On the habits and distribution of birds in the North Atlantic.' *Proc. Bost. Soc. Nat. Hist.*, 40, 233-346.

**White-throated Warbler in Western Victoria.**—The occurrence of this species in this area would appear to be sufficiently rare to be worth recording. On January 12, 1954, at 7 a.m., I was attracted by its notes in my garden where it was singing at about one-minute intervals from thick lilly pilly (*Eugenia*) trees. From there it flew into the lower branches of a eucalypt where I was able to observe it at only ten feet distance. It was heard and seen at intervals that day in the same vicinity, and again at 7 a.m. next morning when I was able to get immediately below it, about six feet above my head. It appeared to be a solitary bird, and was not seen or heard after about 7.30 a.m. on the second day. Its yellow under parts appeared paler than those of the Queensland skins which I have examined at the National Museum in Melbourne.—CLAUDE N. AUSTIN, Coleraine, Vic., 14/2/54.