son, National Museum of Victoria, who identified the barnacle and molluses referred to herein.

REFERENCES

Baker, A. A. 1956. The swallowing of stones by animals. Vic. Nat. 73: 82-95.
Bierman, W. H., and Voous, K. H. 1950. Birds observed and collected during the whaling expeditions of the "Willem Barendz" in the Antarctic, 1946-47 and 1947-48. Ardea. 37: 19.
Budd, G. M. 1964. The ANARE 1963 Expedition to Heard Island. ANARE

Reps. A(1) Pubn. No. 74.

Campbell, A. J. 1900. The Nests and Eggs of Australian Birds, pp. 885-892, Sheffield, England.

Lewis, F. 1946. Feeding habits of the Short-tailed Shearwater (Mutton-Bird).
 Emu. 45: 225-228.
 Mawson, D. 1943. Macquarie Island, its Geography and Geology. Australas.

Antarct. Exp. Sci. Rep. (A)5.
Sutherland, F. L. 1964. Pumice from Tasmania. Aust. J. Sci. 26: 397.

Egg removal by Horsfield Bronze-Cuckoo.—On October 28, 1964, at about 1 p.m. I was strolling on Mt. Pleasant, near Canberra, A.C.T., in company with Mr S. J. Wilson when we saw a Horsfield Bronze-Cuckoo, Chalcites basalis in the vicinity of an already-known nest of the Western Warbler, Gerygone fusca. The nest was built in a eucalypt sapling at 15 to 20 feet above the ground. While we watched through binoculars from a distance of about 25 feet, the cuckoo made its way to the nest, paused at the entrance, then flew away carrying a pink egg in its bill. Neither the Western Warblers, nor any other cuckoo, were seen in the vicinity.-IAN GRANT, 12 Selwyn Street, Hackett, A.C.T.