

Supplementary Material

Short- and long-term outcomes of the absence of protein during bovine blastocyst formation *in vitro*

A. Murillo-Ríos^A, V. Mailló^B, M. Muñoz^A, A. Gutiérrez-Adán^B, S. Carrocera^A, D. Martín-González^A, A. Fernández-Buznego^A and E. Gómez^{A,C}

^AGenética y Reproducción Animal, Centro de Biotecnología Animal, SERIDA, Camino de Rioseco 1225, 33394 Gijón, Spain.

^BDepartamento de Reproducción Animal, INIA, Ctra de la Coruña, km 5.9, 2804 Madrid, Spain.

^CCorresponding author. Email: egomez@serida.org

Table S1. Embryo transfers (N) performed per bull, with expression of Day-40 pregnancies and miscarriage attained

Bull	Sex Sorted	Protein (+)			Protein (-)		
		ET(N)	Day-40	Miscarriage	ET(N)	Day-40	Miscarriage
A	No	5	2	1	9	6	0
A	No				1	1	in course
B	No	2	1	1	6	4	1
C	Female	7	4	1			
C	Male	3	2	1			
F	No	2	1	in course	8	4	in course
G	Female	2	1	1			
H	Female	1	0		1	1	0
H	Female				6	3	in course
I	Female	5	3	1	4	2	0
I	Male	4	1	0			
J	No	10	4	0			
K	Male	3	2	0	2	0	
K	Female	4	1	0			
L	No	2	1	in course	1	1	in course
Total		50	23	6	38	22	1