

10.1071/RD16460_AC

© CSIRO 2017

Supplementary Material: *Reproduction, Fertility and Development*, 2017, 29(12), 2437–2445.

Supplementary Material

Regulation of interleukin-11 expression in ovulatory follicles of the rat ovary

You-Jee Jang^A, Jae-Il Park^B, Seong-Eun Jeong^A, You-Mi Seo^D, Phuong T. M. Dam^A, Young-Woo Seo^B, Bum-Chae Choi^E, Sang-Jin Song^E, Sang-Young Chun^{A,F} and Moon-Kyoung Cho^{C,F}

^ASchool of Biological Sciences and Technology, Faculty of Life Science, Chonnam National University, Gwangju 61186, Republic of Korea.

^BAnimal Facility of Aging Science, Korea Basic Science Institute, Gwangju 61186, Republic of Korea.

^CDepartment of Obstetrics and Gynecology, Chonnam National University Medical School, Gwangju 61469, Republic of Korea.

^DDepartment of Oral Histology–Developmental Biology, School of Dentistry and Dental Research Institute, Seoul National University, Seoul 03080, Republic of Korea.

^ECenter for Recurrent Miscarriage and Infertility, Creation and Love Women's Hospital, Gwangju 61917, Republic of Korea.

^FCorresponding authors. Emails: chomk@jnu.ac.kr; sychun@jnu.ac.kr




Fig. S1. The specific marker genes, Rhox8 (a), Cyp17 α 1 (b) and CD45 (c) for granulosa cells (GCs), theca cells (TCs) and leukocytes, respectively, were analyzed in the ovary and isolated cells of dissected preovulatory follicles by a real-time PCR.