

Supplementary Material

Morphometric and gene expression analyses of stromal expansion during development of the bovine fetal ovary

M. D. Hartanti^A, K. Hummitzsch^A, H. F. Irving-Rodgers^{A,B}, W. M. Bonner^A, K. J. Copping^A, R. A. Anderson^C, I. C. McMillen^D, V. E. A. Perry^E and R. J. Rodgers^{A,F}

^ADiscipline of Obstetrics and Gynaecology, School of Medicine, Robinson Research Institute, The University of Adelaide, Adelaide, SA 5005, Australia.

^BSchool of Medical Science, Griffith University, Gold Coast Campus, Qld 4222, Australia.

^CMedical Research Council Centre for Reproductive Health, University of Edinburgh, Edinburgh, EH16 4TJ, UK.

^DThe Chancellery, University of Newcastle, Callaghan, NSW 2308, Australia.

^ESchool of Veterinary and Medical Science, University of Nottingham, Sutton Bonington, LE12 5RD, UK.

^FCorresponding author. Email: ray.rodgers@adelaide.edu.au

Supplementary Material




Fig. S1. Hartanti *et al.*

Fig. S1. Variation of morphometric analysis. Data are presented as percentage of 3 – 40 fields of view and was counted by dividing standard deviation by the mean from each total number of fields of view to calculate the coefficient of variation (CV). The CV of the percentage of stromal area (A-F) and the proportion of stromal cells that are proliferating (G-L) in the ovarian cortex are presented. CRL 13.5 cm or 80 days (A and G), CRL 15.5 cm or 87 days (B and H), CRL 24 cm or 112 days (C and I), CRL 39 cm of 154 days (D and J), CRL 58 cm or 200 days (E and K) and CRL 80 cm or 245 days (F and L), respectively.