

AUSTRALIAN SYSTEMATIC BOTANY

An international journal devoted to the taxonomy, biogeography
and evolution of all plant groups

Contents Volume 23 Issue 6

23 December 2010

Tasmanian and Victorian populations of the fern *Asplenium hookerianum* result from independent dispersals from New Zealand.

L. R. Perrie, D. J. Ohlsen, L. D. Shepherd, M. Garrett, P. J. Brownsey and M. J. Bayly 387–392

Floral ontogeny in *Wisteria sinensis* (Fabaceae: Faboideae: Millettieae) and its systematic implications.

S. Naghiloo and M. R. Dadpour 393–400

Two new species and one reapplied name in *Sesbania* Scop. (Fabaceae, Sesbanieae) for Australia.

C. L. Gross 401–414

Evolution of nickel hyperaccumulation by *Stackhousia tryonii* (Celastraceae), a serpentinite-endemic plant from Queensland, Australia.

D. O. Burge and W. R. Barker 415–430

Phylogenetics of hopbushes and pepperflowers (*Dodonaea*, *Diplopeltis* – Sapindaceae), based on nuclear ribosomal ITS and partial ETS sequences incorporating secondary-structure models.

M. G. Harrington and P. A. Gadek 431–442

The *Lejeunea tumida* species group (Lejeuneaceae: Jungermanniopsida) in New Zealand.

M. A. M. Renner, E. A. Brown and G. M. Wardle 443–462