

[10.1071/SR21254](https://doi.org/10.1071/SR21254)

Soil Research

Supplementary Material

Effect of irrigation on soil physical properties on temperate pastoral farms: a regional New Zealand study

John J. Drewry^{A,}, Sam Carrick^B, Veronica Penny^B, John L. Dando^A, and Nina Koele^{C,D}*

^ASoils and Landscapes, Manaaki Whenua – Landcare Research, Private Bag 11052, Palmerston North, New Zealand.

^BSoils and Landscapes, Manaaki Whenua – Landcare Research, PO Box 69040, Lincoln, New Zealand.

^CLand Use and Ecosystems, Manaaki Whenua – Landcare Research, PO Box 69040, Lincoln, New Zealand.

^DPresent address: Te Hunga Mārama o te Taiao (Evidence, Data and Insights), Ministry for the Environment, PO Box 10362, Wellington, New Zealand.

*Correspondence to: John J. Drewry Soils and Landscapes, Manaaki Whenua – Landcare Research, Private Bag 11052, Palmerston North, New Zealand Email: drewryj@landcareresearch.co.nz

Supplementary data




Fig. S1. Distributions of available water content, AWC ($\% v v^{-1}$) for individual irrigated and dryland sites, by depth increment.




Fig. S2. Distributions of semi-available water content, SAWC ($\% v v^{-1}$) for individual irrigated and dryland sites, by depth increment.




Fig. S2. Distributions of water content at permanent wilting point, PWP ($\% v v^{-1}$) for individual irrigated and dryland sites, by depth increment.




Fig. S3. Distributions of bulk density ($Mg m^{-3}$) for individual irrigated and dryland sites, by depth increment.