

---

## Supplementary material

### **The influence of food availability, quality and body size on patch selection of coexisting grazer ungulates in western Serengeti National Park**

*John Bukombe<sup>A,G</sup>, Andrew Kittle<sup>B</sup>, Ramadhan B. Senzota<sup>C,\*</sup>, Hamza Kija<sup>D</sup>, Simon Mduma<sup>A</sup>, John M. Fryxell<sup>B</sup>, Flora Magige<sup>C</sup>, Cosmas Mligo<sup>E</sup> and Anthony R. E. Sinclair<sup>F</sup>*

<sup>A</sup>Biodiversity Program, Tanzania Wildlife Research Institute, PO Box 661, +255, 23113 LEMARA, Kikwakwaru ‘A’, Arusha, Tanzania.

<sup>B</sup>University of Guelph, Department of Integrative Biology, 50 Stone Road East, Guelph, Ontario, N1G 2W1, Canada.

<sup>C</sup>University of Dar es Salaam, Zoology and Wildlife Conservation department, PO Box 35064, +255, 14113 Kijitonyama, Mwenge, Dar es Salaam, Tanzania.

<sup>D</sup>Tanzania Wildlife Research Institute, PO Box 661, +255, 23113 LEMARA, Kikwakwaru ‘A’, Arusha, Tanzania.

<sup>E</sup>University of Dar es Salaam, Botany Department, PO Box 35065, +255, 14113 Kijitonyama, Mwenge, Dar es Salaam, Tanzania.

<sup>F</sup>Beaty Biodiversity Centre, Biodiversity Research Centre, 2212 Main Mall, Vancouver, BC Canada V6T 1Z4, University of British Columbia, Vancouver, Canada.

<sup>G</sup>Corresponding author. Email: bukombe2017@gmail.com

**Note:** All correlations were done using “PerformanceAnalytics”) in R version 3.5.1

Fig. S1. Correlation chart of coefficients (\*\*\*) =  $P < 0.0001$ , \* =  $P < 0.05$ ) between variables expressing diet quantity in grazing patches using all five selected ungulate species (buffalo, zebra, wildebeest, topi and impala) in western Serengeti National Park.



