

Evidence of significantly higher island feral cat abundance compared with the adjacent mainland

Patrick L. Taggart^{A,H}, Bronwyn A. Fancourt^{B,C}, Andrew J. Bengsen^D, David E. Peacock^{A,E}, Patrick Hodgens^F, John L. Read^G, Milton M. McAllister^{A,} and Charles G. B. Caraguel^{A,*}*

^ASchool of Animal and Veterinary Sciences, The University of Adelaide, Roseworthy, SA 5371, Australia.

^BPest Animal Research Centre, Biosecurity Queensland, Department of Agriculture and Fisheries, Toowoomba, Qld 4350, Australia.

^CSchool of Environmental and Rural Science, University of New England, Armidale, NSW 2350, Australia.

^DVertebrate Pest Research Unit, Department of Primary Industries, Orange, NSW 2800, Australia.

^EBiosecurity South Australia, Adelaide, SA 5001, Australia.

^FTerrain Ecology, PO Box 966, Kingscote, SA 5223, Australia.

^GSchool of Earth and Environmental Sciences, The University of Adelaide, Adelaide, SA 5005, Australia.

^HCorresponding author. Email: patrick.taggart@adelaide.edu.au

*Joint last author.

Figure S1. Time difference between consecutive cat detections at a particular camera location. Time difference ranges from 0–24 hours and is in increments of three hours from 180 min onwards.

