

Domestic dogs and water-availability effects on non-volant mammals in a protected area, south-eastern Brazil

*Priscila Stéfani Monteiro-Alves^A, Atilla Colombo Ferreguetti^{A,D}, Marina Mello Allemand^B,
Juliane Pereira-Ribeiro^A, Maja Kajin^{A,C}, Carlos Frederico Duarte Rocha^A and
Helena Godoy Bergallo^A*

^ADepartment of Ecology, Rio de Janeiro State University, Rua São Francisco Xavier 524, PHLC sala 220, Maracanã, 20550-900, Rio de Janeiro, RJ, Brazil.

^BDepartment of Biology, Centro Universitário Espírito-Santense/FAESA, Rua Anselmo Serrat, 199, Ilha de Monte Belo, 29053-250, Vitória, ES, Brazil.

^CDepartment of Biology, The University of Ljubljana – UL, Biotechnical Faculty – BF, Večna pot 111, 1000 Ljubljana, Slovenia.

^DCorresponding author. Email: atilla.ferreguetti@gmail.com

Table S1. Pearson correlation matrix of the covariates used in the Non-Metric Multidimensional Scaling (NMDS) and test whether there is an effect on mammalian species richness and abundance by using the generalized linear model (GLM) in the Mata das Flores State Park, Espírito Santo, Brazil.

	canopy	under	water	human	road	edge	dog
canopy	1						
under	0.41	1					
water	0.09	0.35	1				
human	-0.18	-0.18	0.52	1			
road	0.31	0.36	-0.27	-0.09	1		
edge	-0.22	0.31	0.12	0.35	0.38	1	
dog	0.12	-0.14	0.24	0.62	0.41	0.38	1

Covariates include canopy cover (canopy), understory cover (under), distance from water (water), distance from the human settlement (human), distance from the road (road), distance from forest edge (edge), and domestic dog abundance (dog).