

Supplementary Material

Are canid pest ejectors an effective control tool for wild dogs in an arid rangeland environment?

T. L. Kreplins^{A,E}, J. Miller^B and M. S. Kennedy^{C,D}

^ADepartment of Primary Industries and Regional Development, 75 York Road, Northam, WA 6401, Australia.

^BDepartment of Primary Industries and Regional Development, 10 Doney Street, Narrogin, WA 6312, Australia.

^CDepartment of Primary Industries and Regional Development, 3 Baron-Hay Court, South Perth, WA 6151, Australia.

^DPresent address: Department of Agriculture and Fisheries, 203 Tor Street, Toowoomba, Qld 4350, Australia.

^ECorresponding author. Email: tracey.kreplins@dpird.wa.gov.au

Appendix S1. The density (per linear km; \pm SE) of wild dogs, g0, sigma (km) and number of individual wild dogs in the sampled area during each session of the canid pest ejector trials as estimated by the secr linear analysis.

Session	Month	Year	Density (km)	SE	g0	σ (km)	N
1a	September-October	2018	0.38	0.26	0.01	6.31	50
1b	October-November	2018	0.21	0.18	0.01	6.31	45
2a	March-April	2019	0.19	0.05	0.02	4.80	32
2b	April-May	2019	0.18	0.05	0.02	4.80	31
3a	September-October	2019	0.13	0.04	0.00	2.99	19
3b	October-November	2019	0.11	0.04	0.00	2.99	18
4a	March-April	2020	0.23	0.06	0.00	7.71	37
4b	April-May	2020	0.14	0.05	0.00	7.71	30