

Sympathy for the devil: captive-management style did not influence survival, body-mass change or diet of Tasmanian devils 1 year after wild release

Tracey Rogers^{A,C}, Samantha Fox^B, David Pemberton^B and Phil Wise^B

^AEvolution and Ecology Research Centre, School of BEES, The University of New South Wales, Sydney, NSW 2052, Australia.

^BWildlife Monitoring and Management Section, Wildlife Management Branch, Department of Primary Industries, Parks, Water and Environment, GPO Box 44, Hobart, Tas. 7001, Australia.

^CCorresponding author. Email: tracey.rogers@unsw.edu.au

Abstract

Context. The value of captive breeding for recovery programs of endangered carnivorous mammals is often questioned because of low post-release survival reported for founder animals following translocation.

Aims. The aim of the present study was to test the effect of rearing method on survival and body mass of captive-raised Tasmanian devils (*Sarcophilus harrisii*) following release on an offshore island. We also compared the post-release diet of these devils with the diet of wild devils on mainland Tasmania, where a similar array of diet items is available.

Methods. Twenty-eight captive-raised devils were released onto the island; 19 had been raised in intensive captive-management facilities (IC) and nine in free-range (22 ha) enclosures (FRE). Survival and body-mass change were compared between IC and FRE for up to 440 days post-release. Devil diet was assessed via scat analysis.

Key results. A high proportion (96%) of the founders survived 1 year post-release. Pre-release captive-rearing method had no effect. Released devils gained an average of 14% of their original body mass, irrespective of captive-rearing method. There was very little difference in the diet of captive-reared devils released onto Maria Island relative to wild mainland devils: Tasmanian pademelon, *Thylogale billiardieri*, was the primary food item for both.

Conclusions. The intensity of captive rearing did not affect the survival of devils released onto Maria Island. This suggests that even devils held in IC facilities retain the innate behaviour required to scavenge and hunt prey, and therefore maintain bodyweight post-release. The lack of any threatening processes on the island is also likely to have contributed to the high survival rate 2 years post-release.

Implications. Our study provided preliminary evidence that the release of captive-raised Tasmanian devils onto off-shore islands is a viable conservation action. Captive-breeding programs and captive-raised founders can play a viable and valuable role in the conservation action plans for recovery programs of endangered carnivorous mammals.

Additional keywords: applied ecology, conservation biology, endangered species, foraging, vertebrates, wildlife management.

Received 3 December 2015, accepted 2 September 2016, published online 2 November 2016

Introduction

Translocation programs for threatened species have had variable but often low success, especially those involving carnivores (Griffith *et al.* 1989; Wolf *et al.* 1996; Fischer and Lindenmayer 2000; Soorae 2013; Tarszisz *et al.* 2014), and survival rates decrease when captive-raised animals are used (Mathews *et al.* 2005; Jule *et al.* 2008), although the reasons for this are not always clear. In certain taxa (e.g. fish), rapid evolution after a single generation in captivity can result in substantial selection for traits that are beneficial in captivity but reduce the fitness of captive-raised individuals when they are reintroduced into the wild (Christie *et al.* 2012). Captive animals can lose natural

behaviours associated with wild fitness (Snyder *et al.* 1996; Rabin 2003), particularly behaviours specifically associated with socialisation, foraging and hunting in birds (van Heezik and Ostrowski 2001; Vickery and Mason 2003) and mammals (Stoinski *et al.* 2003; Mathews *et al.* 2005). One of the primary underlying causes of translocation failures in captive-raised mammalian carnivores has been starvation (Jule *et al.* 2008) because of the absence of foraging behaviours. This raises important animal-welfare concerns (Mathews *et al.* 2005; Tarszisz *et al.* 2014). Thus, it is critical to examine how captive-bred marsupial carnivores will behave in the wild post-release, particularly at the initial stages of a program.

The Tasmanian devil (*Sarcophilus harrisii*) is a carnivorous marsupial that scavenges on carcasses and hunts for small prey such as insects, lizards, frogs, birds and small mammals (Pemberton and Renouf 1993; Pemberton *et al.* 2008). It was listed as endangered under the Environment Protection and Biodiversity Conservation Act (Hawkins *et al.* 2008) and, across its Tasmanian distribution, it has experienced population declines of 60–80% over a 10-year period (Hawkins *et al.* 2006; McCallum *et al.* 2009; DPIPWE, unpubl. data). The reduced population numbers are largely due to devil facial-tumour disease (DFTD; McCallum *et al.* 2009), which is an infectious fatal cancer (Pearse and Swift 2006). Disease transmission occurs when live tumour cells pass between animals during aggressive interactions over food or during mating (Hamede *et al.* 2013). The disease is spreading west across Tasmania at a rate of ~5–10 km per year, and has now been detected across the majority of the devil's geographic range (DPIPWE, unpubl. data). Recovery of individuals after clinical symptoms present has not been documented (Hamede *et al.* 2013). Alongside DFTD, the synergistic effects of habitat loss, human-related factors (e.g. roadkill) and the increase in introduced predators, such as cats and foxes, are thought to be important in the observed decline. As part of a conservation effort, DFTD-free devils are being bred in captivity in Tasmania and on mainland Australia. This insurance population was established for use in population supplementation projects to boost the genetic diversity and environmental functionality of wild devil populations.

Captive-reared Tasmanian devils released onto the Tasmanian mainland had moderate survival (~42%) 2–8 months post-release (Sinn *et al.* 2014). Twenty-eight DFTD-free captive devils have been released onto Maria Island (a devil-free Tasmanian island), and the present study examines their survival and diet. Maria Island is ~12 km off the south-eastern coast of Tasmania. Although devils have not been recorded there (Hope 1972), the island has a range of the devil's prey species and is large enough to theoretically support a large devil population. Ongoing management is required, through additional releases, to maintain genetic diversity, and to make removals once carrying capacity has been reached (DPIPWE 2011). Maria Island was connected to mainland Tasmania as recently as 5000–8000 years ago; thus, it is likely that devils were on Maria Island in the past during glacial periods.

Tasmanian devils are being bred at 35 captive-management institutions using two different methodologies, namely, free-range enclosures (FRE) (22-ha pens) and intensively managed captive facilities (IC; e.g. zoos, sanctuaries, hand-rearing). We predict that the intensity of captive-rearing will influence survival of released animals because the free-range pens provide opportunities for captive devils to hunt and forage, which is not possible in intensively managed facilities. Any loss of natural hunting behaviours is likely to be greatest among animals that have been raised in IC facilities. Reduced hunting abilities for these animals would mean that they would lose more body mass post-release than do animals raised in FRE facilities.

We further predict that captive-rearing will influence the ability of devils to forage in the wild; thus, captive-raised and released devils will have a diet different from that of wild devils.

As there are no natural devil populations on other offshore islands, we compared the diet of Maria Island devils to that of animals on mainland Tasmania. Nonetheless, the main diet items recorded for mainland devils (e.g. ringtail possum (*Pseudocheirus peregrinus*), Tasmanian bettong (*Bettongia gaimardi*), Tasmanian pademelon, common brushtail possum (*Trichosurus vulpecula*), short-beaked echidna (*Tachyglossus aculeatus*), red-necked wallaby (*Macropus rufogriseus*), common wombat (*Vombatus ursinus*) and eastern grey kangaroo (*Macropus giganteus*)) are known to occur on Maria Island, so prey availability is not likely to differ substantially.

The devil takes a broad range of prey, from as small as 0.01 kg (e.g. frogs and moths) to large macropods (Pemberton and Renouf 1993; Pemberton *et al.* 2008). Compared with other terrestrial mammalian carnivores (Carbone *et al.* 1999; Tucker and Rogers 2014a, 2014b), the devil, with a body mass of ~8–10 kg for a male, consumes relatively large species compared with its body size. For example, the Tasmanian pademelon, red-necked wallaby, common wombat and eastern grey kangaroo are ~80%, 100%, 260% and 420% of the devil's body mass respectively. These large species are available as roadkill on the mainland and the devil presumably acts as a scavenger when it feeds on them. Maria Island is a national park, with little road traffic. The translocated Maria Island devils are less likely to use these large species because they may not be available as carrion. We cannot determine whether the method of captive management (IC or FRE) influences the diversity of prey species because we cannot attribute wild-collected scat to an individual devil.

We propose the following three hypotheses: (1) animals that have been raised in intensively managed facilities will have lower survivorship than do animals raised in free-range pens; (2) animals that have been raised in intensively managed facilities will have greater body-mass losses; and (3) captive-raised devils will have a diet different from that of wild devils on the mainland.

Release of captive-raised animals, particularly mammalian carnivores, for conservation programs is risky; however, it is important to make decisions and implement programs while opportunities to act still remain (Martin *et al.* 2012). In the situation described herein, inaction could result in the loss of the world's largest extant marsupial carnivore, the Tasmanian devil.

Materials and methods

Animal release

Twenty-eight adult (~1 year of age) Tasmanian devils (13 females and 15 males) were released onto devil-free Maria Island. Nineteen of these animals were reared in IC management facility and nine were reared in 22-ha FRE facilities (Table 1). The release site at Four Mile Creek has a clear flowing stream. The release strategy for both groups was the same. There were two releases, with the first 15 animals released on 12 November 2012 and an additional 13 animals released about 1 year later (between 25 October and 1 November 2013; Table 1). Fifteen animals were selected for the first release because this number allowed for adequate post-release monitoring. The number of IC-raised ($n = 7$) and FRE-raised ($n = 8$) animals was similar. The second release was predominantly of IC ($n = 12$) animals ($n = 1$, FRE).

Table 1. Fate and body-mass (BM) change of 28 captive-raised adult Tasmanian devils released on Maria Island

The fate of devils as of 12 January 2015 and covariate data on 28 captive-raised ~12-month-old Tasmanian devils released on Maria Island. Source institutions for each rearing style are given as superscript numbers, as follows: ¹Bridport free range enclosure, ²Freycinet free range enclosure, ³Murdunna–Taroona Intensive Captive Facility (DPIPWE), ⁴Taroona Intensive Captive Facility (DPIPWE), ⁵Trowunna Wildlife Park (Tasmania), ⁶Healesville Sanctuary (Vic), ⁷Monarto Zoo (S.A) and ⁸orphaned wild animals hand-raised in intensive captive institutions. The BM change over time (up to 26 months post-release) was identified for each individual ($n = 26$ devils where sufficient consecutive trapping data) as functions derived using the following coefficients of the polynomial regression equation: b_0 is the coefficient that is a function of the BM at release, b_1 is initial BM change immediately following release, b_2 is BM change several months after release, and b_3 is change at the completion of the study. Individuals with similar patterns of BM change (using the polynomial regression coefficients b_0 , b_1 , b_2 , and b_3 as a proxy) were classified into groups (clusters), initially into two groups (A and B), and then to subclusters (A_{1-4} and B_{1-3}), by using a hierarchical agglomerative cluster analysis. Clusters were linked using Ward's method and separated using squared Euclidean distances. Superscript F, first release; superscript S, second release; FR, free-range 22-ha enclosure; and IC, intensive captive managed, e.g. zoos

ID	Months post-release	Fate	Sex	Rearing style	Release BM (kg)	Clusters	Polynomial regression coefficient			
							b_0	b_1	b_2	b_3
Armin	26 ^F	Alive	M	FR ²						
Manny	1 ^F	Dead	M	IC ⁴	8.6					
Becks	26 ^F	Alive	F	IC ⁴	5.6	A ₁	6	-0.1	0.02	0
China Girl	26 ^F	Alive	F	FR ¹	6.1	A ₁	6.7	-0.18	0.007	0
Florence	26 ^F	Alive	F	FR ²	5.5	A ₁	5.6	-0.2	0.03	-0.001
Lola	26 ^F	Alive	F	FR ¹	5.9	A ₂	5.9	0.22	0.05	-0.001
Lolita	26 ^F	Alive	F	FR ²	5.9	A ₂	5.7	0.3	-0.02	0
Oddity	20 ^F	Alive	M	FR ¹	6.7	A ₂	6.6	0.27	0	-0.005
Dylan	14 ^S	Alive	M	IC ⁶	5.6	A ₂	5.4	0.42	-0.05	0
Muffs	26 ^F	Alive	M	IC ⁴	9	A ₃	8.7	0.02	-0.01	0
Sirius	26 ^F	Alive	M	IC ⁴	7.5	A ₃	7.2	0.29	-0.02	0
Snips	26 ^F	Alive	M	IC ⁴	8.1	A ₃	8	0.04	0.008	0
Toby	14 ^S	Alive	M	IC ⁶	7.9	A ₃	7.5	0.05	-0.003	0.001
Francesca	26 ^F	Alive	F	FR ²	4.9	A ₄	4.8	0.83	-0.08	0.002
Reba	26 ^F	Alive	F	IC ^{3,8}	4	A ₄	3.8	0.26	-0.02	0.001
Remmy	26 ^F	Alive	F	IC ^{3,8}	3.4	A ₄	3.4	0.03	-0.02	0
Jimmy	17 ^F	Presumed dead	M	FR ²	9.6	B ₁	10	-1.26	0.28	-0.015
Mozzie	14 ^S	Alive	F	IC ⁵	6.1	B ₁	6.2	-0.49	0.26	-0.03
Axel	14 ^S	Alive	M	IC ⁶	9.8	B ₂	9.6	-0.76	0.11	-0.004
Boomer	14 ^S	Alive	M	IC ⁶	9.8	B ₂	9.4	-0.67	0.1	-0.004
Bowie	14 ^S	Alive	M	FR ¹	9.4	B ₂	9.4	-0.31	0.12	0
Gateman	14 ^S	Alive	M	IC ⁶	8.8	B ₂	8.8	-0.73	0.09	0
Gus	14 ^S	Alive	M	IC ⁶	9.3	B ₂	9.3	-0.83	0.17	-0.008
Chee	14 ^S	Alive	F	IC ⁶	6.4	B ₃	6.3	-0.41	0.11	-0.006
Emily	14 ^S	Alive	F	IC ⁶	5.7	B ₃	5.6	-0.05	0.13	-0.006
Lilli	14 ^S	Alive	F	IC ⁷	7.6	B ₃	7.5	-0.57	0.06	-0.002
Marumba	14 ^S	Alive	M	IC ⁷	7.6	B ₃	7.5	-0.61	0.13	-0.005
Nutella	14 ^S	Alive	F	IC ⁷	5.7	B ₃	5.6	-0.61	0.14	-0.007

Survivorship and body-mass change

The mean life expectancy of wild Tasmanian devils is 5 years (Jones 2001). A baited-trapping and weighing program was conducted monthly for the first 4 months following each release, and every 3 months thereafter. During each trapping session, up to 40 traps were deployed over seven consecutive nights across accessible areas of Maria Island. All activities were conducted in accordance with the standard operating procedures for trapping and handling wild Tasmanian devils (DPIPWE 2011) approved under the Animal Care and Ethics Committee of the Department of Primary Industries, Parks, Water and Environment (DPIPWE 2011), Hobart. The traps were placed in areas considered conducive to regular devil movement (i.e. the intersection of two or more tracks or beside a track where a culvert/drainage/creek crosses underneath) and located in shaded quiet sites. The traps were custom-designed (N. J. M. Mooney and D. Ralph, unpubl. data) PVC pipe traps ~315 mm in diameter and 875 mm long. The traps were inspected throughout the

night as well as in the early morning hours. The animals were individually identified by scanning for subcutaneous microchip transponders (Trován, Microchips Australia, Melbourne, Victoria, Australia) and then weighed.

Diet

Following release, carcasses in variable quantities were used as lure as part of the ongoing monitoring program; an average of 124 kg (s.d. 124 kg) was used per trapping event following the first release and an average of 32 kg (s.d. 29 kg) per trapping event following the second release, for three months after each release event. Trapping schedule is outlined above in 'Survivorship and body-mass change'. Scat was not collected during the 3 weeks following carcass provisioning. In total, 105 devil-scat samples were collected over 662 days from Maria Island between 19 November 2012 and 12 November 2014. Samples collected in 2014 may have included scats from juvenile animals because young were born on the island. To ensure representative sampling

of wild-devil diet, 161 Tasmanian devil-scat samples were collected from five disparate Tasmanian mainland locations, including Fentonbury ($n=14$, October to December 2004), Bronte ($n=45$, October 2004 to March 2005), Mersey Forest ($n=28$, May 2012), Waratah ($n=15$, November 2011) and Woolnorth ($n=46$, December 2012); scats with unknown locations were also included ($n=13$, June 2005 to August 2008).

It was not possible to distinguish between diet items that were prey (known or likely to have been predated on) and those that had been scavenged from carcasses; thus, we refer to all species detected in the scat as diet items. We attempted to identify diet items in the Tasmanian devil scat to the species level. However, bird species were not distinguishable from the feather remains and were considered a single diet-item group, namely 'feathers'.

To evaluate our prediction that captive-raised devils will have a diet different from that of wild (mainland) devils, we used a range of diversity indices, including the following: species richness (S ; Whittaker 1972); Shannon–Wiener diversity (H') where the H' index measures the value of any species as a diet item as a function of their frequency in the community; effective number of species (ENS); and Brillouin's (HB) index where a high HB value represents a high dietary diversity, suggesting that many different species are in the diet and that they are used more evenly (Brillouin 1956). Indices were calculated as follows:

$$S = \text{number of species,}$$

$$H' = - \sum p_i \times \ln p_i,$$

$$ENS = H' / \ln N_{\max}, \text{ and}$$

$$HB = (\ln N! - \sum \ln n_i!) / N.$$

where n = the total number of individual diet items recorded in scat samples, n_i = the number of individual dietary items in the i_{th} category, N_{\max} = the number of individuals in the most abundant species and $p_i = n_i / N$.

The frequency of occurrence (FO) of a diet item (species or species group) was expressed as a percentage and calculated as the number of occurrences of a diet item in the scat samples, divided by the total number of scat samples collected. The results were expressed as the percentage occurrence. The CO, the 'diet item composition' was the number of occurrences of a species or species group, calculated by dividing the number of times a particular diet item was identified by the total number of occurrences for all diet items.

To convert the FO data to biomass consumption, we multiplied the FO by a linear function (Ackerman's equation) for each diet species (Ackerman *et al.* 1984), as follows:

$$y = 1.980 + 0.035x,$$

where y = the biomass consumed (kg) that produced a single field-collectable scat sample and x = the average bodyweight of the diet species (kg; Table 2).

Ackerman's equation was developed for felids (specifically cougar (*Felis concolor*); Ackerman *et al.* 1984), but has been used for a range of mammalian carnivores, including the crab-eating fox (*Cerdocyon thous*), ocelot (*Leopardus pardalis*), puma (*Puma concolor*) and jaguar (*Panthera onca*; Farrell *et al.* 2000). Although the digestive physiology of the Tasmanian devil is likely to differ from that of eutherian carnivores, this equation was used as an approximation.

Statistics

We used a mixed-model ANOVA to examine the mean change in BM (relative to BM at release) over 440 days post-release of

Table 2. Diet species consumed by wild and captive-raised Tasmanian devils

Diet species in the scat samples of Tasmanian mainland (wild, $n=161$ scats) and captive-raised Maria Island (MI, $n=105$ scats) Tasmanian devils are presented as follows: %FO, relative frequency of occurrence; %CO, relative composition; and %biomass, relative biomass consumed of each species. The P -values reported are from chi-square analysis, performed where values were 5 or greater, so as to examine whether there were significant differences in the proportion of each species used between the wild and Maria Island populations. The relative biomass consumed (%) was estimated with Ackerman's equation (Ackerman *et al.* 1984), using the frequency of occurrence of prey in the faeces and the body mass of the prey (Jones *et al.* 2009). P–P BM, predator–prey body mass; mass, mass of species consumed by Tasmanian devils in kg; %, the percentage of the prey's body mass relative to the body mass of the Tasmanian devil (10 kg)

Diet species	%FO			%CO			%Biomass			P–P BM Mass, %
	Wild	MI	P	Wild	MI	p	Wild	MI	P	
Cattle	1	0		1	0					
Sheep	4	0		4	0					
Dog	1	0		1	0					
Rabbit	1	0		1	0					
House mouse	1	0		1	0					
Feathers	7	21	0.05	7	18	0.05				
Ringtail possum	9	4		9	3		9	3	0.03	1, 10
Tasmanian bettong	0	1		0	1		0	1		2, 20
Brush-tail possum	17	29	0.36	18	25	0.44	18	25	0.03	3, 30
Echidna	1	3		1	2		1	2		5, 50
Tasmanian pademelon	29	27	0.22	29	23	0.25	29	23	0.35	8, 80
Red-necked wallaby	23	10	0.003	24	9	0.004	24	9	0.0001	10, 100
Wombat	6	19	0.05	6	17	0.04	6	17	0.00004	26, 260
Eastern grey kangaroo	0	2		0	2		0	2		42, 420
<i>Macropus</i> sp.	7	0		8	0					

26 of the captive-raised animals (Table 1). Two of the original 28 released animals were excluded from the BM-change analysis, because there were fewer than 4 monthly trapping weights available post-release (trap-shy animals). The animals excluded from analysis were both from the first release, one was an IC and one FRE reared.

The factors for the mixed-model ANOVA corresponded to a hierarchy of levels, with correlated measurements of the dependent variable (relative BM change) occurring at each level. The fixed factors were assumed to be (1) sex and (2) the rearing style (assuming that the animals were not randomly selected from a larger population of captive animals). Time (months) was assumed to be a random factor, because the trapping of the animals was not a fixed event occurring at regular time intervals, with no missing values. The results of the mixed-model ANOVA were not compromised by violations of the assumptions of normality, because the residuals were normally distributed, or by heteroskedasticity, because the variances were homogeneous across the levels (Levene's test $P=0.376$).

Cubic polynomial regression equations (examples for two devils, Fig. 1*a, b*) of the following form were computed to describe the patterns of BM change (kg) of each of 26 animals against time (months) post-release:

$$\text{Body mass (kg)} = b_0 + b_1 \times \text{months} + b_2 \times \text{months}^2 + b_3 \times \text{months}^3,$$

where b_0 , b_1 , b_2 and b_3 are polynomial regression coefficients. The constant b_0 is a function of the BM of the devils at the time of release. The first-order regression coefficient b_1 is a function of the initial change in BM, immediately following release. The second-order regression coefficient b_2 is a function of a subsequent change in BM, several months after release. The




Fig. 1. Body-mass change and diet diversity of captive-raised devils following translocation. Polynomial regressions were used to define the post-release BM change that classified the captive-reared devils into clusters; an example from (a) Cluster A, an adult male (Muffs) from the first release; and (b) Cluster B, an adult male (Gus) from the second release.

third-order regression coefficient b_3 is a function of a subsequent change in BM towards the end of the study. For the devils in the first release, in 2012, this is at 825 days post-release, whereas for the devils in the second release, in 2013, this is at 440 days post-release (Table 1). The data were good fits to the polynomial model (indicated by R^2 -values up to 99.9%).

The polynomial equations defined above do not describe the growth of the animals, but, rather, reflect temporal fluctuations in the BM. The devils were released as adults, so it is presumed that BM change reflects a change in body condition (e.g. loss or gain of soft tissue such as reserves of adipose and muscle tissue) rather than growth. BM gain is presumed to reflect successful foraging, whereas BM loss reflects unsuccessful foraging.

To classify the 26 devils into groups according to their BM-change patterns, we used a hierarchical agglomerative cluster analysis that used the coefficients of the polynomial regression equations (b_0 , b_1 , b_2 and b_3). The clusters were linked using Ward's method and separated using squared Euclidean distances. A dendrogram was constructed, in which, the further apart the clusters were, the greater were the differences between the patterns of BM change of the animals in each cluster.

Diet

Species diversity curves (Shannon–Wiener (H') index (Fig. 2*a*), representing diversity; Brillouin's (HB) index (Fig. 2*b*), representing dominant species) showed that the number of diet




Fig. 2. Diet diversity of captive-raised and wild devils. Cumulative observations of the frequency of occurrence of diet items in the scat samples. Diet diversity is represented by (a) the Shannon–Wiener (H') index and (b) the Brillouin index (HB). The Shannon–Wiener index represents the value of any species as a diet item as a function of their frequency in the community, whereas the Brillouin's index represents dietary diversity. Scats are ordered according to date collected. Circles represent cumulative values, black circles represent captive-raised Maria Island devils, and white circles represent wild mainland devils.

items in the scat reached an asymptote after analysing ~40 scat samples. This result indicated that the diets of Maria Island ($n=105$) and mainland ($n=161$) devils were adequately sampled (Fig. 2).

Results

Survival

Of the 15 animals in the first release, most (87%, 13 of 15) were alive 825 days post-release (Table 1). One animal, a 2-year-old male (Manny) reared in an IC-management facility, died within the first month after release. A second animal, a 5-year-old male (Jimmy) reared in an FRE facility, was presumed dead because he had not been observed in the trapping program after 460 days. Of the 28 captive-raised devils that were released, 96% (27 of 28) were alive 365 days later (Table 1). The rearing style (IC or FRE) did not influence survival at least up to the time of this report. The mean age of the surviving founders was 3 years (0.9 s.d., $n=26$).

Body-mass change

The released devils had an average proportional BM gain of 0.14 (0.17 s.d., range 0.22–0.78, $n=150$) over the post-release period of 440 days. The mixed-model ANOVA indicated that none of the factors significantly (at the 0.05 α -level) influenced the mean BM change (relative to BM at release) of the captive-reared devils over the first 440 days post-release (Table 3). However, the effect size was very small (adjusted $R^2=11.85\%$).

The BM changes in each individual animal over time were explored using non-linear regression analysis. Polynomial regression coefficients defined mostly sigmoid-shaped trajectories to explain BM-change patterns (Fig. 1a, b). A dendrogram of a hierarchical cluster analyses, which used the coefficients of the polynomial regression equations (b_0 , b_1 , b_2 and b_3), displayed that one major dichotomy divided the devils into two groups labelled Clusters A and B. The Clusters A and B were separated predominantly according to release date (Table 1), but not according to any of the other factors (e.g. sex, birth year or rearing style). Most animals in Cluster A were from the first group of devils released in 2012. Only the one devil in Cluster A, namely Dylan, had been from the second group released in 2013. All but the one animal in Cluster B were released in 2013, as part of the second release. Jimmy, the oldest devil, a 3-year-old male at the time of release, was the only animal in Cluster B from the first release.

The values of coefficient b_0 , a function of the body mass at release, was lowest in a group of females in Cluster A, and highest in males cluster within Clusters B and A (Table 1).

Table 3. Mixed-measures ANOVA of the body mass change (relative to the body mass at release) over 440 days post-release for 25 captive-reared Tasmanian devils

Parameter	d.f.	SS	MS	F	P
Captive management style	1	0.002	0.002	0.07	0.792
Months since release	6	0.657	0.11	4.01	0.058
Sex	1	0.079	0.079	2.88	0.141
Sex \times months since release	6	0.164	0.027	1.11	0.362

This is unsurprising because the Tasmanian devil is sexually dimorphic.

Diet

The Maria Island devils consumed significantly (χ^2 : FO, $P=0.0007$; CO, $P=0.001$; biomass, $P<0.1000$; d.f.=4; Table 2) different proportions of some species as diet items. They consumed significantly greater proportions of wombat (by FO, CO and biomass), bird (by FO and CO) and brushtail possum (by biomass), but significantly smaller proportions of red-necked wallaby and ringtail possum (by biomass; Table 2) than did the devils from the wild mainland population.

The H' index was lower for the Maria Island population ($H'=1.82$), indicating that 6.2 species was the effective number of species (ENS) used, compared with an ENS of 7.2 species for the mainland population ($H'=1.97$; Table 2). The HB from all mainland sites ($HB=1.84$, $n=161$) was slightly higher than that for the Maria Island ($HB=1.70$, $n=105$) devils. Wombat and bird were important for Maria Island devils, whereas red-necked (Bennet's) wallaby and ringtail possum were used by mainland devils (Table 2). The captive-raised devils and their offspring did not consume significantly ($\chi^2=0.7$, d.f.=1, $P=0.4$; Table 2) different numbers of species ($S=9$) compared with the wild mainland devils ($S=11$).

Discussion

Survival

The method of captive-rearing did not influence survivorship post-release. In contrast to our prediction, Tasmanian devils raised in IC facilities did not experience lower survivorship. The proportion of captive-bred founders surviving on Maria Island was higher (~96% over 2 years) than that observed for animals released on the mainland (~42% over 2–8 months) in the study of Sinn *et al.* (2014).

A review of 45 carnivore reintroduction programs (17 species across 5 families) that used captive-raised animals as founders found a low (~30%) founder survival success 12 months post release (Jule *et al.* 2008). More recent translocation programs have reported higher post-release founder success for captive-raised mammalian carnivores than in the studies reviewed by Jule *et al.* (2008), namely, 61% for the Vancouver Island marmot (*Marmota vancouverensis*; Aaltonen *et al.* 2009), 50% for the European mink (*Mustela lutreola*; Maran *et al.* 2009) and 53–67% for the Iberian lynx (*Lynx pardinus*; Simón *et al.* 2013). Largely because of this low survival of released captive-bred animals, a great deal of debate has surrounded the value of captive-raised founders for conservation programs (Armstrong and Seddon 2008; Jule *et al.* 2008). The generally low success rate in releases of captive-raised founders, including devils (Sinn *et al.* 2014), has been a concern for managers of the Tasmanian devil program. Over the long-term, captive-raised devils are an important part of the conservation strategy to maintain devil insurance populations, while DFTD moves through the wild population. Thus, the survivability of captive-raised devils in the wild post-release must be maximised. The Maria Island population, including the devils released, and subsequently born, has risen to more than 90 Tasmanian devils (DPIPWE, unpubl. data, part of a further study) in the 2.5 years

since the commencement of the release program. The high post-release survival of the captive-raised devils, which showed little influence from pre-release management methods, is promising; however, several favourable factors influenced the outcome of this translocation program. Human-related activities (shooting, poisoning and car collisions) have directly caused the death of captive-raised founders in over 50% of all translocation programs for mammalian carnivores (Jule *et al.* 2008) and roadway-related mortalities are significant for devils on the Tasmanian mainland; however, Maria Island has little vehicular traffic. Predation and inter-specific competition and aggression have also reduced founder survivorship in other mammalian carnivore translocations, with ~50% of captive-raised European mink killed by other carnivores and raptors within the first few months following release (Maran *et al.* 2009), and newly released captive-raised African wild dogs (*Lyacon pictus*) killed by lions (Moehrensclager and Somers 2004). Although young devils, less than 12 months old, are vulnerable to predation, adult Tasmanian devils have few natural predators in Tasmania. Because the devils that were released on Maria Island were at least 12 months old, they were less vulnerable to predation.

Body-mass change

In contrast to our prediction, rearing style (intensive or free-range management) did not influence the body condition of devils post-release. On average, the captive-raised devils gained weight during the release period. Although rearing style did not influence BM change, the year of release did. Although devils in the first release gained mass immediately following release, the devils in the second release, 1 year later, lost weight immediately following release. Releases occurred at the same time, in the austral spring (October 2012 and 2013), timed to maximise food availability for newly released devils. The differences in BM change following release between years may have occurred because of a shift in species availability or climatic variability.

Diet

In contrast to our prediction, the captive-raised devils did not have a diet substantially different from that of wild mainland devils. The captive-raised founders and their offspring used a similar broad diversity of diet species and consumed the same dominant food items typical of wild devils from five different Tasmanian locations. The Tasmanian pademelon was the primary food item used by both Maria Island and mainland devils. The Maria Island devils consumed different proportions of some species, such as, for example, greater quantities of wombat, bird and brushtail possum and lower proportions of red-necked wallaby and ringtail possum. This may reflect differences in species availability on the island and the mainland. Although red-necked (Bennett's) wallaby and ringtail possum are present on the island (DPIPWE data) and important in the diet of mainland devils, their remains were less important in the diet of Maria Island devils.

Bird feathers were found more frequently in and represented a greater proportion of the scats of the Maria Island population than on the mainland. Maria Island is an important nesting site

for ground-nesting birds, including short-tailed shearwaters (*Puffinus tenuirostris*), little penguins (*Eudyptula minor*) and Cape Barren geese (*Cereopsis novaehollandiae*). Unfortunately, because the majority of bird bones and feathers were small, they could not be identified to the species level. However, devils on the island were observed by the authors to prey on all three of these ground-nesting bird species.

Devils are considered scavengers that hunt (Owen and Pemberton 2005), and their massive skull can produce high bite forces for their size (Attard *et al.* 2011), thus allowing them to crack open bones. A large proportion of the diet items used by devils on mainland Tasmania are believed to be carrion. The current Tasmanian mainland landscape has been greatly altered, and a large quantity of available carrion is created by anthropogenic activity, such as road mortality or farmland-generated waste (Owen and Pemberton 2005). Although Maria Island is heavily modified, it does not have these anthropogenically derived resources. As less carrion is available, devils have less opportunity to scavenge on Maria Island than on the mainland, and yet, the larger-sized species, such as the Tasmanian pademelon, common wombat and eastern grey kangaroo, remain important diet items. Thus, it is likely that Maria devils hunt large prey.

Conclusions

Although the survivorship of released captive-raised endangered carnivores has traditionally been low for translocation programs, our study indicated high founder survivorship following the first stage of a release program that employed captive-raised Tasmanian devils. The high post-release survival of the captive-raised devils was irrespective of pre-release rearing style (intensive or free-range). The rearing method did not affect the body-condition change, and captive-raised devils maintained and further increased their weight post-release. The captive-raised founders were not susceptible to starvation as a result of a lack of foraging skills; they had a broad diet similar to the diet of wild mainland devils. Although the program is at an early stage (approximately 2 years post-release), the preliminary indicators are that captive propagation with the subsequent release of captive-reared individuals on devil-free islands can play a viable role in the conservation action for the Tasmanian devil. In specific contexts, captive-breeding programs, including both intensive and free-range management styles, are valuable for recovery programs of endangered carnivorous mammals.

Acknowledgements

Funding was provided by Australian Federal Government to the Save the Tasmanian Devil Program. Phil Iles, Nick Mooney and Oliver Gales are thanked for scat collections and Barbara Triggs for her scat analysis. Oliver and Nick are also thanked for discussions on diet of Tasmanian fauna. T. R. was supported by a grant from Australian Research Council (LP0989933).

References

- Aaltonen, K., Bryant, A. A., Hostetler, J. A., and Oli, M. K. (2009). Reintroducing endangered Vancouver Island marmots: survival and cause-specific mortality rates of captive-born versus wild-born individuals. *Biological Conservation* **142**, 2181–2190. doi:10.1016/j.biocon.2009.04.019

- Ackerman, B. B., Lindzey, F. G., and Hemker, T. P. (1984). Cougar food habits in southern Utah. *The Journal of Wildlife Management* **48**, 147–155. doi:10.2307/3808462
- Armstrong, D. P., and Seddon, P. J. (2008). Directions in reintroduction biology. *Trends in Ecology & Evolution* **23**, 20–25. doi:10.1016/j.tree.2007.10.003
- Attard, M. R. G., Chamoli, U., Ferrara, T. L., Rogers, T. L., and Wroe, S. (2011). Skull mechanics and implications for feeding behaviour in a large marsupial carnivore guild: the thylacine, Tasmanian devil and spotted-tailed quoll. *Journal of Zoology* **285**, 292–300. doi:10.1111/j.1469-7998.2011.00844.x
- Birillouin, L. (Ed.) (1956). 'Science and Information Theory.' (Academic Press: New York.)
- Carbone, C., Mace, G. M., Roberts, S. C., and Macdonald, D. W. (1999). Energetic constraints on the diet of terrestrial carnivores. *Nature* **402**, 286–288. doi:10.1038/46266
- Christie, M. R., Marine, M. L., French, R. A., and Blouin, M. S. (2012). Genetic adaptation to captivity can occur in a single generation. *Proceedings of the National Academy of Sciences, USA* **109**, 238–242. doi:10.1073/pnas.1111073109
- DPIPWE (2011). Proposal to establish a managed population of Devil facial tumour disease (DFTD)-free Tasmanian devils (*Sarcophilus harrisii*) on Maria Island. Department of Primary Industries, Parks, Water and Environment Internal Document, Tasmania. Available at <http://www.parks.tas.gov.au/file.aspx?id=26885> [Accessed: 18 August 2016].
- Farrell, L. E., Roman, J., and Sunquist, M. E. (2000). Dietary separation of sympatric carnivores identified by molecular analysis of scats. *Molecular Ecology* **9**, 1583–1590. doi:10.1046/j.1365-294x.2000.01037.x
- Fischer, J., and Lindenmayer, D. B. (2000). An assessment of the published results of animal relocations. *Biological Conservation* **96**, 1–11. doi:10.1016/S0006-3207(00)00048-3
- Griffith, B., Scott, J. M., Carpenter, J. W., and Reed, C. (1989). Translocation as a species conservation tool: status and strategy. *Science* **245**, 477–480. doi:10.1126/science.245.4917.477
- Hamede, R. K., McCallum, H., and Jones, M. (2013). Biting injuries and transmission of Tasmanian devil facial tumour disease. *Journal of Animal Ecology* **82**, 182–190. doi:10.1111/j.1365-2656.2012.02025.x
- Hawkins, C. E., Baars, C., Hesterman, H., Hocking, G. J., Jones, M. E., Lazenby, B., and Mann, D. (2006). Emerging disease and population decline of an island endemic, the Tasmanian devil *Sarcophilus harrisii*. *Biological Conservation* **131**, 307–324. doi:10.1016/j.biocon.2006.04.010
- Hawkins, C. E., McCallum, H., Mooney, N., Jones, M., and Holdsworth, M. (2008). *Sarcophilus harrisii*. In 'IUCN 2014. IUCN Red List of Threatened Species. Version 2014.1'. Available at <http://www.iucnredlist.org/details/40540/0> [Accessed: 18 August 2015].
- Hope, J. H. (1972). Mammals of the Bass Strait Islands. *Proceedings of the Royal Society of Victoria* **85**, 163U195.
- Jones, M. E. (2001). Large marsupial carnivores. In 'The New Encyclopedia of Mammals'. (Ed. D. Macdonald.) pp. 814–817. (Oxford University Press: Oxford, UK.)
- Jones, K. E., Bielby, J., Cardillo, M., Fritz, S. A., O'Dell, J., Orme, C. D. L., Safi, K., Sechrest, W., Boakes, E. H., Carbone, C., and Connolly, C. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**, 2648. doi:10.1890/08-1494.1
- Jule, K. R., Leaver, L. A., and Lea, S. E. G. (2008). The effects of captive experience on reintroduction survival in carnivores: a review and analysis. *Biological Conservation* **141**, 355–363. doi:10.1016/j.biocon.2007.11.007
- Maran, T., Pödra, M., Pölma, M., and Macdonald, D. W. (2009). The survival of captive-born animals in restoration programmes: case study of the endangered European mink *Mustela lutreola*. *Biological Conservation* **142**, 1685–1692. doi:10.1016/j.biocon.2009.03.003
- Martin, T. G., Nally, S., Burbidge, A. A., Amall, S., Garnett, S. T., Hayward, M. W., Lumsden, L. F., Menkhurst, P., McDonald-Madden, E., and Possingham, H. P. (2012). Acting fast helps avoid extinction. *Conservation Letters* **5**, 274–280. doi:10.1111/j.1755-263X.2012.00239.x
- Mathews, F., Orros, M., McLaren, G., Gelling, M., and Foster, R. (2005). Keeping fit on the ark: assessing the suitability of captive-bred animals for release. *Biological Conservation* **121**, 569–577. doi:10.1016/j.biocon.2004.06.007
- McCallum, H., Jones, M., Hawkins, C., Hamede, R., Lachish, S., Sinn, D. L., Beeton, L., and Lazenby, B. (2009). Transmission dynamics of Tasmanian devil facial tumour disease may lead to disease induced extinction. *Ecology* **90**, 3379–3392. doi:10.1890/08-1763.1
- Moehrenschrager, A., and Somers, M. (2004). Canid reintroductions and metapopulation management. In 'Canids: Foxes, Wolves, Jackals, and Dogs: Status Survey and Conservation Action Plan'. (Ed. C. Sillero-Zubiri, M. Hoffman and D. W. Macdonald.) pp. 289–297. (IUCN/SSC Canid Specialist Group: Gland, Switzerland.)
- Owen, D., and Pemberton, D. (2005). 'The Tasmanian Devil: a Unique and Threatened Animal.' (Allen and Unwin: Sydney.)
- Pearse, A. M., and Swift, K. (2006). Transmission of devil facial-tumour disease. *Nature* **439**, 549. doi:10.1038/439549a
- Pemberton, D., and Renouf, D. (1993). A field-study of communication and social-behavior of the Tasmanian devil at feeding sites. *Australian Journal of Zoology* **41**, 507–526. doi:10.1071/ZO9930507
- Pemberton, D., Gales, S., Bauer, B., Gales, R., Lazenby, B., and Medlock, K. (2008). The diet of the Tasmanian devil, *Sarcophilus harrisii*, as determined from analysis of scat and stomach contents. *Papers and Proceedings of the Royal Society of Tasmania* **142**, 13–22.
- Rabin, L. A. (2003). Maintaining behavioural diversity in captivity for conservation: natural behaviour management. *Animal Welfare* **12**, 85–94.
- Simón, M. A., Arenas-Rojas, R., Rodríguez-Siles, J., García-Tardío, M., Perez-Marin, J., Gil-Sánchez, J. M., and Lopez, G. (2013). Re-introduction of the Iberian lynx, Andalusia, Spain. In 'Global Re-introduction Perspectives 2013. Further Case Studies from around the Globe'. (Ed. P. S. Soorae.) pp. 120–124. (IUCN/SSC Re-introduction Specialist Group Abu Dhabi and Environment Agency: Abu Dhabi.)
- Sinn, D. L., Cawthen, L., Jones, S. M., Pukk, C., and Jones, M. E. (2014). Boldness towards novelty and translocation success in captive-raised, orphaned Tasmanian devils. *Zoo Biology* **33**, 36–48. doi:10.1002/zoo.21108
- Snyder, N. F., Derrickson, S. R., Beissinger, S. R., Wiley, J. W., Smith, T. B., Toone, W. D., and Miller, B. (1996). Limitations of captive breeding in endangered species recovery. *Conservation Biology* **10**, 338–348. doi:10.1046/j.1523-1739.1996.10020338.x
- Soorae, P. S. (Ed.) (2013). 'Global Re-introduction Perspectives 2013. Further Case Studies from around the Globe.' (SSC Re-introduction Specialist Group and Abu Dhabi, UAE Environment Agency: Abu Dhabi.)
- Stoinski, T. S., Beck, B. B., Bloomsmith, M. A., and Maple, T. L. (2003). A behavioral comparison of captive-raised, reintroduced golden lion tamarins and their wild-born offspring. *Behaviour* **140**, 137–160. doi:10.1163/156853903321671479
- Tarsisz, E., Dickman, C. R., and Munn, A. J. (2014). Physiology in conservation translocations. *Conservation Physiology* **2**, doi:10.1093/conphys/cou054
- Tucker, M. A., and Rogers, T. L. (2014a). Examining the prey mass of terrestrial and aquatic carnivorous mammals: minimum, maximum and range. *PLoS One* **9**, e106402. doi:10.1371/journal.pone.0106402
- Tucker, M. A., and Rogers, T. L. (2014b). Examining predator-prey body size, trophic level and body mass across marine and terrestrial mammals. *Proceedings of the Royal Society B: Biological Sciences* **281**, 2014–2103. doi:10.1098/rspb.2014.2103

- van Heezik, Y., and Ostrowski, S. (2001). Conservation breeding for reintroductions: assessing survival in a captive flock of *houbara bustards*. *Animal Conservation* **4**, 195–201. doi:[10.1017/S1367943001001238](https://doi.org/10.1017/S1367943001001238)
- Vickery, S. S., and Mason, G. J. (2003). Behavioural persistence and its implication for reintroduction success. *Ursus* **14**, 35–46.
- Whittaker, R. H. (1972). Evolution and measurement of species diversity. *Taxon* **21**, 213–251. doi:[10.2307/1218190](https://doi.org/10.2307/1218190)
- Wolf, C. M., Griffith, B., Reed, C., and Temple, S. (1996). Avian and mammalian translocations: update and reanalysis of 1987 survey data. *Conservation Biology* **10**, 1142–1154. doi:[10.1046/j.1523-1739.1996.10041142.x](https://doi.org/10.1046/j.1523-1739.1996.10041142.x)