

[10.1071/BT22076](https://doi.org/10.1071/BT22076)

Australian Journal of Botany

Supplementary Material

Seed dormancy alleviation by warm stratification progressively widens the germination window in Mediterranean climate Rutaceae

Michael Just^{A,}, Adam T. Cross^{A,B}, Wolfgang Lewandrowski^{C,D}, Shane R. Turner^A, David J. Merritt^{C,D}, and Kingsley Dixon^A*

^ASchool of Molecular and Life Science, Curtin University, Kent Street Bentley, WA 6102, Australia.

^BEcoHealth Network, 1330 Beacon St, Suite 355a, Brookline, MA 02446, USA.


^CKings Park Science, Department of Biodiversity, Conservation and Attractions, 2 Kattidj Close, Kings Park, WA 6005, Australia.

^DSchool of Biological Sciences, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

*Correspondence to: Michael Just School of Molecular and Life Science, Curtin University, Kent Street Bentley, WA 6102, Australia Email: michael.just@curtin.edu.au

Supplementary Table S1 Akaike information criterion obtained from the AIC function for each model applied to each parameter. Model with the lowest value was selected as the best fit.

Parameter	Model	df	AIC
Germination proportion	Beta	31	-154.6
Germination proportion	Broken stick	25	-113.5
Germination proportion	Yan and hunt	19	-166.4



Supplementary Figure S1 Germination proportion ($\pm 95\%$ confidence interval) of *Rhadiotothamnus anceps* seeds incubated at 5, 10, 15 or 20°C following warm stratification at either 20 or 25°C for 0, 1, 2, 4, 8 and 12 week.