

[10.1071/CP22255](https://doi.org/10.1071/CP22255)

Crop & Pasture Science

Supplementary Material

Seed priming and soil application of zinc decrease grain cadmium accumulation in standard and zinc-biofortified wheat cultivars

Ayta Umar^A, and Shahid Hussain^{A,}*

^ADepartment of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan 60800, Pakistan.

*Correspondence to: Shahid Hussain Department of Soil Science, Faculty of Agricultural Sciences and Technology, Bahauddin Zakariya University, Multan 60800, Pakistan Email: shahid.hussain@bzu.edu.pk

Supplementary Table S1. Post-harvest soil DTPA extractable Zn and Cd

Main effect	Levels	Soil DTPA extractable Zn (mg kg ⁻¹)	Soil DTPA extractable Cd (mg kg ⁻¹)
Cultivars	Jauhar-2016	0.73±0.12 A	0.14±0.01 A
	Zincol-2016	0.74±0.12 A	0.13±0.01 A
Zinc treatments	Control (un-primed)	0.63±0.02 B	0.13±0.01 A
	Hydro-primed	0.62±0.02 B	0.14±0.01 A
	Zinc-primed	0.61±0.02 B	0.13±0.01 A
	Soil Zn (un-primed)	0.84±0.02 A	0.14±0.01 A
	Hydro-primed + soil Zn	0.83±0.03 A	0.13±0.01 A
	Zinc-primed + soil Zn	0.85±0.03 A	0.13±0.01 A

Separately for each parameter, different letters after means ± s.d. of each parameter indicate significant differences based on Tukey's HSD test ($P \leq 0.05$).