

Bernhard Hermann Neumann 1909–2002

Cheryl E. Praeger

School of Mathematics and Statistics, University of Western Australia, WA 6009, Australia.

Email: cheryl.praeger@uwa.edu.au


Bernhard Hermann Neumann was born and educated in Berlin. He held doctorates from Berlin and Cambridge, and mathematical positions at universities in Cardiff, Hull, Manchester, and the Australian National University (ANU) in Canberra. Whereas his move to the UK in 1933 was a result of the difficulties he faced as a Jew in finding employment in Germany, his move to Australia in 1962 was to set up a new research Department of Mathematics at the Institute of Advanced Studies at the ANU. Bernhard Neumann was famous for both his seminal research work in algebra and his strong support of all endeavours in mathematics. His scholarly publications span more than seventy years. His honours include Fellowship of the Royal Society and of the Australian Academy of Science, appointment as Companion of the Order of Australia, and numerous honorary doctorates. To Bernhard it was important to share and spread the joy of doing mathematics.

Bernhard Hermann Neumann was born in Berlin, Germany, on 15 October 1909. After completing his doctorate in Berlin, he moved to the UK to seek employment, and undertook a second doctorate in Cambridge. His academic work as a mathematician in the UK was interrupted by a period of internment, and then service in the British army, during the Second World War. Bernhard Neumann was head-hunted from his position in Manchester to become, in 1962, foundation professor and head of a new research mathematics department at the Australian National University. He had an enormous positive influence on the development of mathematics in Australia and the Asia-Pacific region, not only during his time as departmental head at the ANU but also in the decades following his retirement in 1975. Bernhard died suddenly on 21 October 2002, in Canberra, Australia, shortly after his 93rd birthday. He is survived by his second wife Dorothea and the five children of his first marriage (to Hanna, also a mathematician and a Fellow of the Australian Academy of Science at the time of her death in 1971).

Early Life and Education

Bernhard was named after his two grandfathers. His paternal grandfather Bernhard Neumann was a hardware merchant in Karlsruhe in the south of Germany, and Bernhard's father Richard was the youngest of four children, born on 15 January 1876. Bernhard's maternal grandfather Hermann Aronstein had a large farm, called a *Rittergut*, in Westphalia, and Bernhard's mother Else was Hermann's third daughter. Both sides of Bernhard Neumann's family were thus solidly middle-class.

Richard Neumann married Else in 1905 in Berlin. He studied engineering, spent two years in the USA as an engineer and then held a job until the late 1930s in Berlin, in the AEG—Allgemeine Elektrizitätsgesellschaft. Bernhard was the second child and only son. His elder sister Edel-Agathe (known as Eta) was born in 1906 in Berlin, studied physics in Berlin, worked in patent law and patent physics in north-east England until she married Gösta Lindskog, a Swedish pastor and school teacher, and then lived in Sweden with Gösta until her death from cancer in 1958. Both of Bernhard's



parents were Jewish. Richard was not practising, while Else went to synagogue for the important festivals. Bernhard was also not practising, but despite this he and his first wife Hanna sat on the Council of Jews and Christians in their Manchester days in the late 1950s, wishing to bring reconciliation between the two religions.

Bernhard was four years old when the First World War began. The blockade against Germany caused severe shortages and strict rationing, and Bernhard remembered feeling very hungry for much of the war. Thus he knew privation of a type not often met with in Europe today. Bernhard remembered with some affection the arrival of Quakers immediately after the war; they went around schools in Germany feeding the children.

Bernhard attended the *Herderschule* in Berlin-Charlottenburg from 1916 to 1928. He claims that he was dull as a young child to the extent that an aunt consoled his mother with the words, ‘he might yet make an artisan of sorts!!’¹ A tonsillectomy that happened at about the time he entered the *Herderschule* changed this perception. From then on he took delight in sums and particularly mental arithmetic. Bernhard’s father had a student text on calculus that fascinated him. He did all the exercises and thus learned differential calculus (then a university subject) at a very early age, and loved studying the beautiful curves contained in the book. Later, in Year 10 of school, he invented three-dimensional analytic geometry for himself.

At school, Felix W. Behrend taught Bernhard mathematics and physics. His teacher’s son Felix A. Behrend, a life-long friend of Bernhard, became an Associate Professor of Mathematics at the University of Melbourne (see B73). The *Herderschule* also produced G. P. Hochschild, later to become

Professor of Mathematics at the University of California, Berkeley. Throughout his life, Bernhard made friends quickly and easily, and kept in contact with a large number of people. His remarkable memory was no doubt aided by his famous pocket diary in which he recorded birthdays and other significant details.

Bernhard left the *Herderschule* with the *Abitur*, the qualification that was an entitlement to university entrance. His university education began with two semesters at Freiburg University in 1928–29, studying under Alfred Loewy, Gustav Mie and Lothar Heffter. Heffter, Bernhard’s favourite teacher, was mathematically active until his 100th year. Bernhard dedicated his paper B68 for Heffter’s 100th birthday. However on submitting the paper Bernhard learned from the editor ‘by return’ that Heffter had just died, less than six months before his 100th birthday (B120, p. 587). So the paper is dedicated to his memory. Bernhard was always keenly interested in people who continued to be very active academically after retirement, and later loved the fact that his own life was so long. When a mutual friend died, Bernhard is reported as saying, ‘but he was only 85, wasn’t he?’ The other side of the coin was that he became one of the great supporters of the young, as we see later.

After the Freiburg experience, Bernhard returned home to Berlin to study at the Friedrich-Wilhelms-Universität, later to become the Humboldt University. He was delighted to have a degree from both incarnations of the institution, the doctorate from Humboldt being an honorary one. Bernhard studied under such luminaries as Erhard Schmidt, Issai Schur, Robert Remak and Heinz Hopf. He began with pure mathematics, physics and philosophy. Later he concentrated on pure mathematics.

Bernhard was introduced to group theory in the autumn semester of the academic year 1930–31 by Remak, who had given a course on his own (then unpublished) work

¹ Much of the material for this part of the memoir has been drawn from the record of interview [1] and the essays written by Bernhard in the *Selected Works* of B. H. Neumann and Hanna Neumann B120.

on direct products of groups. Apparently the attendance started at thirty and ended up with Bernhard as the single attendee. Groups were new and exciting to him, and this excitement never wavered throughout his life. He came across the subject that eventually became part of his Dr.phil. dissertation while attending a seminar on geometry and topology led by Hopf and Georg Feigl.

Bernhard had read a paper of Jakob Nielsen on automorphisms of free groups (B120, p. 1); Nielsen had found a finite presentation for the automorphism group of a free group of finite rank. Bernhard discovered that he could reduce the number of generators and relations. Indeed he found the minimum generating numbers for free groups of rank at least 4. He told Hopf, who suggested that the proof should be written down; and when Hopf saw the resulting manuscript, he suggested that it might be used as Bernhard's doctoral dissertation. Bernhard thought that he was too young and the result too slim, but after comings and goings involving Hopf, Schur and Schur's assistant Alfred Brauer, Schur eventually agreed that it was perhaps a little on the slim side. Schur suggested that Bernhard investigate the minimum generating numbers of what are now called permutational wreath products of finitely presented groups and symmetric groups. In a fortnight, Bernhard had done the required work—'the size doubled' (B120, p. 1)—and now there was enough for the Dr.phil., which Bernhard took at the early age of 22. Bernhard's first published paper B1 came out of this.

Life in England up to 1946

Bernhard left Germany for England in August 1933, and on his arrival was given refugee status, somewhat to his surprise. He found that job prospects in England were no better than they had been in Germany, so he 'camouflaged' the fact that he was unemployed. He became a research student again, at Fitzwilliam House (now

Fitzwilliam College), Cambridge, financed at first by his parents in Germany and then by an uncle in South Africa. His supervisor was Philip Hall and weekly supervision sessions took the form of dinner in King's College followed by discussion in Hall's rooms in college, 'where we would drink his sherry, smoke his cigarettes and talk about rhizomorphs of plants and political history in Germany and everything'. At about ten o'clock the discussion would drift towards mathematics. After completing his PhD in 1935, Bernhard stayed on in Cambridge for a further two years until he was appointed in late 1937 as a temporary assistant lecturer in mathematics at University College, Cardiff. He taught in Cardiff until 1940.

Bernhard had met Hanna von Caemmerer in January 1933. Hanna was born in Berlin on 12 February 1914, the youngest of three children of Hermann and Katharina von Caemmerer. Her father had a doctorate in history, and was well on the way to establishing himself as an archivist and academic historian when he was killed in the first days of the 1914–18 war. As a result she and her mother, brother Ernst (born 1908) and sister Dorothea (Dora, born 1910) lived impecuniously on a war pension and Hanna contributed to the family income by coaching younger school students for up to fifteen periods a week.

Bernhard and Hanna got to know each other through the Mathematisch-Physikalische Arbeitsgemeinschaft, a working group within the university in Berlin. Although Hanna was not Jewish, she was very much anti-Nazi. Bernhard and Hanna remained in regular contact by correspondence after Bernhard moved to England. As open correspondence was too dangerous, their letters went secretly via various friendly channels. They became engaged in 1934 and met for a couple of weeks in Denmark in 1936 as Bernhard was returning to England from the International Congress of Mathematicians in Oslo. Then in 1938 Hanna, deciding that war was not unlikely,

moved to the UK where she and Bernhard were married. However such ‘mixed’ marriages between Jews and non-Jews were against the law in Germany, and Bernhard and Hanna felt they could not openly marry until his parents were safe from possible reprisals. Thus after their marriage Hanna lived in Bristol and Bernhard in Cardiff. Bernhard’s father could not believe that the level of evil abroad in Germany since 1933 could possibly last, and it was not until February 1939 that Bernhard’s parents left Germany. From that time, the four Neumanns lived together in a small house in Cardiff, where Bernhard’s and Hanna’s first child Irene was born in August 1939, just before the beginning of the Second World War.

During the first part of 1940, Bernhard and Hanna fell into the class of ‘least restricted aliens’ and Bernhard continued to work as a lecturer in Cardiff as this was a ‘reserved occupation’ not liable for army service. However, after the Dunkirk evacuation a large part of the coast was barred to all aliens and they were required to leave Cardiff. They moved to Oxford, and within a week Bernhard was interned. After several months he was released into the British army. He was trained in Yorkshire and then transferred to the south of England. Most of the time he was within reach of Oxford and could visit his family occasionally. In particular, he obtained special leave when his son Peter was born in Oxford at the end of December 1940.

Bernhard served in the Pioneer Corps until 1943, when he was allowed to volunteer for combatant service. He was transferred to the Royal Artillery, then to the Artillery Survey where he used theodolites and did some numerical work. Later he entered the Intelligence Corps. During this time Bernhard completed several mathematical papers (B14, B15, B16, see B120 pp. 279 and 740).

Meanwhile Hanna submitted her Dr.phil. thesis in Oxford in mid-1943, and soon

afterwards she was allowed to return to Cardiff, where their third child Barbara was born in November 1943. Hanna returned to Oxford for her oral examination in April 1944 and Bernhard, by then in the Intelligence Corps and stationed not very far away, was able to join her for a celebratory lunch after the examination.

During 1945, when the European war was over, Bernhard volunteered to go to Germany with the Intelligence Corps. On a long weekend leave (replacing his leave for VJ Day that he had missed) he was able to visit Hanna’s sister and mother. He was demobilized later that year and returned to England.

Life in England 1946–1962

The University College in Cardiff wanted Bernhard to return to his teaching post there, but Bernhard did not feel obliged to do so because they had not helped to get him out of internment. Instead he obtained a Temporary Lectureship at the University College in Hull (now the University of Hull) at the beginning of 1946, as replacement for Jacob Bronowski. At the same time his fourth child Walter was born. For the next academic year, Bernhard was appointed to a permanent lectureship and, as Bronowski decided not to return to Hull, Hanna applied for and was appointed to the vacant position as a Temporary Assistant Lecturer.

Hanna stayed in Hull for twelve years, but in 1948 Bernhard was enticed to a position at the University of Manchester by M. H. A. Newman. This entailed ten years of commuting between Manchester and Hull in all the vacations and for many weekends. Bernhard was not only a keen everyday cyclist, but also a long-distance one: on several occasions he made the hundred-mile trip across the Pennines between Manchester and Hull by bicycle, and once he rode from Manchester to give a lecture in London because there was a train strike. In 1951, during this period, his fifth child Daniel was born.

This period saw also the publication of a number of highly influential research papers. Foremost is the famous joint paper B19 (1949) of Bernhard and Hanna with Graham Higman introducing HNN-extensions. In the same year Bernhard's paper B17 on ordered groups appeared. Published shortly afterwards were the paper B20 (1950) in which he answered one of the 1948 Prijsvraagen of the Wiskundig Genootschap te Amsterdam, and Bernhard's celebrated essay B38 for which he won the Adams Prize in 1952. Then in 1954 his papers B36 and B41 on BFC groups appeared (see Section F.5).

In 1958 Hanna obtained a Lectureship in the Faculty of Technology at the Manchester College of Science and Technology (now, after several name changes, part of the University of Manchester). At the time the College was part of the University of Manchester but independently financed and physically separate, with its own department of mathematics. Hanna's role was to set up an honours programme in mathematics.

By then Bernhard had the rank of Reader in Mathematics, and in the following year, 1959, he was elected to the Fellowship of the Royal Society for his numerous and influential contributions to the theory of infinite groups. Bernhard served as a Vice-President of the London Mathematical Society 1957–59 and as an Editor of the *Proceedings of the London Mathematical Society* 1959–61. Bernhard supervised eight PhD students in Manchester, seven of whom became long-term university teachers and researchers in mathematics and the other a senior secondary teacher of mathematics and mathematics editor. Six of them became professors of mathematics: in Egypt, England, Wales, the USA and two in Australia.

During 1959 Bernhard was granted sabbatical leave from Manchester. He spent three months visiting all the universities in Australia, including a week at the Australian

National University in Canberra. Following his Australian tour he spent the 'Monsoon Term' of 1959 at the Tata Institute of Fundamental Research in Bombay, and notes of his lectures in Bombay were subsequently published by the Tata Institute B61. These travels are perhaps an early manifestation of his combining travelling activities with the spreading of mathematics.

Bernhard held the position in Manchester until 1962, but the final academic year was spent on study leave working at the Courant Institute of Mathematical Sciences in New York. The year in New York was fruitful mathematically, for Bernhard and also for Hanna and their son Peter who was by then an undergraduate student in mathematics in Oxford. They wrote the '3N' paper B67 on product varieties of groups and embarked on a joint project with Gilbert Baumslag, one of Bernhard's former PhD students, which culminated a year later in the 'B+3N' paper B75 on varieties generated by a finitely generated group.

The Careers of Bernhard's Five Children

After completing a Master of Arts in English literature at the University of Manchester, Irene (Neumann Brown) was a high school teacher in Aberdeen, Scotland, and then a lecturer in English at New Mexico State University. Peter became a mathematician at the Queen's College, Oxford, with research contributions ranging over a number of areas of algebra and its history. He was awarded the Lester R. Ford Award by the Mathematical Association of America in 1987 and the Senior Whitehead Prize by the London Mathematical Society in 2003, and was appointed an Officer in the Order of the British Empire in 2008 for services to education. On completing a mathematics degree at the University of Sussex, Barbara (Cullingworth) taught at Saint Bernard's Convent School in Slough, England, specializing in the teaching of statistics, and



Figure 1. Bernhard Neumann and his first wife Hanna, around the time of Hanna's appointment as professor at the Australian National University, Canberra, 1964.

having an active role in the Mathematical Association. Walter studied mathematics in Adelaide and Bonn. His mathematical interests are in topology, algebraic geometry and geometric group theory, and his academic career has been an international one, with posts in Germany, Maryland, Ohio, Melbourne and New York. He is presently Full Professor at Barnard College, Columbia University, New York. Daniel completed degrees in ancient Greek and pure mathematics at Monash University, and in psychology at the University of New England and Swinburne University. For many years he was a professional musician in Orchestra Victoria, and he now works as a psychologist.

Life in Australia 1962–1974

As a result of Bernhard's Australian visit in 1959, and with some strong encouragement especially from Sir Mark Oliphant, Bernhard and Hanna in 1961 accepted positions at the Australian National University (ANU), Bernhard as the foundation Professor of Mathematics at the Institute of Advanced Studies, a position

he held until his retirement at the end of 1974, and Hanna as Reader in the department Bernhard was appointed to set up. Later Hanna became the first female professor of mathematics in Australia, and Foundation Head of the Department of Pure Mathematics in the School of General Studies at the ANU (see Fig. 1). She was also the first woman mathematician elected a fellow of the Australian Academy of Science (in 1969). Sadly, she died in 1971 while on a lecture tour in Canada.

Bernhard's major task on taking up his position at ANU in 1962 was to build up the new Department of Mathematics, and to establish a healthy PhD programme in mathematics there. He attracted active researchers to his new department. During Bernhard's headship (1962–74) about fifty graduate students completed their degrees, of whom eight were directly supervised by him. He always had an open door and fostered a feeling of family. The students sometimes found his expectation of intellectual rigour daunting but came to appreciate it. They have gone on in diverse ways—quite a few of them made their careers in



Figure 2. Bernhard Neumann and his second wife Dorothea, photographed in Canberra in 1974 by Professor Kurt Mahler FRS FAA.

Australian universities, and Bernhard lived to see many of them make a significant mark.

Bernhard married Dorothea Frieda Auguste Zeim in 1973 (see Fig. 2). Dorothea was born in Berlin on 20 January 1939. Her father Albert Zeim was a classics scholar and secondary school teacher. Her mother Hildegard Rockstroh studied modern languages (French and English) but did not finish her doctorate for family reasons. She started working, as a contract teacher at secondary school, in the 1960s when her three children were all at university. Dorothea had known the Neumann family for many years. Her mother had gone to school with Hanna's elder sister Dora, and Dorothea had spent some time with the Neumann family in Manchester working as an *au pair* and becoming familiar with English as part of her education. She had remained in contact with the family from that time. After her marriage Dorothea completed a PhD in linguistics at the Australian National University.

Bernhard travelled widely, giving many talks, attending many conferences and

spreading the message that good mathematics was being done in Australia and that Australia was a good place in which to do mathematics.

Bernhard expected intellectual rigour not only in mathematics but more widely. He was known for his rigour in other activities such as editing typescripts and recording of minutes of meetings.

The first two international conferences in Australia on a mathematical topic were held under his leadership, on the theory of groups, in 1965 and 1973. Both were notable for the quality of the main speakers and the range of countries from which they came. The first was also notable because Bernhard was able to arrange for young people from overseas to earn their way to Australia by teaching at an Australian university. A third international conference on the theory of groups was held in 1989 to mark his 80th birthday.

Bernhard was active in many aspects of mathematical life in Australia, holding numerous leadership positions that were later appropriately acknowledged. He became the fifth President of the

Australian Mathematical Society (1964–66), was elected an honorary member in 1981, and was further honoured by having a prize named after him for the most outstanding talk by a student at the Annual Meeting of the Society. Two of his mathematical grandchildren (doctoral students of his own PhD students) have won that prize. As founding editor (1969–79) of the *Bulletin of the Australian Mathematical Society*, which he developed as a quality international journal, Bernhard succeeded in creating a vehicle for fast publication of good mathematics, though he did not quite achieve his hoped-for five months, on average, from submission to publication. Bernhard also helped found the Australian Association of Mathematics Teachers and was its first President (1966–68).

Bernhard was elected to the fellowship of the Australian Academy of Science in 1964. The citation noted his contributions to group theory and that he had established himself as one of the leaders of mathematics in the country, and his determination to foster mathematics on a national scale. He served on Council (1968–71), was a Vice-President (1969–71), and gave the Matthew Flinders Lecture in 1984. He served an extended term on the National Committee for Mathematics (1963–75) and on Australian delegations to many meetings of the International Mathematical Union (IMU), held in conjunction with International Congresses of Mathematicians (he attended thirteen of these congresses; surely close to a record). He gave an invited lecture at the Nice congress in 1970. Also, at the Nice congress, he was appointed to improve communication among mathematicians. This led to his founding the *IMU Canberra Circular*, which he edited almost single-handedly from 1972 to 1999; it provided timely information about mathematical meetings as well as announcements of honours and deaths within the mathematical community internationally. Its circulation

rose to more than 1100 before becoming largely electronic, and was especially valued by colleagues in countries with little contact with the wider world. Bernhard also served (1975–79) on the Exchange Commission of the IMU.

Through the Australian Academy of Science, Bernhard initiated the Australian Subcommittee of the International Commission on Mathematical Instruction (ICMI), chaired it (1968–75) and was the Australian representative on ICMI. He was a member at large of ICMI (1975–82) and of its Executive Committee (1979–82). This was the basis for Australia's hosting of the Fifth International Congress on Mathematical Education (ICME5) in Adelaide in 1984. In addition he was active in getting the Academy involved in providing materials for schools, and after a long gestation period, six volumes of *Mathematics at Work* appeared in 1980–81.

Life in Australia 1974–2002

On retiring as Professor and Head of the Department of Mathematics at the end of 1974, Bernhard was made Professor Emeritus and an Honorary Fellow of ANU. He was also appointed a Senior Research Fellow at the Commonwealth Scientific and Industrial Research Organisation (CSIRO) for three years and then became an Honorary Research Fellow, reappointed annually until his death. In 1975 he also became an Honorary Member of the Canberra Mathematical Association, the Australian Association of Mathematics Teachers, and the New Zealand Mathematical Society.

In 'retirement' he continued his own research in mathematics and his support of work in mathematics by others, directly, through teaching and editorial and committee work. For example, he served as a member of the Academic Advisory Council of the Royal Australian Naval College. He continued to be an ambassador for mathematics and for Australia.




Figure 3. Bernhard Neumann cycling from his CSIRO office located in the Division of Forest Research, Yarralumla, ACT. Photographed in 1978 by Alan G. Edward AIAP. Image courtesy of CSIRO.

Together Bernhard and Dorothea provided a steady welcoming and supportive environment especially for visitors and young people, in mathematics, in music and quite generally. The annual Neumann wine and cheese parties were legendary. Initially they were yearly gatherings of Bernhard's and Hanna's students. By the 1970s the guest list included at least 100 students and former students, together with their friends and family.

Dorothea gave Bernhard outstanding non-mathematical support by joining him in exploring and pursuing activities outside mathematics. In particular, she joined him in regular chamber music evenings and in an orchestra in which Bernhard played the cello, and strongly encouraged him to continue. Bernhard's approach to new challenges and suggestions was always

'why not'. After his son Daniel and grandchildren took him camping for the first time when he was over seventy, Dorothea and he planned and executed numerous camping expeditions that sometimes included mathematics conferences. Dorothea developed ingenious plans to enable Bernhard to keep cycling up to a few months before his death (see Fig. 3), and to continue his active enjoyment of the Australian bush. On his eightieth birthday he went ballooning; and on his ninetieth he went abseiling. One cannot help but wonder what he would have done had he made it to his hundredth birthday!

Bernhard continued to support activities aimed at stimulating and developing mathematics talent. He gave considerable encouragement to Peter O'Halloran and his colleagues involved in the formative stages of what is now the Australian Mathematics Competition and maintained an active interest in it. He chaired the Australian Mathematical Olympiad Committee from its inception in 1980 till 1986. In his role as Chair of a Site Committee (1981–83) during his term on the Executive Committee of ICMI (1979–82), he ensured better structure and operation of the International Mathematical Olympiads. The holding of the 1988 IMO in Australia during Australia's bicentennial year owed much to him. Through the B. H. Neumann Awards, the Australian Mathematics Trust recognizes significant and sustained contributions to the enrichment of mathematics learning in Australia and its region. When the Australian Mathematics Trust commissioned the Sydney portrait artist Judy Cassab to paint his portrait, Bernhard was delighted not only by the painting but also by Judy Cassab's approach to portrait painting: 'she talked all the time and asked me questions, and I talked all the time—and so the portrait is not a photograph, it's myself'.

Most of his mathematical life was spent in Australia. During that time he contributed much to mathematics in Australia and became a much loved and

respected figure. He was a positive influence on an immense number of people. He was made a Companion of the Order of Australia in 1994 for service to the advancement of research and teaching in mathematics.

He was awarded honorary doctorates from a number of universities: the University of Newcastle (NSW), Monash University (Vic), the University of Western Australia and the ANU within Australia; and the University of Hull (UK), the University of Waterloo (Canada) and the Humboldt University of Berlin (Germany). Bernhard was well known in circles outside mathematics. On his first visit to Australia he played in many chess clubs. At the time of his death he was the oldest rated chess player in the country. He was known to the broader Canberra public as a familiar figure cycling on its roads, in more recent times wearing an electric blue helmet. He was for many years Vice-President of the Friends of the Canberra School of Music; he helped judge an annual chamber music competition the day before he died. He enjoyed exploring the countryside and showing visitors around it. In spite of all this visible activity, he was perhaps at his best giving quiet, often unnoticed and unrecognized help to individuals.

Bernhard Neumann was a strong supporter of all endeavours in mathematics—he supported people who did mathematics for its own sake, people who applied mathematics and people who taught mathematics. To him it was important to share and spread the joy of doing mathematics. Bernhard Neumann was the right person at the time for mathematics in Australia with his energy, enthusiasm and commitment to the subject and to all people involved in mathematics. During his lifetime, Bernhard saw Australia change from a mathematically under-developed country to one with a significant mathematical profile, and he was instrumental in that change.

Scientific Work

The scientific work of Bernhard Neumann ranges widely in mathematics. The core is his published research. In addition he made many other contributions through his editorial work, his lectures, his reviews (of which he wrote many hundreds), his training of students, his participation in and organization of conferences and his ready availability to all who sought him out.

The published research of Bernhard Neumann is primarily in algebra with occasional excursions into geometry and other parts of mathematics. Within algebra his main work was on the Theory of Groups, though there was often work in more general contexts such as division rings, near-rings, semigroups, universal algebra and ordered algebras. Groups are mathematical structures satisfying a small number of axioms that formalize the essential aspects of symmetry. Thus one may view the focus of Bernhard Neumann's research as the 'Mathematical Science of Symmetry'.

Much of Bernhard Neumann's published work continues to be cited regularly up to the present: most notably the HNN-embedding paper B19 and the 'Mal'cev-Neumann construction' B18 for ordered division rings, both dating from 1949; his seminal paper B6 on infinite groups and his paper B9 on varieties of groups, both published in 1937; his essay on group products with amalgamations B38; and his papers on group coverings B36, B39, on various finiteness conditions for groups B24, B42 and on the 'Erdős Problem' B104. Other of his 'mathematical inventions', such as the twisted wreath product, are now household names in mathematics and are routinely used without explicit reference.

Bernhard Neumann was punctilious about publication names. For example, in 1970, when I was writing my first research article arising from an undergraduate summer vacation research project supervised by

Bernhard, he counselled me about keeping one publication name throughout my career. Moreover that name, he said, should as far as possible uniquely identify me as author. Thus I became Cheryl E. Praeger and, as Bernhard explained, he was from 1935 onwards B. H. Neumann in print. Therefore, in this scientific evaluation, I will refer to Bernhard as BHN, a name used by his colleagues when discussing Bernhard as mathematician.

Moreover, as a tribute to BHN, the numbering system used for listing his publications in the Bibliography and citing them in the text is that used by BHN himself in the *Selected Works* (B120, 1988). That is to say, publication number xy in the list is referenced as Bxy , both there and in the text. In particular this overcomes the problem of distinguishing the publications of BHN from those in the list of other publications. Moreover use of other references has been kept to a minimum. Where possible, such references have been restricted to material that can be readily traced using the mathematical reviewing journals *Mathematical Reviews* [2] and *Jahrbuch über die Fortschritte der Mathematik* [3], or the bibliography of the monograph by Chandler and Magnus [4] (from which one may gain a good view of BHN's contribution to combinatorial group theory). Publications reviewed by *Mathematical Reviews* or the *Jahrbuch* will be referenced in the text by their review number and year of publication. As examples of this usage, the history by Chandler and Magnus [4] might be referenced as Chandler and Magnus (MR0680777, 1982), and the paper of Burnside mentioned in Section C.1 as Burnside (JFM33.0149.01, 1902).

Within Group Theory, a number of topics are specifically associated with BHN. He is noted for the inventiveness of his constructions and his analysis of constructions and their application to embedding problems. He is a founder of the theory of varieties of groups.

A number of his contributions have seen his name used (in conjunction with others) and these are discussed in the indicated section:

the Birkhoff–Neumann embedding	Section E.1
the Douglas–Neumann Theorem	Section D.3
the Grushko–Neumann Theorem	Section F.3
the HNN construction/embedding	Section E.2
the Macintyre–Neumann Theorem	Section F.6
the Mal'cev–Neumann construction	Section E.5
the Moser–Neumann question	Section D.2
the Neumann–Rado Theorem	Section D.1
the Neumann semidirect product	Section E.3

A Automorphisms and Endomorphisms

A.1 Automorphism Groups of Free Groups

As mentioned already, BHN's interest in groups was aroused by a course in Berlin given by Remak in 1930. While he was participating in a seminar on geometry and topology conducted by Feigl and Hopf he came across a paper by Nielsen (MR1512188, 1924) in which finite presentations for the automorphism groups of finitely generated free groups are determined. BHN was able to reduce these presentations. In particular he showed that the automorphism groups of the free groups of rank at least 4 can all be generated by two elements. This was the core of his doctoral dissertation (1931) in Berlin and of his first paper B1 (1932). The heart of the work is manipulation of group presentations. This is something which continued to be a significant part of his work for his whole career. Presentations for the automorphism class groups of the finitely generated

free groups are determined in paper B23 (1951) with Hanna. The main thrust of that paper is characteristic subgroups and systems of generating vectors. Characteristic subgroups are also studied in B82 (1966). The theme is extended to general categories in B89 (1969).

A.2 Fixed-point-free Automorphisms

BHN reported in the *Selected Works* (B120, introductions to Chapters 2 and 9) that questions in the foundations of geometry had a revival of interest in the 1920s and 1930s arising from work on the 7th edition of Hilbert's *Grundlagen der Geometrie*. BHN also reported that much of the work he did on these foundational questions was never published. BHN was fascinated by the algebra underlying geometrical systems. In B11 (1940) he studied what are now called near-rings, generalizing results of Zassenhaus about finite near-fields to the infinite case. The questions were boiled down to proving that a group satisfying certain conditions is abelian. One of these conditions is the existence of an automorphism with order 2, which leaves fixed only the identity element; nowadays usually referred to as a fixed-point-free automorphism. BHN proved that a group which admits a fixed-point-free automorphism with order 2 and in which every element has a unique square root is abelian. In B14 (1943) he showed that the existence, but not uniqueness, of square roots is not sufficient. In B44 (1956) BHN gave a description of finite groups which admit a fixed-point-free automorphism with order 3. The study of groups and other algebraic systems admitting automorphisms with few fixed points has developed into a substantial topic; see, for example, monographs by Khukhro (MR1224233, 1993; MR1615819, 1998).

A.3 Finite Groups

In B43, B45 (1956) Ledermann and BHN studied automorphism groups of finite

groups. They were inspired by a problem of Birkhoff and Hall (MR1501860, 1936) which asked for a lower bound for the orders of automorphism groups of finite groups. In B43 Ledermann and BHN showed that large finite groups have large automorphism groups or more precisely: there is a bound $f(n)$ such that every finite group with order not less than $f(n)$ has at least n automorphisms. Recent work of Bray and Wilson (MR2243245, 2006) sheds further light on this. In B45 Ledermann and BHN considered what they called a local version and showed that, for each prime p , there is a function $g(h)$ such that every group whose order is divisible by $p^{g(h)}$ has p^h dividing the order of its automorphism group. An important ingredient in the work on the local problem is Schur's multiplier. Not long after, Green (MR0081901, 1956) made improvements that show that the function $h(h+3)/2+1$ suffices. There have been some further improvements in the bound but not asymptotically. Nevertheless there is a popular conjecture, seemingly dating from about the same time, that the order of a non-abelian group with prime-power order divides the order of its automorphism group (Schenkman MR0067111, 1955; Mann MR1716701, 1999).

A.4 Semigroups

The story is quite different in semigroups as Dlab and BHN showed in B91 (1969). They exhibited arbitrarily large finite semigroups with just eight endomorphisms, and their question of whether a smaller number can be achieved was answered by Kublanovsky (MR734514, 1983) who showed that for semigroups containing at least four elements, the minimum possible number of endomorphisms is four, and there are infinitely many finite semigroups with exactly four endomorphisms. Dlab and BHN also exhibited an infinite rigid semigroup: one whose only endomorphism is the identity.

B Presentation Manipulation

Presentation manipulation is a significant feature of BHN's work. It is the core of his paper B1, and in particular is a critical ingredient of B6 (1937). About the paper B6, Baumslag and Miller (MR2554768, 2009) say:

In [this] brief, remarkable paper [BHN] established several facts which are now widely known and used in combinatorial group theory: (i) a proper subgroup of a finitely generated group is contained in a maximal proper subgroup; (ii) if a group can be defined by finitely many relations on one finite set of generators, then it can be defined by finitely many relations on any other finite set of generators; (iii) there exist finitely generated groups which are not finitely presented; and (iv) there exists [by construction] a family of continuously many non-isomorphic two generator groups.

This important construction is discussed further in Section E.1.

Part of his joy with presentation manipulation came via coset enumeration. BHN was one of the early practitioners and he continued to enjoy it for a long time. He saw that it would benefit from the use of electronic computers. He argued for computational resources at ANU in the 1960s and established a post to support computation. He set interesting challenges for the development of better implementations (see B107c for an example and MR1775892, 2000). He continued to record such manipulations in his notebooks and to publish consequences he thought of interest.

In B47 BHN considered the question of whether a finite group with trivial Schur multiplier has a presentation with an equal number of generators and relations, in other words has deficiency zero. This is a growing subject to which BHN contributed further in B106, dedicated to Coxeter who was another great presentation manipulator, B117, B122 and B123.

In the course of proving that it is possible to adjoin n 'th roots of elements to semi-groups in B85 and B86, BHN suggested a

computational procedure, analogous to the Todd–Coxeter procedure for groups, to enumerate the elements of a finitely presented monoid. The procedure was proved by Jura (MR0486223, 1978), and various variants have been proposed, culminating in a general framework of Linton (MR1344920, 1995) that encompasses double coset enumeration in groups and vector enumeration, as well as the Todd–Coxeter procedure for groups and for monoids as special cases.

C Varieties

C.1 Varieties of Groups

BHN initiated the study of identical relations in groups in his Cambridge dissertation (1935), *Identical relations in groups*. The investigation was set in motion from his study of Boolean matrices B2 (1932). The introduction to the paper B7 (1937) that arose from the thesis suggests that the famous paper (JFM33.0149.01, 1902) of Burnside played a part in setting the direction of the work. At that time not much progress had been made on the questions Burnside had posed though they were being thought about (see footnotes on p. 506 of B7). What mattered for BHN was that the groups satisfied a relation $x^k = 1$ identically and, related to that, the idea of a 'maximal group' generated by n elements and having the order of all its elements divide a fixed number k . BHN asserted (B7, p. 507):

The identical relations that hold in a given group are well worth being closely studied; whereas the relations which connect the generators of a group depend on the choice of the generators, the identical relations clearly depend only on the group itself: they are an invariant property of the group.

The paper B7 is labelled I; it deals with the general theory. It introduces 'word-groups', now called verbal subgroups, and the V -groups $V_n(G)$, now called relatively free groups—that is, the n -generator group defined by the identical relations in n variables that hold in G . BHN proved that for a

finite group G the groups $V_n(G)$ are finite; more precisely, the order of $V_n(G)$ divides $|G|^{|G|^n}$. He also observed that $V_n(G)$ can be determined from G and n in a finite number of steps. He then said (B7, p. 520):

But to get *all* the identical relations we have to know something about $V_n(G)$ for *every* n . Whether this question can still be reduced to a finite problem cannot be decided now.

This comment evolved into the *finite basis problem*: does every group have a finite basis for its identical relations, that is, is there a finite set of identical relations from which all the identical relations follow?

The second part of the thesis, which contains detailed computations of the identical relations or laws of a number of finite metabelian groups, was never edited for publication (B120, p. 144). BHN also pointed out (B7, p. 509) that in Hall's paper (JFM62.0082.02, 1936) on Eulerian functions the V -groups for the icosahedral group are implicitly determined.

Nowadays this field of study is given the name 'varieties of groups'. The term variety in this context was introduced by P. Hall in 1949.

Graham Higman and BHN opened another aspect of the subject by showing in B28 (1952) that the variety of groups can be described by a single law in terms of the binary operation of right division.

Encouraged by Padmanabhan, BHN returned in B111 and B118 to finding single laws for the variety of groups this time in terms of the natural operations. His technique is based on the use of left and right mappings. This theme has been taken up for other varieties and in particular using automated theorem provers.

All BHN's other publications on varieties deal with problems that arose in Hanna's major paper (MR0078374, 1956) on varieties.

BHN showed in B49 (1956) that there is a 3-generator non-metabelian group all of whose 2-generator subgroups are metabelian and a 4-generator

non-metabelian group all of whose 3-generator subgroups are metabelian. These groups have exponent 8, and were the first of several examples demonstrating the complexity of groups with exponent 8. The most recent is by Krasilnikov (MR1994688, 2003).

The visit of BHN, Hanna and their son Peter to New York for the academic year 1961–2 provided a flourish of activity on varieties that is reported in two papers. In B67 (1962) the three Neumanns settled questions about the structure of the lattice and algebra of varieties left open in MR0078374. In particular the semigroup of varieties of groups (omitting the variety of all groups and that of trivial groups) is free and freely generated by indecomposable varieties, a result obtained independently by Shmel'kin (MR0151539, 1963). The important new idea used is to relate wreath products of groups to products of varieties of groups.

In B75 (1964) Baumslag and the three Neumanns considered conditions under which a variety is generated by a relatively free group of finite rank. They introduced, and made effective use of, the notion of a 'discriminating' set of groups. For example, they showed that the variety of all groups with given soluble length is generated by the relatively free group of rank 2.

In 1967 BHN gave an invited lecture to the American Mathematical Society in which he surveyed work on varieties referring his listeners to Hanna's monograph that was about to appear; B83 is the text.

The finite basis problem was the impetus for a lot of work which established 'varieties of groups' as a separate discipline in the 1950s (see the index of [2]) with its status reported in a monograph by Hanna (MR0215899, 1967). Oates and Powell, students of Higman, proved in 1964 that the identical relations of a finite group always have a finite basis (MR0161904, 1964). Then in 1970, in quick succession, Ol'shanskii, Adian and Vaughan-Lee

(MR0286872, MR0257189, MR0276307) published quite different proofs that there are groups that do not have a finite basis for their identical relations and indeed there are continuously many varieties of groups.

Many of the results about varieties of groups have been transferred to varieties of Lie algebras especially over infinite fields.

BHN and Macdonald took up a theme from BHN's PhD—to study the interdependence and independence of commutator laws in groups in a series of papers B121, B124, B127.

C.2 Varieties of Other Algebras

BHN and Evans (B30, 1953), answering a question in B25, showed that there are continuously many varieties of groupoids and loops. BHN and Christine Wiegold (B81, 1966) reported on semigroup representations of varieties of algebras. In B87 (1968) BHN applied the techniques of B28 to improve a result of Grätzer concerning a variety \mathfrak{R} of algebras in a certain species such that \mathfrak{R} is defined by a finite set of laws. Grätzer had constructed a new species by adding two binary operations, and a variety \mathfrak{R}_1 in that species, such that both \mathfrak{R}_1 restricted to \mathfrak{R} in the original species, and also \mathfrak{R}_1 could be defined by two laws. In B87 BHN showed that \mathfrak{R}_1 can be defined by a single law. BHN was an early contributor in B116 (1984) to the now extensive theory of knot quandles. BHN also studied other varieties of algebras B109, B125, B133, B134.

D A Little Geometry

The foundations of geometry were a significant source of inspiration early in BHN's career. His interest in geometry went back to his early days as an undergraduate at Freiburg in 1928. Most of his publications related to geometry deal with algebraic questions that arise—primarily in foundations B11, B17, B18, B44. There are two other themes—convex plane regions and

Napoleon's Theorem. The former was the basis for various lectures including the Matthew Flinders Lecture to the Australian Academy of Science. It has also led, via a question popularized by J. Moser, to developments in dynamical systems—the theory of outer billiards. The approach of BHN to Napoleon's Theorem is quite algebraic.

D.1 Convex Regions

BHN (B10, 1939) showed that a certain parameter associated with the lengths of chords of a plane closed convex curve must lie between $1/3$ and $1/2$. Then in B16 (1945, 1946) BHN found that the same bounds hold for an analogous parameter related to the area of intersection with a half-plane of a closed finite planar region. His proof applies without alteration if the planar region is replaced by an appropriate planar mass distribution. Moreover in the case of convex planar regions BHN gave a simplified proof, improved the lower bound to $4/9$, and conjectured that a lower bound of $(3 - \sqrt{5})/2$ may hold. Recent computational work by Kaiser (MR1424359, 1996) suggests that the conjecture is true and may be exact in some cases. The main result from B16 and its generalization by Rado (MR0021962, 1946) for higher dimensions is now viewed as a result on measures in Euclidean spaces, called the Neumann–Rado Theorem (see Dolnikov MR1187871, 1992), and has been generalized by many others, most notably by Tverberg (MR0187147, 1966).

D.2 Outer Billiards

Around 1960 BHN gave some popular lectures on Dupin curves under the title: 'Sharing ham and eggs'. There is a written account B52b of one such lecture in the magazine *Iota* published in Manchester. In this account BHN asked a question about what is now known as outer or dual billiards. This question was popularized by Moser as a crude model for planetary motion

(MR0442980, 1978). Recently Schwartz (MR2299242, 2007) answered what he now calls the Moser–Neumann question and published a monograph on outer billiards (MR2562898, 2009).

D.3 Douglas–Neumann Theorem

The (Petr–)Douglas–Neumann theorem is an ‘astounding’ generalization (see MR1959194, 2003) of what is usually referred to as Napoleon’s Theorem: If on the sides of a triangle are erected (outside or inside) equilateral triangles, then the centroids of these triangles form an equilateral triangle with the same centroid as the original triangle. The generalization can be stated (see Theorem 3.2 of B12): If isosceles triangles with base angles $\pi/2 - 2\pi\nu/2n$ are erected on the sides of an arbitrary polygon Π , and if this process is repeated with the polygon formed from the free vertices of the triangles, but with a different value of ν , and so on until all values of ν except 0 and one other (arbitrary) value, say μ , have been used in arbitrary order, then a regular figure is obtained. Its centroid coincides with that of the vertices of Π , and its sides are equal in length and (parallel but) opposite in direction to the sides of the polygon that would have been obtained from Π by taking $\nu = \mu$. Jesse Douglas was one the (two) first Fields Medallists in 1936 for his work on the plateau problem. BHN’s interest stemmed from a problem about electrical transformers which he came across in proof-reading his father’s book, *Symmetrical Component Analysis of Unsymmetrical Polyphase Systems* (1939). BHN’s paper attracted the interest of the well-known Cambridge geometer Baker (MR0008453, 1942) who gave a direct proof. BHN in turn polished it further B13. BHN gave a number of lectures on this topic. There is a nice, more recent, account in B113. A Java script that illustrates the theorem can be found on the web (<http://www.maa.org/editorial/knot/Napoleon.html> (*sic*)).

E Constructions and Embeddings

A major topic of interest to BHN was the embedding of groups or algebras in other groups or algebras, frequently with the preservation of given properties. Many of his embedding results are achieved by ingenious constructions, and may be viewed either as embedding or construction theorems. Most notable are the famous Higman–Neumann–Neumann (HNN) construction for groups and the Mal’cev–Neumann construction for division rings. Other important examples include the twisted wreath product, the non-Hopfian group constructions, and the ‘Neumann semidirect product’ for semigroups.

E.1 Subgroups of Direct Products

Several authors remark on a construction in B6 (1937). It is described in detail in a recent book by de la Harpe (MR1786869, 2000). BHN used it to construct continuously many pairwise non-isomorphic finitely generated groups. They are all subgroups of unrestricted direct products of alternating groups which contain the restricted direct product of these groups. Since there are only countably many finitely presented groups, most of BHN’s examples must be not finitely presented. Indeed, recently Baumslag and Miller showed that all of them are not finitely presented. Baumslag (and others) have used these groups to provide examples of non-trivial discriminating groups (MR2288461, 2007). They are significant in the theory of amenable groups because they show there are continuously many finitely generated elementary amenable groups.

BHN made other use of his insight into unrestricted direct products. The embedding of a relatively free group $V_n(G)$ into the direct product of $|G|^n$ copies of G introduced by Birkhoff (1935) and by BHN (B7, 1937) has been called the Birkhoff–Neumann embedding (see MR2245589, 2006). The restricted product is usually

not complemented in the unrestricted one (B76, 1965). BHN and Yamamuro used it (B80, 1965) to show that for one family of subgroups, the Boolean powers, all countable quotients are finite. In an addendum they remarked that this fits into a broader context.

E.2 The Higman–Neumann–Neumann (HNN) Construction

Given a group G with subgroups A and B , when does there exist a group H containing G such that A and B are conjugate in H ? Necessarily A and B must be isomorphic. The HNN construction (B19, 1949) demonstrates that this is sufficient, constructing groups that are now known as HNN extensions. They are closely related to amalgamated free products and have many analogous properties. HNN extensions are used in B19 to embed any countable group in a 2-generator group, and to prove that any group G can be embedded in a group in which all elements of G of the same order are conjugate. They are at the heart of G. Higman's famous embedding theorem (MR0130286, 1961) showing that a finitely generated group can be embedded in a finitely presented group if and only if it is recursively presented. Most important in their application these days is Britton's Lemma (MR0168633, 1963) giving a normal form for an HNN extension. The HNN construction is of considerable importance in combinatorial and geometric group theory. It is used in essentially all work on unsolvable decision problems. The use of HNN embedding plays a significant role in the theory of amenable and non-amenable groups, and there are many other recent applications (for example MR2294245 and MR2270456, 2007).

E.3 Wreath Products and Generalizations

The wreath product construction has a long history and remains a fundamental tool in both abstract group theory and applications

of group actions, in algebra, combinatorics and statistics. BHN and Hanna pioneered the use of wreath products in their work on embeddings in B54 (1959) and B59 (1960). In B54, BHN and Hanna showed that every countable soluble group G is embeddable into a soluble 2-generated group H . This embedding theorem was extended to generalized soluble groups in B77 (1965) (with Kovács). In B59, BHN showed that if the (soluble) group G is fully ordered, then the 2-generated (soluble) group H can be made fully ordered with G order isomorphic to its image in H . This result was extended to generalized soluble groups with a full order by Mikaelian (MR1898374, 2002; MR1970055, 2003).

In B68 (1963) BHN introduced a generalization of this construction that he called the *twisted wreath product* of which the wreath product is a special case. The general notion leads in B65 to a sharpened form of a theorem of Auslander and Lyndon on the lower central series of normal subgroups of free groups. The twisted wreath product gives its name to one of the small number of types of finite primitive permutation groups in the O'Nan–Scott scheme. A number of other generalizations of the wreath product have appeared in the literature. Of these the *crown product* introduced in B48 (1956) is used to study ascending derived series of groups.

BHN was the first to introduce a semidirect product construction for semigroups. In B57 he used it to define a wreath product of semigroups, to prove that every finite semigroup can be embedded in a finite 2-generated semigroup, and also to give a new proof of Evans' theorem (MR0050566, 1952) that every countable semigroup can be embedded in a 2-generator semigroup. The 'Neumann semidirect product' turned out to be a very powerful tool for semigroup theory. For example, Preston (MR0837162, 1986) investigated a more general product definition and showed that only the direct product and the Neumann semidirect

product have the crucial property that the product of two semigroups is always a semigroup.

E.4 Group Amalgams

Group amalgams arose in Hanna's work on free products with amalgamated subgroups (MR0026997, 1948) and the amalgam concept was introduced shortly afterwards by Baer (MR0030953, 1949). Amalgams play an important role in the study of finite simple groups and associated geometries (for example see MR1705272, 1999). A group amalgam is an 'incomplete group' in that its set of elements is the union of groups G_i , for $i \in I$, such that for distinct i, j the intersection $H_{ij} = G_i \cap G_j$ is a subgroup of both G_i and G_j . In particular, the product of two elements is only assumed to exist if some G_i contains both elements. An amalgam is embeddable if there is a one-to-one homomorphism from it to some group. BHN in collaboration with Hanna undertook an intense study of group amalgams. In B21 (1950) they studied a family of amalgams in which the subgroups H_{ij} are central in both G_i and G_j , and constructed what are now called *central products* of the groups G_i to prove that such amalgams are embeddable. In B32 (1953) they introduced uniform notation to state all the then-known embedding results and showed that each was best possible. In particular they asked whether an embeddable amalgam consisting of finite groups is embeddable in a finite group. This question was answered in the negative by Brown (MR1230631, 1992). In B58 (1960) BHN introduced the permutational product of an amalgam of two [sub]groups as an embedding group for the amalgam. He studied the extent to which various finiteness properties of the amalgam carry over to the permutational product. Using this construction he showed (B56, 1960) by example that the properties of local finiteness, or being periodic need not carry over to the embedding group. BHN (B60, 1960) explored existence questions

for linked products of given groups (introduced by Hanna and Wiegold, MR0124386, 1960), and (B66, 1962) demonstrated that there are no necessary and sufficient criteria involving only commutators, intersections, and multiplications of subgroups for the existence of a generalized free nilpotent product, or a generalized free soluble product, of the amalgam of two groups.

E.5 Mal'cev–Neumann Theorem

It was the role of ordered groups and ordered division rings in the foundations of geometry that drew BHN's interest (B120, p. 800). In B17 (1949) BHN generalized some necessary conditions, and some sufficient conditions proposed by Levi for a group to be orderable, producing some general constructions for ordered groups, and the first example of a perfect ordered group. Ordered division rings are even more important in geometry than ordered groups, and in B18 (1949) BHN constructed a formal power series division ring and used this to prove that every ordered group can be embedded in an ordered division ring. The construction made independently by Mal'cev (MR25457, 1948) is now known as the Mal'cev–Neumann construction and is of central importance in the theory of division rings (see Chapter 2 of Cohn's book MR1349108, 1995). It has had a far-reaching impact in areas as diverse as classical analysis, diophantine equations, combinatorics and commutative algebra.

E.6 Non-Hopfian Constructions

A *Hopf group* is one that is not isomorphic to any of its proper quotient groups. Hopf had conjectured in the 1930s that all finitely generated groups should be what we now call Hopf groups. The first suggestion that this was not so came in a paper by Baer that 'shook the group-theoretical world' (B120, p. 905). Combining Baer's basic ideas with use of the newly developed HNN-extension of B19, BHN constructed (B22,

1950) the first finitely generated non-Hopf group (a 2-generator group). In B31 (1953) BHN and Hanna answered another question of Hopf negatively by constructing two non-isomorphic (finitely presented) groups, each a homomorphic image of the other. Later (B54, 1959) BHN and Hanna constructed the first example of a soluble non-Hopf group.

E.7 Essay for the Adams Prize

The largely expository prize-winning essay (B38, 1954) has been widely read and has had substantial influence. Its origins lie in an appendix that BHN was invited to write as an up-date for the 1953 German translation of Kurosh's book *Teoriya Grupp*, to 'describe some of the then new results that had been obtained and that answered many of the problems in Kurosh's book' (B120, p. 281). The appendix was updated, translated into English, and submitted to the University of Cambridge for the Adams Prize, which it won. The published essay B38 contains basic facts about amalgamated free products with all sorts of applications and examples, including the HNN construction, 2-generator embedding, and 'three remarkable groups of G. Higman', in a very readable exposition.

E.8 More on Embeddings of Groups and Algebras

In B14 (1943), written while on leave from the British army, BHN studied the problem of solving equations in groups and adjoining the necessary solutions. He developed an elegant procedure, proving that every group can be embedded in a divisible group (one in which, for every group element g and every positive integer n , the equation $x^n = g$ has a solution x in the group).

In B17 (1949) BHN noted that every ordered group can be embedded in the multiplicative group of an ordered division ring, and in the early 1950s, BHN asked whether every totally ordered group can

be embedded in a totally ordered divisible group. Despite forming the topic for PhD theses in 1974 and 2002, and being listed as the first and most notorious problem in the Black Swamp Problem Book (a problem book on ordered groups carried around to conferences in this area of mathematics and containing contributions from participants), this problem was not solved until Bludov (MR2213301, 2005) produced a totally ordered group for which no such embedding exists.

In B25 (1951), arising from discussions with Evans, BHN showed that a non-associative ring R is embeddable in a division ring D with unique left and right division if and only if R has no zero divisors. Moreover, D may be chosen to retain certain properties of R such as commutativity or being idempotent.

In a subsequent paper (B37, 1954) BHN derived from a theorem of Steenrod a general embedding result for a large class of universal algebras that contains all (partial) semigroups. He deduced a number of striking consequences: a group can be fully ordered if and only if all its finitely generated subgroups can be fully ordered; a ring can be embedded in a division ring if and only if every finitely generated subring can be so embedded; the edges of a graph can be coloured with k colours so that no incident edges have the same colour provided the same is true for every finite subgraph. Many of these consequences would these days be derived from the Compactness Theorem of first order logic. However several now-senior mathematicians recall the high impact B37 had on them as students, one finding the paper 'very beautiful and clever', another finding it had 'changed forever [his] view of the mathematical world to see that one could take such a global view of the world and still prove significant theorems'.

In B70 (1963) with Tekla Taylor, BHN characterized those semigroups that can be embedded in nilpotent groups, a

characterization similar to an earlier one by Mal'cev (MR75959, 1953).

Other papers of BHN may be regarded as a prelude to B19 (B20 written in June 1948—see the endnote to B20 on SW118), or a development of B19 (B27, B40). In this context, Problem 11.67 in the Kourovka notebook by BHN asks: does there exist a torsion-free group with exactly 3 classes of conjugate elements such that no non-trivial conjugate class contains a pair of inverse elements?

F Other Contributions

F.1 A Group-theoretic Arithmetic Problem

B3 (1933) shows the existence of continuously many maximal non-parabolic subgroups of the modular group. The existence of such subgroups was needed by Arnold Schmidt for his doctoral thesis on the axiomatics of geometry (B120, p. 65). There is an extensive account in a monograph by Wilhelm Magnus on *Noneuclidean Tessellations (sic) and Their Groups* (MR0352287, 1974).

F.2 A Problem like Burnside's

BHN considered, though not as part of his thesis work, a problem similar to Burnside's that he called the *bounded orders problem*. In B8 he considered the simplest non-trivial case: groups all of whose elements have order at most 3.

The last thirty years have seen an extension of the bounded orders problem. The set of orders of elements of a periodic group has been called the *order spectrum* of the group. Given a divisibility-closed subset Ω of the positive integers the extended problem asks for a description of the periodic groups whose order spectrum is Ω , preferably in the strong form of determining the groups up to isomorphism. There is a recent survey on this topic by Mazurov and Shi (MR2491731, 2009; see also also MR1676650, 1999).

There are three types of order spectrum for groups all of whose elements have order at most 3. The solution of the bounded orders problem for groups with order spectrum $\{1, 2\}$ was known when Burnside posed his questions. BHN solved the bounded orders problem for finitely generated groups with order spectrum $\{1, 2, 3\}$. The strong form of the problem has still not been solved for groups with order spectrum $\{1, 3\}$, that is groups with exponent 3.

F.3 The Grushko–Neumann Theorem

The Grushko–Neumann Theorem asserts that the number of generators needed to generate a free product of groups is the sum of the number of generators for each of the free factors. The BHN proof (B15, 1943) of this result was written while he was on leave from the Pioneer Corps and published in 1943, responding to a question of Levi. During the war both communication and publication were difficult, and we note that the proofs by Grushko and BHN were obtained independently even though the former (MR0003412) dates from 1940. In [4, p. 106] the theorem is described as the next development in the theory of free products after the work of Kurosh. This important result has attracted intense interest since then with alternative proofs offered by Wagner (the infinitely generated case, MR0008809, 1957), Lyndon (MR0178073, 1965), Stallings (MR0188284, 1965) and Imrich (MR0773183, 1984).

F.4 The Frattini Subgroup

In B6 (1937) BHN proved fundamental properties of what is now called the Frattini subgroup for not necessarily finite groups.

In B35 (1954) BHN and Higman solved two questions of Itô about the Frattini subgroup. They showed that the Frattini subgroup of a group need not be nilpotent and that the Frattini subgroup of a non-trivial free product is trivial. They pointed out that, for free products with an amalgamated

subgroup, the Frattini subgroup can coincide with the amalgamated subgroup, and they asked if it can be larger. It has been shown independently and by different methods that the Frattini subgroup is contained in the amalgamated subgroup, by Gelandner and Glasner (MR2377495, 2008) if the free product is countable, and by Allenby and Tang (MR2419010, 2008) if the amalgamated subgroup is countable.

In B52 (1958) BHN studied ascending Frattini series. He concluded by asking whether a finite group that is the Frattini subgroup of some group is also the Frattini subgroup of a finite group. This question was answered in the affirmative by Eick (MR1427706, 1997).

F5 Almost Finite and Almost Abelian Groups

BHN's work over many years on groups that are in some sense close to being finite, or close to abelian, has been very influential. In B24 and B53 he studied groups in which the conjugacy classes of elements are all finite, in particular proving an extension of the Sylow theorems for these infinite groups. In B36 and B41, BHN studied BFC-groups, groups for which there is a finite upper bound on the sizes of the conjugacy classes of elements, and proved that these are precisely the groups with finite derived subgroup (a measure of commutativity). The question in B36, p. 239, asking for an explicit upper bound for the order of the derived group in terms of the upper bound on the conjugacy class sizes of such a group has been investigated over many years. BHN's review of a paper by Segal and Shalev (MR1726791, 1999) gives an account of recent results.

In similar vein BHN proved in B42 equality of the class of groups whose centre has finite index (another measure of commutativity) with the class of groups all of whose conjugacy classes of subgroups are finite. Baer's observation that this class is

also equal to the class of groups which can be covered by finitely many abelian subgroups prompted BHN's investigation (B39, B41) of groups coverable by finitely many cosets of proper subgroups. If such a covering is minimal in the sense that no proper subset of the cosets covers the group, then BHN's results in B36 imply that all the cosets correspond to subgroups of finite index, and bounds for their indices are given in B41. BHN's work on coverings, in the contexts of both abstract groups and group actions, has been applied and developed in many ways by many mathematicians up to the present (see MR0401882, 1976; MR1328918, 1994).

A more combinatorial measure of commutativity is provided by the non-commuting graph $\Gamma(G)$ of a group G in which elements of G are joined by an edge if and only if they do not commute. In 1975, Erdős asked of groups G where $\Gamma(G)$ contains no infinite complete subgraph or, equivalently, groups in which every infinite subset of elements contains two that commute: does there exist an integer $n(G)$ such that every complete subgraph of $\Gamma(G)$ has at most $n(G)$ vertices? In B104 BHN characterized such groups G as those for which the centre Z has finite index, and proved that $n(G) \leq |G|/|Z| - 1$ (see also MR0506252, 1978). This result inspired many further investigations and is much cited (see the review MR0419283, 1996; the sequel B135; and MR2260490, 2006).

F6 Algebraic Closure: the Macintyre–Neumann Theorem

BHN's work on algebraic closure, also called existential closure, in groups and general algebras began with a question of Scott (MR0040299, 1951). Scott had introduced notions of algebraic closure and weak algebraic closure for groups. Using the generalized free product construction, BHN proved (B26, 1952) that these notions are equivalent, and moreover

that each algebraically closed group is simple. Recognising that these notions belong to universal algebra rather than group theory, BHN proved analogous results for algebraically closed semigroups (B93, 1971) and related results for cancellative semigroups (B92, 1970; see SW1116).

BHN's second major paper (B97, 1969/1973) on this theme was widely circulated as a preprint several years before it was published. In it he proved (among other things) that a finitely generated group with soluble word problem can be embedded in every algebraically closed group—one half of the Macintyre–Neumann Theorem. Macintyre (1972) proved the converse for finitely generated groups (and more—see Schupp's review MR0414671 of B97). At that time logicians from Abraham Robinson's school of model theory were developing generic structures using forcing and were interested in existentially closed structures because of work of Macintyre. This result led to work on existentially closed groups by Simmons, Macintyre, Ziegler and others.

F.7 Properties of Countable Character

In B99, dedicated to the memory of Mal'cev and written a little before B96 (see B120, p. 1259), BHN studied group properties of countable character. These are properties possessed by a group if and only if all the countable subgroups possess them. BHN showed in B99 that residual finiteness is a property of countable character. 'Properties of countable character' is the title of BHN's invited lecture at the International Congress of Mathematicians in Nice in 1970. In the account B96 of that lecture BHN considered this theme for general algebras.

F.8 Means in Groups

In B33, BHN simplified necessary and sufficient conditions, due to Scott, for the existence on a group of a set of 'Schimmack means'. In doing so he constructed the

first example of a nonabelian group admitting such means. Subsequently Wiegold (MR120277, 1960) showed that both Scott's and BHN's conditions are equivalent to requiring that the centre of the group and its factor group are both direct products of groups isomorphic to the additive group of rational numbers.

F.9 Unsolvability and the Axiom of Choice

BHN learnt from Remak in Berlin to be alert to the use of the Axiom of Choice when working with infinite sets and this was reinforced by lectures there by von Neumann. There are signs of this influence in quite a lot of his papers, for example in B6. It surfaces explicitly in B102 where Hickman and BHN showed that the answer to a question of Babai depends on the underlying set theory. Another publication at the confluence of algebra and logic arose out of a visit by Boone to Manchester in 1958. Their discussions together with Baumslag led to results on the algorithmic unsolvability of questions about groups B55.

Acknowledgements

Many people have helped in the preparation of this memoir. Foremost I thank M. F. (Mike) Newman to whom I am especially indebted and without whose help and scholarship over several years this memoir could not have been written. In particular, together with L. G. (Laci) Kovács, he prepared the BHN bibliography below, and I thank both Mike and Laci for their patience and dedication in presenting this accurate listing of BHN's published work. I thank members of Bernhard's family for their patience and assistance checking innumerable details and reading drafts: Dorothea Neumann, Peter Neumann, Walter Neumann, Barbara Cullingworth, Daniel Neumann.

I am greatly indebted to James Wiegold, Martin Taylor, and many others who have helped in various ways: Reg Allenby, Gilbert Baumslag, Keith Burns, Slava Grigorchuk, Hermann Heineken, Charles

Holland, Andrei Kelarev, Ann Chi Kim, C. F. (Chuck) Miller III, Sidney Morris, Akbar Rhemtulla, Douglas Rogers, K. P. Shum, Anne Penfold Street, Zhi-Wei Sun, F. C. Y. Tang, Helge Tverberg, Mikhail Volkov, Sheila Williams, J. S. Wilson.

In addition to generous personal contributions, I am indebted to sources of biographical information such as the record of an interview [1] of BHN by Professor Bob Crompton in 1998, and the obituaries (MR0342340, 1974) and (MR1983807, 2003) of Hanna Neumann and BHN.

The formal portrait of B. H. Neumann was taken for the Royal Society in 1984.

Students

BHN was the supervisor of the work of seventeen students for doctorates. He was the formal supervisor of several other students. He was interested in the work of a wider group of students. He also gave help to some students of which the most significant was that given to Kamal Riad Yacoub (see B120, pp. 65–66).

0. Yacoub 1953

Manchester

1. Chehata Chehata	1953
2. John Britton	1954
3. Christine Dale	1957
4. Gilbert Baumslag	1958
5. James Wiegold	1959
6. Ian Macdonald	1960
7. Michael Newman	1960
8. László Kovács	1961

ANU

9. Narain Gupta	1965
10. Terence Gagen	1967
11. Ahmad Shafaat	1968
12. Neil Williams	1969
13. Andrew Brunner	1973
14. Kok Yéo	1973
15. Lambertus Hesterman	1974
16. Colin Fox	1974
17. Tim Brook	1974

References

1. Professor Bernhard Neumann (1909-2002), Mathematician, Interviews with Australian Scientists, Interviewed by Professor Bob Crompton in 1998. <http://www.science.org.au/scientists/bn.htm>
2. *Mathematical Reviews*, The American Mathematical Society, Providence, RI. For *Mathematical Reviews on the Web* see <http://www.ams.org/mathscinet/>
3. *Jahrbuch über die Fortschritte der Mathematik (JFM)* (1868–1942), available electronically through the Electronic Research Archive for Mathematics at <http://www.emis.de/projects/JFM/>
4. Bruce Chandler and Wilhelm Magnus, *The History of Combinatorial Group Theory: A Case Study in the History of Ideas*. Studies in the History of Mathematics and Physical Sciences, 9. Springer-Verlag, New York, 1982.

Publications of B. H. Neumann

Several entries in this list have a lower case roman letter as an additional label following the numeral, and represent papers published or submitted at roughly the same time as the ‘unadorned entry’, for example [B 4a] and [B 4] were both published in 1935. The entries [B 1a] and [B 5a] are the two doctoral theses. All other such entries up to [B 115a] are numbered as in the *Selected Works* (B120, 1988). BHN explained there that he added letters for items that he considered of lesser importance, and this was the guiding principle for later entries.

- | | |
|-------|---|
| B 1. | Bernhard Neumann, Die Automorphismengruppe der freien Gruppen, <i>Math. Ann.</i> 107 (1932), 367–386. |
| B 1a. | Bernhard Neumann, Die Automorphismengruppe der freien Gruppen, Inaugural-Dissertation, Friedrich-Wilhelms-Universität zu Berlin, 1932. |
| B 2. | Bernhard Neumann, Matrizenkalkül in der Booleschen Algebra, <i>Jber. Deutsche Math.-Verein.</i> 42 (1932), 126–130 (1933). |
| B 3. | Bernhard Neumann, Über ein gruppentheoretisch-arithmetisches Problem, <i>Sitzungsber. Preuss. Akad. Wiss. phys.-math. Kl.</i> X (1933), 429–444. |
| B 4. | B. H. Neumann, Decomposition of groups, <i>J. London Math. Soc.</i> 10 (1935), 3–6. |

- B 4a. Bernhard H. Neumann, Aufgabe 188, *Jber. Deutsche Math.-Verein.* **45** (1935), 22.
- B 5. B. H. Neumann, A remark on primary ideals, *J. London Math. Soc.* **10** (1935), 111–112.
- B 5a. Bernhard Hermann Neumann, Identical Relations in Groups, Ph.D. thesis, University of Cambridge, 1936.
- B 6. B. H. Neumann, Some remarks on infinite groups, *J. London Math. Soc.* **12** (1937), 120–127.
- B 7. B. H. Neumann, Identical relations in groups. I., *Math. Ann.* **114** (1937), 506–525.
- B 8. B. H. Neumann, Groups whose elements have bounded orders, *J. London Math. Soc.* **12** (1937), 195–198.
- B 9. B. H. Neumann, Identical relations in groups, *Comptes Rendus Congrès Internat. des Math.* (Oslo 1936), Tome 2, A.W. Brøgger's Boktrykkeri, Oslo, 1937, pp. 18–19.
- B 10. B. H. Neumann, On some affine invariants of closed convex regions, *J. London Math. Soc.* **14** (1939), 262–272.
- B 11. B. H. Neumann, On the commutativity of addition, *J. London Math. Soc.* **15** (1940), 203–208.
- B 12. B. H. Neumann, Some remarks on polygons, *J. London Math. Soc.* **16** (1941), 230–245.
- B 13. B. H. Neumann, A remark on polygons, *J. London Math. Soc.* **17** (1942), 165–166.
- B 14. B. H. Neumann, Adjunction of elements to groups, *J. London Math. Soc.* **18** (1943), 4–11.
- B 15. B. H. Neumann, On the number of generators of a free product, *J. London Math. Soc.* **18** (1943), 12–20.
- B 16. B. H. Neumann, On an invariant of plane regions and mass distributions, *J. London Math. Soc.* **20** (1945), 226–237 (1946).
- B 17. B. H. Neumann, On ordered groups, *Amer. J. Math.* **71** (1949), 1–18.
- B 18. B. H. Neumann, On ordered division rings, *Trans. Amer. Math. Soc.* **66** (1949), 202–252.
- B 19. G. Higman, B. H. Neumann, and Hanna Neumann, Embedding theorems for groups, *J. London Math. Soc.* **24** (1949), 247–254.
- B 20. B. H. Neumann, On a special class of infinite groups, *Nieuw Arch. Wiskunde* (2) **23** (1950), 117–127.
- B 21. B. H. Neumann and Hanna Neumann, A remark on generalized free products, *J. London Math. Soc.* **25** (1950), 202–204.
- B 22. B. H. Neumann, A two-generator group isomorphic to a proper factor group, *J. London Math. Soc.* **25** (1950), 247–248.
- B 23. B. H. Neumann and H. Neumann, Zwei Klassen charakteristischer Untergruppen und ihre Faktorgruppen, *Math. Nachr.* **4** (1951), 106–125.
- B 24. B. H. Neumann, Groups with finite classes of conjugate elements, *Proc. London Math. Soc.* (3) **1** (1951), 178–187.
- B 25. B. H. Neumann, Embedding non-associative rings in division rings, *Proc. London Math. Soc.* (3) **1** (1951), 241–256.
- B 26. B. H. Neumann, A note on algebraically closed groups, *J. London Math. Soc.* **27** (1952), 247–249.
- B 26a. B. H. Neumann, On some interesting sets of circles, *Math. Gaz.* **36** (1952), 121–122.
- B 27. B. H. Neumann and Hanna Neumann, Extending partial endomorphisms of groups, *Proc. London Math. Soc.* (3) **2** (1952), 337–348.
- B 28. G. Higman and B. H. Neumann, Groups as groupoids with one law, *Publ. Math. Debrecen* **2** (1952), 215–221.
- B 29. B. H. Neumann and H. Neumann, On a class of abelian groups, *Arch. Math.* **4** (1953), 79–85.
- B 30. T. Evans and B. H. Neumann, On varieties of groupoids and loops, *J. London Math. Soc.* **28** (1953), 342–350.
- B 31. B. H. Neumann, On a problem of Hopf, *J. London Math. Soc.* **28** (1953), 351–353.
- B 32. B. H. Neumann and Hanna Neumann, A contribution to the embedding theory of group amalgams, *Proc. London Math. Soc.* (3) **3** (1953), 243–256.
- B 33. B. H. Neumann, A note on means in groups, *J. London Math. Soc.* **28** (1953), 472–476.
- B 34. B. H. Neumann, Anhang, in *Gruppentheorie* by A. G. Kurosch (Mathematische Lehrbücher und Monographien. I. Abt. Band III.), Akademie-Verlag, Berlin, 1953, pp. 333–381.
- B 34a. B. H. Neumann, Függelék, Csoportok általánosított szabad szorzatai, in *Csoportelmélet* by A. G. Kurosch (Hungarian,

- translated by Sándor Gacsályi, Akadémiai Kiadó, Budapest, 1955, pp. 433–502.
- B 35. G. Higman and B. H. Neumann, On two questions of Itô, *J. London Math. Soc.* **29** (1954), 84–88.
- B 36. B. H. Neumann, Groups covered by permutable subsets, *J. London Math. Soc.* **29** (1954), 236–248.
- B 37. B. H. Neumann, An embedding theorem for algebraic systems, *Proc. London Math. Soc.* (3) **4** (1954), 138–153.
- B 38. B. H. Neumann, An essay on free products of groups with amalgamations, *Philos. Trans. Roy. Soc. London Ser. A* **246** (1954), 503–554.
- B 39. B. H. Neumann, Groups covered by permutable subsets, *Proc. Internat. Congr. Math.* (Amsterdam 1954), vol. 2, North-Holland, Amsterdam, 1954, p. 45.
- B 40. B. H. Neumann and H. Neumann, Partial endomorphisms of finite groups, *J. London Math. Soc.* **29** (1954), 434–440.
- B 41. B. H. Neumann, Groups covered by finitely many cosets, *Publ. Math. Debrecen* **3** (1954), 227–242 (1955).
- B 42. B. H. Neumann, Groups with finite classes of conjugate subgroups, *Math. Z.* **63** (1955), 76–96.
- B 43. W. Ledermann and B. H. Neumann, On the order of the automorphism group of a finite group. I, *Proc. Roy. Soc. London Ser. A* **233** (1956), 494–506.
- B 44. B. H. Neumann, Groups with automorphisms that leave only the neutral element fixed, *Arch. Math.* **7** (1956), 1–5.
- B 45. W. Ledermann and B. H. Neumann, On the order of the automorphism group of a finite group. II, *Proc. Roy. Soc. London Ser. A* **235** (1956), 235–246.
- B 46. B. H. Neumann, On a question of Gaschütz, *Arch. Math.* **7** (1956), 87–90.
- B 47. B. H. Neumann, On some finite groups with trivial multiplier, *Publ. Math. Debrecen* **4** (1956), 190–194.
- B 48. B. H. Neumann, Ascending derived series, *Compositio Math.* **13** (1956), 47–64.
- B 49. B. H. Neumann, On a conjecture of Hanna Neumann, *Proc. Glasgow Math. Assoc.* **3** (1956), 13–17.
- B 50. B. H. Neumann and J. A. H. Shepperd, Finite extensions of fully ordered groups, *Proc. Roy. Soc. London Ser. A* **239** (1957), 320–327.
- B 51. B. H. Neumann, Corrigendum and addendum to “Ascending derived series”, *Compositio Math.* **13** (1956), 128.
- B 52. B. H. Neumann, Ascending verbal and Frattini series, *Math. Z.* **69** (1958), 164–172.
- B 52a. B. H. Neumann, Review: *The Theory of Groups* by A. G. Kurosh; Review: *Gruppen-theorie* by Wilhelm Specht, *Math. Gaz.* **42** (1958), 151–154.
- B 52b. B. H. Neumann, Sharing ham and eggs, *Iota* (1958), no. 1, 14–18.
- B 53. B. H. Neumann, Isomorphism of Sylow subgroups of infinite groups, *Math. Scand.* **6** (1958), 299–307 (1959).
- B 54. B. H. Neumann and Hanna Neumann, Embedding theorems for groups, *J. London Math. Soc.* **34** (1959), 465–479.
- B 54a. B. H. Neumann, Review: *Structure of Rings* by Nathan Jacobson, *Math. Gaz.* **43** (1959), 73.
- B 55. Gilbert Baumslag, W. W. Boone, and B. H. Neumann, Some unsolvable problems about elements and subgroups of groups, *Math. Scand.* **7** (1959), 191–201.
- B 56. B. H. Neumann, On amalgams of periodic groups, *Proc. Roy. Soc. London Ser. A* **255** (1960), 477–489.
- B 56a. B. H. Neumann, Review: *Abelian Groups* by L. Fuchs, *Math. Gaz.* **44** (1960), 150–151.
- B 57. B. H. Neumann, Embedding theorems for semigroups, *J. London Math. Soc.* **35** (1960), 184–192.
- B 58. B. H. Neumann, Permutational products of groups, *J. Austral. Math. Soc.* **1** (1959/1960), 299–310.
- B 59. B. H. Neumann, Embedding theorems for ordered groups, *J. London Math. Soc.* **35** (1960), 503–512.
- B 60. B. H. Neumann and Hanna Neumann, On linked products of groups, *Acta Sci. Math. Szeged* **21** (1960), 197–205.
- B 61. B. H. Neumann, Lectures on topics in the theory of infinite groups, Notes by M. Pavman Murthy. Tata Institute of Fundamental Research Lectures on Mathematics, No. 21, Tata Institute of Fundamental Research, Bombay, 1960, reissued 1968, iii+267+iv pp.
- B 62. L. G. Kovács, B. H. Neumann, and H. de Vries, Some Sylow subgroups, *Proc. Roy. Soc. Ser. A* **260** (1961), 304–316.
- B 62a. B. H. Neumann, Review: *Boolean Algebras* by Roman Sikorski, *Math. Gaz.* **45** (1961), 365–366.

- B 63. B. H. Neumann, *Special topics in Algebra. Universal algebra*, Lectures delivered in the Fall Semester 1961–62, Notes by P. M. Neumann, Courant Institute of Mathematical Sciences, New York, 1962, Cyclostyled notes, ii+78 pp.
- B 64. B. H. Neumann, *Special topics in Algebra. Order techniques in algebra*, Lectures delivered in the Spring Semester 1962, Notes by P. M. Neumann, Courant Institute of Mathematical Sciences, New York, 1962, Cyclostyled notes, ii+77 pp.
- B 65. B. H. Neumann, On a theorem of Auslander and Lyndon, *Arch. Math.* **13** (1962), 4–9.
- B 66. B. H. Neumann and J. Wiegold, On certain embeddability criteria for group amalgams, *Publ. Math. Debrecen* **9** (1962), 57–64.
- B 67. B. H. Neumann, H. Neumann, and Peter M. Neumann, Wreath products and varieties of groups, *Math. Z.* **80** (1962), 44–62.
- B 67a. B. H. Neumann, Ferment in school mathematics, (Opening address at the) *Summer School for Mathematics Teachers*, University of New South Wales, 1963, pp. 0.1–0.10.
- B 68. B. H. Neumann, Twisted wreath products of groups, *Arch. Math.* **14** (1963), 1–6.
- B 68a. B. H. Neumann, Products of groups (Lectures delivered at the Summer Research Institute, Canberra, 1963; notes by Tekla Taylor), The Australian National University, 1963, i+55 pp.
- B 69. B. H. Neumann, A further note on means in groups, *J. London Math. Soc.* **38** (1963), 226–227.
- B 70. B. H. Neumann and Tekla Taylor, Subsemigroups of nilpotent groups, *Proc. Roy. Soc. Ser. A* **274** (1963), 1–4.
- B 71. B. H. Neumann, Felix Adalbert Behrend, *J. London Math. Soc.* **38** (1963), 308–310.
- B 72. B. H. Neumann and R. Rado, Monotone functions mapping the set of rational numbers on itself, *J. Austral. Math. Soc.* **3** (1963), 282–287.
- B 73. T. M. Cherry and B. H. Neumann, Felix Adalbert Behrend, *J. Austral. Math. Soc.* **4** (1964), 264–270.
- B 74. B. H. Neumann, Subsemigroups of nilpotent groups: an acknowledgement, *Proc. Roy. Soc. Ser. A* **281** (1964), 436.
- B 74a. B. H. Neumann, On decimal coinage, *Matrix* (1964), 16–18.
- B 75. G. Baumslag, B. H. Neumann, Hanna Neumann, and Peter M. Neumann, On varieties generated by a finitely generated group, *Math. Z.* **86** (1964), 93–122.
- B 76. B. H. Neumann, Supplements of direct powers in cartesian powers, *Math. Z.* **87** (1965), 17–18.
- B 77. L. G. Kovács and B. H. Neumann, An embedding theorem for some countable groups, *Acta Sci. Math. (Szeged)* **26** (1965), 139–142.
- B 78. L. G. Kovács and B. H. Neumann, On the existence of Baur-soluble groups of arbitrary height, *Acta Sci. Math. (Szeged)* **26** (1965), 143–144.
- B 79. G. Baumslag, L. G. Kovács, and B. H. Neumann, On products of normal subgroups, *Acta Sci. Math. (Szeged)* **26** (1965), 145–147.
- B 80. B. H. Neumann and S. Yamamuro, Boolean powers of simple groups, *J. Austral. Math. Soc.* **5** (1965), 315–324.
- B 81. B. H. Neumann and E. C. Wiegold, A semigroup representation of varieties of algebras, *Colloq. Math.* **14** (1966), 111–114.
- B 82. B. H. Neumann, On characteristic subgroups of free groups, *Math. Z.* **94** (1966), 143–151.
- B 82a. L. G. Kovács and B. H. Neumann, editors, *Proceedings of the International Conference on the Theory of Groups* (held at the Australian National University, Canberra, 10–20 August, 1965), Gordon and Breach Science Publishers, New York, 1967, xvii+397 pp.
- B 82b. B. H. Neumann, Making new finite simple groups, *Some problems and results in the theory of groups, II* (Notes of a mini-conference, Oxford, 1966), pp. 3–4.
- B 83. B. H. Neumann, Varieties of groups, *Bull. Amer. Math. Soc.* **73** (1967), 603–613.
- B 84. I. D. Macdonald and B. H. Neumann, A third-Engel 5-group, *J. Austral. Math. Soc.* **7** (1967), 555–569.
- B 85. B. H. Neumann, Some remarks on semigroup presentations, *Canad. J. Math.* **19** (1967), 1018–1026.
- B 86. B. H. Neumann, Some remarks on semigroup presentations: Corrigendum and addendum, *Canad. J. Math.* **20** (1968), 511.
- B 87. B. H. Neumann, On a problem of G. Grätzer, *Publ. Math. Debrecen* **14** (1967), 325–329.

- B 88. B. H. Neumann, Embedding theorems for groups, *Nieuw Arch. Wisk.* (3) **16** (1968), 73–78.
- B 89. B. H. Neumann, On characteristic morphisms, *J. Austral. Math. Soc.* **9** (1969), 478–488.
- B 90. B. H. Neumann, Remarque sur une note de A. Sade, *Univ. Lisboa Revista Fac. Ci. A* (2) **12** (1968/1969), 199–200.
- B 91. V. Dlab and B. H. Neumann, Semigroups with few endomorphisms, *J. Austral. Math. Soc.* **10** (1969), 162–168.
- B 91a. B. H. Neumann, Editorial, *Bull. Austral. Math. Soc.* **1** (1968), 1–2.
- B 91b. B. H. Neumann, Spreading the news, *Manifold* **8** (1970), 39–40.
- B 92. B. H. Neumann, Some remarks on cancellative semigroups, *Math. Z.* **117** (1970), 97–111.
- B 92a. B. H. Neumann, Quelques remarques sur les demi-groupes cancellatifs, *Algèbre et Théorie des nombres, Sem. P. Dubreil, M.-L. Dubreil-Jacotin, L. Lesieur et C. Pisot, année 23 (1969/70) Demi-groupes*, Exposé no.11, 2 pp., Secrétariat mathématique, Paris, 1970.
- B 93. B. H. Neumann, Algebraically closed semigroups, *Studies in Pure Mathematics* (Presented to Richard Rado), Academic Press, London, 1971, pp. 185–194.
- B 94. B. H. Neumann, Group Constructions, Faculty of Mathematics, University of Waterloo, Ontario, Canada, 1971, 25 pp.
- B 95. L. G. Kovács, Joachim Neubüser, and B. H. Neumann, On finite groups with ‘hidden’ primes, *J. Austral. Math. Soc.* **12** (1971), 287–300.
- B 95a. B. H. Neumann, Review: *Permutationstrukturen* by Olaf Tamaschke, *Zbl. Didaktik Math.* **4** (1971), 123.
- B 96. B. H. Neumann, Properties of countable character, Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1, Gauthier-Villars, Paris, 1971, pp. 293–296.
- B 97. B. H. Neumann, The isomorphism problem for algebraically closed groups, *Word problems: decision problems and the Burnside problem in group theory* (Conf. on Decision Problems in Group Theory, Univ. California, Irvine, Calif. 1969; dedicated to Hanna Neumann), *Studies in Logic and the Foundations of Math.*, vol. 71, North-Holland, Amsterdam, 1973, pp. 553–562.
- B 98. B. H. Neumann, Byron’s daughter, *Math. Gaz.* **57** (1973), no. 400, 94–97.
- B 99. B. H. Neumann, Group properties of countable character, *Selected questions of algebra and logic* (Collection dedicated to the memory of A. I. Mal’cev), Nauka, Novosibirsk, 1973, pp. 197–204.
- B 100. B. H. Neumann, Some groups I have known, *Proc. Second Internat. Conf. Theory of Groups* (Australian Nat. Univ., Canberra, 1973), Lecture Notes in Math., vol. 372, Springer-Verlag, Berlin, 1974, pp. 516–519.
- B 100a. B. H. Neumann, The day of a mathematician, *Austral. Math. Soc. Gaz.* **1** (1974), 72–73.
- B 100b. B. Neumann, Mathematics in New Zealand, *CSIRO Division of Mathematics and Statistics Newsletter* (1975), no. 10, 1.
- B 101. B. H. Neumann, Teaching teachers of teachers, *J. Japan. Soc. Math. Educ.* **57** (1975), Supplementary issue (Proc. ICME–JSME Regional Conference on Curriculum and Teacher Training for Mathematical Education, 1974), 24–28.
- B 102. J. L. Hickman and B. H. Neumann, A question of Babai on groups, *Bull. Austral. Math. Soc.* **13** (1975), no. 3, 355–368.
- B 103. B. H. Neumann, Questions and examples on group covering, *Acta Math. Acad. Sci. Hungar.* **26** (1975), no. 3–4, 291–293.
- B 104. B. H. Neumann, A problem of Paul Erdős on groups, *J. Austral. Math. Soc. Ser. A* **21** (1976), no. 4, 467–472.
- B 105. B. H. Neumann, The reflection principle, *Math. Chronicle* **5** (1976), no. 1–2, 1–7.
- B 106. B. H. Neumann, Some group presentations, *Canad. J. Math.* **30** (1978), no. 4, 838–850.
- B 107. B. H. Neumann, A problem of I. D. Macdonald, *Math. Gaz.* **62** (1978), no. 422, 298–299.
- B 107a. B. H. Neumann, Some new rumours in group theory, *Math. Medley* **6** (1978), no. 3, 100–103.
- B 107b. B. H. Neumann, On a problem of Ian D. Macdonald, *Math. Medley* **6** (1978), no. 3, 104–108.
- B 107c. B. H. Neumann, Proofs, *The Mathematical Intelligencer* **2** (1979), 18–19.
- B 107d. B. H. Neumann, Review: *Combinatorial Group Theory* by Roger C. Lyndon, Paul E. Schupp, *The Mathematical Intelligencer* **2** (1979), 42.
- B 108. B. H. Neumann and L. G. Wilson, Some sequences like Fibonacci’s, *Fibonacci*

- Quart.* **17** (1979), no. 1, 80–83; Corrigenda, *ibid.* **21** (1983), no. 3, 229.
- B 108a. B. H. Neumann, On some group presentations, *Word problems II* (Conf. on Decision Problems in Algebra, Oxford, 1976) (S. I. Adian, W. W. Boone, and G. Higman, eds.), Stud. Logic Foundations Math., vol. 95, North-Holland, Amsterdam, 1980, pp. 297–298.
- B 109. Ann Chi Kim, B. H. Neumann, and A. H. Rhemtulla, More Fibonacci varieties, *Bull. Austral. Math. Soc.* **22** (1980), no. 3, 385–395.
- B 110. B. H. Neumann, Formal power series, *Proc. Fourth Biennial Meeting of SEAMS on Modern Applications of Mathematics* (Bangkok, 1978) (Sawai Nualtaranee and Yupaporn Kemprasit, eds.), 1980, pp. 14–18.
- B 111. B. H. Neumann, Another single law for groups, *Bull. Austral. Math. Soc.* **23** (1981), no. 1, 81–102.
- B 112. B. H. Neumann, Not quite inner automorphisms, *Bull. Austral. Math. Soc.* **23** (1981), no. 3, 461–469.
- B 112a. B. H. Neumann, A supernova in Australian mathematical education, *Proc. 2nd South-East Asian Conf. on Mathematical Education* (Kuala Lumpur, 1981), pp. 6–8.
- B 112b. B. H. Neumann, Editor's introduction to J. Nielsen, On calculation with non-commutative factors and its applications in group theory, *Math. Scientist* **6** (1981), 73.
- B 112c. B. H. Neumann, Review: *Continued fractions: Analytic theory and applications* by William B. Jones and W. J. Thron, *Search* **12** (1981), 371.
- B 112d. B. H. Neumann, Review: *Emmy Noether, 1882–1935* by Auguste Dick (translated by H. I. Blocher), *Bull. London Math. Soc.* **14** (1982), 155–156.
- B 113. B. H. Neumann, Plane polygons revisited, *J. Appl. Probab.* (Special volume: Essays in statistical science) **19A** (1982), 113–122.
- B 114. A. Kertész, L. G. Kovács, and B. H. Neumann, Pure subgroups of non-abelian groups, *Publ. Math. Debrecen* **30** (1983), no. 1–2, 1–30.
- B 114a. B. H. Neumann, Review: *The history of combinatorial group theory: a case study in the history of ideas* by Bruce Chandler and Wilhelm Magnus, *Ann. Sci.* **41** (1984), 96–99.
- B 115. B. H. Neumann, Augustus De Morgan, *Bull. London Math. Soc.* **16** (1984), no. 6, 575–589.
- B 115a. A. C. Kim and B. H. Neumann, editors, *Groups—Korea 1983* (Proceedings of a conference on combinatorial group theory held at Kyoungju, August 26–31, 1983), Lecture Notes in Math., vol. 1098, Springer-Verlag, Berlin, 1984, viii+183 pp.
- B 116. B. H. Neumann, Commutative quandles, *Groups—Korea 1983* (Kyoungju, 1983), Lecture Notes in Math., vol. 1098, Springer-Verlag, Berlin, 1984, pp. 81–86.
- B 117. B. H. Neumann, Some finite groups with few defining relations, *J. Austral. Math. Soc. Ser. A* **38** (1985), no. 2, 230–240.
- B 118. B. H. Neumann, Yet another single law for groups, *Illinois J. Math.* **30** (1986), no. 2, 295–300.
- B 119. B. H. Neumann and Alf van der Poorten, Kurt Mahler: 1903–1988, *Austral. Math. Soc. Gaz.* **15** (1988), no. 2, 25–27.
- B 120. *Selected works of B. H. Neumann and Hanna Neumann*. Vol. I–VI, Selected Papers Series, Charles Babbage Research Centre, Winnipeg, MB, 1988, Edited by D. S. Meek and R. G. Stanton. With a foreword by Narain Gupta. With an obituary of Hanna Neumann by M. F. Newman and G. E. Wall, lii+1371 pp.
- B 121. I. D. Macdonald and B. H. Neumann, On commutator laws in groups, *J. Austral. Math. Soc. Ser. A* **45** (1988), no. 1, 95–103.
- B 122. J. L. Mennicke and B. H. Neumann, More on finite groups with few defining relations, *J. Austral. Math. Soc. Ser. A* **46** (1989), no. 1, 132–136.
- B 123. B. H. Neumann, Yet more on finite groups with few defining relations, *Group Theory* (Singapore, 1987), de Gruyter, Berlin, 1989, pp. 183–193.
- B 123a. A. C. Kim and B. H. Neumann, editors, *Groups—Korea 1988* (Proceedings of the Second International Conference on Group Theory held in Pusan, August 15–21, 1988), Lecture Notes in Math., vol. 1398, Springer-Verlag, Berlin, 1989, vi+189 pp.
- B 124. I. D. Macdonald and B. H. Neumann, On commutator laws in groups, 2, *Combinatorial group theory* (College Park, MD, 1988), Contemp. Math., vol. 109, Amer. Math. Soc., Providence, RI, 1990, pp. 113–129.

- B 125. Ann Chi Kim and B. H. Neumann, Laws in a class of groupoids, *Discrete Math.* **92** (1991), no. 1–3, 145–158.
- B 125a. B. H. Neumann, Review: *School mathematics: The challenge to change* by N. F. Ellerton and M. A. Clements, *Zbl. Didaktik Math.* **23** (1991), no. 4, 132–133.
- B 125b. B. H. Neumann, David Hilbert, *Math. Spectrum* **25** (1993), 70–73.
- B 126. B. H. Neumann, A large nilpotent group without large abelian subgroups, *Bull. London Math. Soc.* **25** (1993), no. 4, 305–308.
- B 126a. B. H. Neumann, Group-theoretic introduction, *Proc. Internat. Conf. Geometry Bangkok*, 21 January–2 February 1991 (V. Sa-yakanit, W. Sritrakool, et al., eds.), Chulalongkorn University Press, N.D., 1994, pp. 281–289.
- B 126b. B. H. Neumann, Plane transformations, *Proc. 22nd Annual Iranian Math. Conf.* 12–15 March 1991 (M. R. R. Moghaddam and M. A. Pourabdollah, eds.), Ferdowsi University of Mashhad, p. 262.
- B 126c. B. H. Neumann, Commutator laws, *Proc. 22nd Annual Iranian Math. Conf.* 12–15 March 1991 (M. R. R. Moghaddam and M. A. Pourabdollah, eds.), Ferdowsi University of Mashhad, p. 262.
- B 126d. B. H. Neumann, What use is mathematics?, *Quality Mathematics Education in Developing Countries* (Proc. South Pacific Conf. on Math. and Math. Education, Univ. Papua New Guinea, 24–26 June 1992) (O. P. Ahuja, J. C. Renaud, and R. M. Sekkappan, eds.), UBS Publishers’ Distributors, New Delhi etc, 1995, pp. 1–3.
- B 126e. B. H. Neumann, Plane transformations, *Quality Mathematics Education in Developing Countries* (Proc. South Pacific Conf. on Math. and Math. Education, Univ. Papua New Guinea, 24–26 June 1992) (O. P. Ahuja, J. C. Renaud, and R. M. Sekkappan, eds.), UBS Publishers’ Distributors, New Delhi etc, 1995, pp. 201–220.
- B 126f. B. H. Neumann, Interaction between German and Australian mathematicians, *German-Australian Cultural Relations Since 1945* (Brisbane, September 20–23, 1994) (Manfred Jurgensen, ed.), German-Australian Studies, vol. 9, Peter Lang, Bern etc, pp. 321–325.
- B 127. I. D. Macdonald and B. H. Neumann, On commutator laws in groups, *J. Austral. Math. Soc. Ser. A* **58** (1995), no. 1, 126–133.
- B 128. J. T. Buckley, John C. Lennox, B. H. Neumann, Howard Smith, and James Wiegold, Groups with all subgroups normal-by-finite, *J. Austral. Math. Soc. Ser. A* **59** (1995), no. 3, 384–398.
- B 129. B. H. Neumann, Covering groups by subgroups, *Groups—Korea ’94* (Pusan), de Gruyter, Berlin, 1995, pp. 249–250.
- B 130. B. H. Neumann, Hardy – some reminiscences, *Austral. Math. Soc. Gaz.* **25** (1998), no. 3, 138–141.
- B 131. L. G. Kovács, B. H. Neumann, and M. F. Newman, A problem of Grätzer and Wehrung on groups, *Southeast Asian Bull. Math.* **22** (1998), no. 2, 155–156.
- B 132. B. H. Neumann, Commutator laws in algebraic systems, *Semigroups* (Kunming, 1995), Springer, Singapore, 1998, pp. 245–250.
- B 133. B. H. Neumann, Implicative identities in groups, *Algebras and combinatorics* (Hong Kong, 1997), Springer, Singapore, 1999, pp. 359–365.
- B 133a. B. H. Neumann, Review: *Memoirs of a maverick mathematician* by Zoltan Paul Dienes, *Math. Gaz.* **84** (2000), 348–350.
- B 134. B. H. Neumann, Some semigroup laws in groups, *Canad. Math. Bull.* **44** (2001), no. 1, 93–96.
- B 135. B. H. Neumann, Ensuring commutativity of finite groups, *J. Aust. Math. Soc.* **71** (2001), no. 2, 233–234, Special issue on group theory.
- B 135a. B. H. Neumann, An ancient mathematician remembers, *Newsletter, British Soc. for the History of Math.* **43** (2001), 3–7.
- B 135b. B. H. Neumann, Review: *Calls from the past* by Zoltan Paul Dienes, *Math. Gaz.* **85** (2001), 534; **86** (2002), 347.
- B 135c. B. H. Neumann, An ancient mathematician remembers, *Math. Spectrum* **35** (2002/2003), 2–3.
- B 136. Bernhard Neumann, Some personal recollections, *Studies in Memory of Issai Schur* (Chevaleret/Rehovot, 2000), Progr. Math., vol. 210, Birkhäuser Boston, Boston, MA, 2003, pp. xxxi–xxxii.
- B 136a. B. H. Neumann, Review: *Notable women in mathematics, a biographical dictionary* by Charlene Morrow and Teri Perl; Review: *Women becoming mathematicians. Creating a professional identity in post-world war II America* by Margaret A. M. Murray, *Math. Gaz.* **87** (2003), 183–184.