


CSIRO Publishing



Marine & Freshwater Research

VOLUME 53, 2002

© CSIRO 2002

A journal for the publication of original contributions
in physical oceanography, marine chemistry,
marine and estuarine biology and limnology

All enquiries and manuscripts should be directed to:

Marine and Freshwater Research
CSIRO Publishing
PO Box 1139 (150 Oxford St)
Collingwood, Vic. 3066, Australia



CSIRO
PUBLISHING

Telephone: +61 3 9662 7618
Fax: +61 3 9662 7611
Email: publishing.mfr@csiro.au

Published by CSIRO Publishing
for CSIRO and the Australian Academy of Science

www.publish.csiro.au/journals/mfr

Table. Summary of tagging results for non-migratory, sub-tidal reef fishes showing proportion that had moved away from the tagging site, the maximum distance traveled and source of information

Temperate^T and juvenile^J study subjects are noted accordingly

Species	% Moved	Max. Distance	N	Source
Acanthuridae				
	6%	0.5 km	65	Randall (1962)
<i>Acanthurus bahianus</i>				
	4%	0.4 km	25	Randall (1962)
<i>A. chirurgus</i>				
	18%	0.5km	44	Randall (1962)
<i>A. coeruleus</i>				
Lutjanidae				
<i>Lutjanus campechanus</i>	6%	5 km	17	Fable (1980)
<i>Lutjanus campechanus</i>	25%	32 km	37	Szedlmayer and Shipp (1994)
<i>L. apodus</i>	22%	1.3 km	18	Randall 1962
<i>Rhombolites aurorubens</i>	5%	10-20 km	39	Fable (1980)
	27%	11 km	11	Randall (1962)
<i>Ocyurus chrysurus</i>				
Scaridae				
	10%	13 km	20	Randall (1962)
<i>Scarus coeruleus</i>				
Scorpaenidae				
<i>Sebastes caurinus</i> ^T	0%	0	11	Mathews and Barker (1983)
<i>S. caurinus</i> ^T	0%	0	75	Gowan (1983) in Mathews and Barker (1983)
<i>S. maliger</i> ^T	9%	2.8km	11	Mathews and Barker (1983)
<i>S. melanops</i> ^T	38%	400km	8	Mathews and Barker (1983)
<i>S. flavidus</i> ^{JT}	80%	144km	10	Mathews and Barker (1983)

Serranidae

<i>Cephalopholis fulva</i>	3.4%	0.5 km	29	Randall (1962)
<i>Epinephelus adscensionis</i>	8%	1.8 km	12	Randall (1962)
<i>E. guttatus</i>	8.3%	0.5 km	24	Randall (1962)
<i>E. morio</i>	28%	100 km	403	Wilson and Burns (1996)
<i>E. striatus</i>	6.7%	+ 50 km	74	Randall (1962)
<i>Mycteroperca microlepis</i>	14%	458 km	132	Heinisch and Fable (1999)
<i>M. phenax</i>	53%	-	22	Wilson and Burns (1996)
<i>M. venosa</i>	22%	6.4 km	27	
<i>Plectropomus leopardus</i>	10%			Beinssen 1989 in Roberts and Polunin 1991

Sparidae

<i>Chrysoblephus laticeps</i> ^T	26%	39 km	240	This study
<i>C. laticeps</i>	36%	2 km	11	Buxton and Allen (1989)
<i>C. gibbiceps</i> ^T	7%	33 km	70	This study
<i>Petrus rupestris</i> ^{IT}	33%	1000 km	111	This study
<i>Chiemerius nufar</i>	43%	84 km	240	This study
<i>Argyrozona argyrozona</i> ^{IT}	32%	95 km	44	This study
Inshore				
<i>A. argyrozona</i> ^J	43%	300 km	23	Nepgen (1977)
Inshore				
<i>A. argyrozona</i>	16%	110 km	158	This study
Offshore				
<i>Pachymetopon blochii</i> ^T	43%	40 km	7	Nepgen (1977)
<i>Pagrus auratus</i>	21%	120 km	227	Jones (1981)

References listed in Table

- Buxton, C.D., and Allen J.C.** (1990). Mark and recapture studies of two reef sparids in the Tsitsikamma Coastal National Park. *Koedoe* **32**, 39-45.
- Fable, W.A.** (1980). Tagging studies of red snapper (*Lutjanus campechanus*) and vermilion snapper (*Rhombolites aurorubens*) off the south Texas coast. *Contributions to Marine Science* **23**, 115-121.
- Heinisch, B.V., and Fable, W.A.** (1999). Movement of gag, *Mycteroperca microlepis*, in and from St. Andrew Bay, Florida. *Bulletin of Marine Science* **64**, 501-508.
- Jones, K.** (1981). Biological research on snapper (*Chrysophrys auratus* sun. unicolor) and an analysis of the fishery in northern Spencer Gulf. *SAFIC* **5**(6), 5-8.
- Mathews, S.B., and Barker, M.W.** (1983). Movements of rockfish (*Sebastes*) in Northern Puget Sound, Washington. *Fishery Bulletin U.S.* **82**, 916-922.
- Nepgen C.S. DeV.** (1977). The biology of the Hottentot *Pachymetopon blochii* (Val.) and the Silverfish *Argyrozona argyrozona* (Val.) along the south-west coast. *Investigational Reports of the Division of Sea Fisheries, South Africa* **105**, 1-35.

- Randall, J.E.** (1962). Tagging reef fishes in the Virgin Islands. In (J.B. Higman ed) Gulph and Caribbean Fisheries Institute, Proceedings of the 14th Annual Session, University of Miami., Florida, November 1961. 201-241.
- Roberts, C.M., and Polunin, N.V.C.** (1991). Are marine reserves effective in management of reef fisheries? *Reviews in Fish Biology and Fisheries* **1**, 65-91.
- Szedlmayer, S.T., and Shipp, R.L.** (1994). Movement and growth of red snapper, *Lutjanus campechanus*, from an artificial reef area in the northeastern Gulf of Mexico. *Bulletin of Marine Science* **55**, 887-896.
- Wilson, R.R., and Burns, K.M.** (1996). Potential survival of released groupers caught deeper than 40m based on shipboard and in-situ observations, and tag-recapture data. *Bulletin of Marine Science* **58**, 234-247.