

Supplementary material

Immature individuals dominate elasmobranch fisheries of the Bali Strait

Sara Tabea Winter^{A,H}, *Fahmi*^B, *Didik Rudianto*^C, *Betty J. L. Laglbauer*^{C,D,E}, *Isabel Ender*^{A,F}
and *Colin A. Simpfendorfer*^{A,G}

^ACollege of Science and Engineering, James Cook University,
Townsville, Qld 4811, Australia.

^BResearch Centre for Oceanography, LIPI (Indonesian Institute of Sciences),
Jakarta 14430, Indonesia.

^CMobula Project Indonesia, Jl. Brawijaya, Gang Kurnia, Dusun Krajan Desa Tembokrejo,
Muncar, Banyuwangi 68472, Jawa Timur, Indonesia.

^DUniversity of the Azores, Rua Frederico Machado, 9900-862, Horta, Portugal.

^ESchool of Biomedical Science, The University of Queensland,
Saint Lucia, Qld 4072, Australia.

^FManta Trust, Catemwood House, Corscombe, Dorchester, Dorset, DT2 0NT, UK.

^GCentre for Sustainable Tropical Fisheries and Aquaculture, James Cook University,
Townsville, Qld 4811, Australia.

^HCorresponding author. Email address: sara.winter1@my.jcu.edu.au




Fig. S1. Species specific contribution to the monthly catch of the 15 most commonly caught species landed in Muncar from September 2017 to February 2018.