

Do vertebral chemical signatures distinguish juvenile blacktip shark (*Carcharhinus limbatus*) nursery regions in the northern Gulf of Mexico?

Justin P. Lewis^{A,B,E}, William F. Patterson, III^{A,B}, John K. Carlson^C
and Katherine McLachlin^D

^AUniversity of South Alabama, Department of Marine Sciences, 307 North University Boulevard, Mobile, AL 36688, USA.

^BDauphin Island Sea Lab, 101 Bienville Boulevard, Dauphin Island, AL 36528, USA.

^CNational Marine Fisheries Service, Southeast Fisheries Science Center, Panama City Laboratory, 3500 Delwood Beach Road, Panama City, FL 32408, USA.

^DElectro Scientific Industries, 685 Old Buffalo Trail, Bozeman, MT 59715, USA.

^ECorresponding author. Present address: Center for Fisheries Research and Development, Gulf Coast Research Laboratory, The University of Southern Mississippi, 703 East Beach Drive, Ocean Springs, MS 39564, USA. Email: jplewis86@gmail.com

Abstract. Identifying and protecting shark nurseries is a common management strategy used to help rebuild overfished stocks, yet we know little about connectivity between juvenile and adult populations. By analysing trace metals incorporated into vertebral cartilage, it may be possible to infer natal origin based on nursery-specific chemical signatures. To assess the efficacy of this approach, we collected juvenile blacktip sharks (*Carcharhinus limbatus*; $n = 93$) from four regions in the Gulf of Mexico in 2012 and 2013 and analysed their vertebral centra with laser ablation–inductively coupled plasma–mass spectrometry. We observed significant regional differences in six element : Ca ratios in both 2012 and 2013. Multi-element chemical signatures were significantly different among regions and between year-classes. Year-class-specific linear discriminant function analysis yielded regional classification accuracies of 81% for 2012 and 85% for 2013, although samples were not obtained from all four regions in 2012. Combining year-classes resulted in an overall classification accuracy of 84%, thus demonstrating the usefulness of this approach. These results are encouraging yet highlight a need for more research to better evaluate the efficacy of vertebral chemistry to study elasmobranch population connectivity.

Additional keywords: laser ablation, natal origin, shark nurseries, vertebral chemistry.

Received 1 March 2015, accepted 1 December 2015, published online 11 February 2016

Introduction

Nurseries are areas that contribute a disproportionate number of juvenile recruits to adult populations, thus the conservation benefit of protecting nurseries is evident. However, inconsistent application of the term ‘nursery’ prompted the development of guidelines to empirically compare the value of different habitats or systems in the context of population maintenance and growth (Beck *et al.* 2001; Heupel *et al.* 2007). Proxies that indicate an area provides favourable conditions for subsequent recruitment to the adult population (e.g. high juvenile density, growth rate and survival) are important for understanding the ecological processes of source versus sink regions (for a review, see Beck *et al.* 2001). However, the ultimate character that distinguishes a functional nursery is a disproportionately high contribution of recruits to the adult population, thus determining where adults spent their early life is the ultimate test of whether a given habitat or system is, in fact, a nursery.

Identifying nursery areas based on estimates of connectivity to adult populations is particularly challenging for coastal shark species. Conventional mark–recapture studies are limited because recapture rates are typically low (e.g. $\leq 10\%$; Kohler and Turner 2001) and home ranges in the order of tens of square kilometres (e.g. Yeiser *et al.* 2008) make it difficult to mark enough individuals to compensate for high mortality during the first year of life. In the case of bony fishes, natural biogeochemical tags based on chemical constituents of calcified structures (e.g. otoliths, scales, fin rays and bone) provide an alternative approach to study movement patterns and population connectivity (for a review, see Elsdon *et al.* 2008). During the biomineralisation process, divalent cations, such as Sr^{2+} and Ba^{2+} , substitute for Ca^{2+} relative to their environmental availability (Wells *et al.* 2000), whereas other elements may become incorporated interstitially or associated with the protein matrix (Campana 1999). Thus, the chemical constituents of calcified structures are representative

of environmental conditions at the time of deposition. In bony fishes, otoliths are the preferred structure for chemical analysis of natural biogeochemical tags because they are inert once formed and their biomineralisation pattern produces alternating translucent and opaque zones that provide a chronometer that accompanies the recorded chemical histories (Campana 1999). Material at the core of the otolith represents deposition during the first year of life; thus, chemical signatures found in the core can act as a natural tag to be used to estimate the proportion of adults derived from different nurseries (e.g. Vasconcelos *et al.* 2011).

The incorporation of trace metal impurities into calcified structures of other marine organisms has also been shown to reflect the environmental conditions at the time of mineralisation (e.g. Pitts and Wallace 1994) and may be used to address a variety of ecological questions. Results from recent field (Scharer *et al.* 2012) and experimental (Smith *et al.* 2013) studies show some elements deposited in the hydroxyapatite matrix ($\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$) of elasmobranch vertebrae are also influenced by abiotic factors (e.g. salinity, water chemistry and temperature). These results indicate that chemical signatures in vertebrae may serve as natural biogeochemical tags of elasmobranch juvenile habitat that could be used to examine sources of recruits to adult populations. However, their usefulness for studying connectivity between juvenile and adult populations is contingent on our ability to accurately classify individuals to their natal nurseries based on vertebral chemical signatures.

In the present study, blacktip sharks (*Carcharhinus limbatus*) were examined as a model coastal shark species to test whether vertebral chemical signatures distinguish nursery regions in the northern Gulf of Mexico (Gulf) and whether differences can be used to infer nursery origin. The blacktip shark is a cosmopolitan species that is found throughout coastal waters of the Gulf and is one of the more economically important shark species in the region (Southeast Data, Assessment, and Review 2012). Females reproduce biennially following a gestation period of ~12 months, with parturition peaking from March to May in coastal nurseries throughout the Gulf (Baremore and Passerotti 2013). Juvenile blacktips remain in natal coastal embayments up to the first 6 months of life before migrating to wintering grounds (Heupel *et al.* 2004; Hueter and Tyminski 2007). Tagging data indicate individuals up to the age of 3 years return to their nurseries during summer months (Heupel and Simpfendorfer 2002; Hueter *et al.* 2005), whereas analysis of mitochondrial (mt)DNA and nuclear DNA microsatellites indicates restricted gene flow that is consistent with female philopatry to natal regions (Keeney *et al.* 2005). Development of region-specific natural biogeochemical tags based on vertebral chemistry may permit further testing of site fidelity and philopatry for this species, as well as facilitate examination of connectivity between juvenile and adult populations.

Materials and methods

Sample collection and preparation

Juvenile blacktip sharks were collected opportunistically in 2012 and 2013 (Table 1) during the Gulf of Mexico States Shark Pupping and Nursery Survey (GULFSPAN). The GULFSPAN survey is a fisheries-independent gill net survey that was initiated by the National Marine Fisheries Service (NMFS) Laboratory in

Table 1. Sample size and size ranges of juvenile blacktip sharks (2012 and 2013 year-classes) from four nursery regions in the northern Gulf of Mexico (Gulf)

Average size at birth for blacktip sharks in the Gulf is 380-mm fork length (FL) (Baremore and Passerotti 2013). EG, eastern Gulf; NEG, north-eastern Gulf; NCG, north-central Gulf; WG, western Gulf

Year-class	Region	Sample size	Size range (mm FL)
2012	NEG	14	465–610
	NCG	15	530–660
	WG	11	604–693
2013	EG	19	420–580
	NEG	10	430–570
	NCG	7	499–589
	WG	16	503–666

Panama City (FL, USA) in 2003 and occurs annually within several coastal areas of the Gulf. Samples were also obtained from fisheries-independent surveys by the Mote Marine Laboratory (Sarasota, FL, USA), Florida State University Coastal and Marine Laboratory (St Teresa, FL, USA), Dauphin Island Sea Laboratory (Dauphin Island, AL, USA), University of Southern Mississippi Gulf Coast Research Laboratory (Ocean Springs, MS, USA) and Texas Parks and Wildlife Department (Port O'Connor, TX, USA) using gill nets or longlines. Samples of the 2012 and 2013 year-classes were collected from four regions in the Gulf (Fig. 1): (1) the eastern Gulf (EG) off the south-west coast of Florida; (2) the north-eastern Gulf (NEG) from the Big Bend west through the Florida panhandle; (3) the north-central Gulf (NCG) within the state boundaries of Alabama and Mississippi; and (4) the western Gulf (WG) in San Antonio Bay, Texas.

Thoracic vertebrae were removed in the field or in the laboratory and stored frozen until processed. Excess tissue and haemal and neural arches were excised with a scalpel (Fig. 2a). Individual vertebrae were sectioned along the sagittal plane with a Buehler (Lake Bluff, IL, USA) Isomet low-speed saw, resulting in ~0.5-mm sections (Fig. 2b). Vertebral sections were placed in acid-leached polystyrene cells filled with ultrapure water (18 MΩ cm⁻¹) and sonicated for 1 h. Following sonication, sections were rinsed with ultrapure water, transferred to acid-leached cell wells and dried under a Class 10 laminar flow clean hood.

Sample analysis

Vertebral chemistry was analysed with laser ablation–inductively coupled plasma–mass spectrometry (LA-ICP-MS). The LA-ICP-MS system consisted of a Nd:YAG NWR213 (Electro Scientific Industries, Portland, OR, USA) laser ablation system coupled with an Agilent Technologies (Santa Clara, CA, USA) 7700x ICP-MS. Helium (flow rate 0.7 L min⁻¹) was used to sweep ablated material from the ablation cell then mixed with argon (flow rate 0.79 L min⁻¹) and transported the ICP-MS. Sample order was randomised and LA-ICP-MS analysis consisted of three parallel laser ablation transects through the corpus calcareum (Fig. 2c). Transects were spaced 120 μm apart and each transect was preablated (laser spot size 100 μm, speed 110 μm s⁻¹, repetition rate 5 Hz, and fluence 0.41 J cm⁻²) before analysis. Ablation transects consisted of a spot size of 80 μm, a scan speed of 11 μm s⁻¹, a repetition rate of 10 Hz and fluence of




Fig. 1. Sampling locations of juvenile blacktip sharks in the Gulf of Mexico.




Fig. 2. Digital images of a vertebral centrum from a juvenile blacktip shark. (a) Cleaned whole centrum, (b) sagittal section and (c) the corpus calcareum, with white lines denoting ablation transects.

4.0 J cm^{-2} . Background counts were collected 30 s before each transect and subtracted from the signal. Reference materials from the United States Geological Survey (USGS MACS3) and National Institute of Standards and Technology (NIST 612) were ablated every 45 min to evaluate analytical precision and correct for instrument drift. Counts per second (CPS) were converted to units of concentration using the Trace ElementIS data reduction scheme in Iolite (version 2.5, School of Earth Sciences, University of Melbourne, Melbourne, Australia) using Ca (35 wt% in shark vertebrae; Tillett *et al.* 2011) as an internal standard. We monitored eight elements (^7Li , ^{24}Mg , ^{43}Ca , ^{44}Ca ,

^{31}P , ^{55}Mn , ^{88}Sr , ^{137}Ba and ^{208}Pb). Because the concentration of ^{31}P is not certified in the MACS3, it was excluded from subsequent analyses. The concentrations of the remaining elements were normalised to ^{43}Ca (element:Ca) and are expressed as micromoles per mole (Li:Ca, Mn:Ca, Ba:Ca, Pb:Ca) or millimoles per mole (Mg:Ca and Sr:Ca).

Data analysis

The portion of each transect that represented postnatal growth was measured with a Nikon (Tokyo, Japan) SMZ1500 dissecting microscope and Nikon's NIS-elements BR 4.00.07 imaging

software. The minimum amount of postnatal vertebral tissue analysed among samples was 170 μm . Therefore, data from the first 170 μm immediately following the birth mark were averaged among the three transects of each vertebral section to estimate chemical signatures. Assuming a mean birthday of 1 May (Carlson *et al.* 2006), this region represents ~ 1 month of life.

Univariate normality was assessed graphically and homogeneity of variance was tested using Levene's test. Individual element : Ca ratios that did not meet parametric assumptions were natural log transformed, but transformation of Li : Ca and Pb : Ca for the 2013 year-class did not result in equal variances among regions. Because the results of non-parametric and parametric univariate tests yielded the same significance and produced identical multiple comparison results, we only present the results of parametric tests. One multivariate outlier was identified and removed from the dataset. Graphical assessment of multivariate dispersion using the Euclidian distances confirmed that homogeneity of variances was met for the multivariate dataset when using raw data of Li : Ca and Pb : Ca.

Differences in multi-elemental signatures among regions (excluding the EG, which was only sampled in 2013) and between years were tested with a two-way multivariate analysis of variance (MANOVA) with Pillai's Trace as the test statistic. Regional and year-class differences for individual element : Ca ratios were tested with a two-way analysis of variance (ANOVA), again for all regions except the EG. A second MANOVA was computed among all regions sampled in 2013, which was followed by an ANOVA for individual element : Ca ratios. Significant pairwise differences in element : Ca ratios within year-classes were identified with Tukey's honestly significant difference (HSD) tests. A linear discriminant function analysis (LDFA) was computed for each year-class separately, and then for all data combined, to assess the ability of vertebral chemical signatures to distinguish sample regions. The accuracy of LDFA models was assessed via cross-validated classification accuracies. Initial LDFAs included all six elements; however, previous studies of the abiotic and biotic factors that influence vertebral chemistry suggest Mn : Ca is primarily derived from dietary sources and may be influenced by internal nutrient reserves (see Discussion). Thus, LDFAs were also run without Mn : Ca. All statistical analyses were performed in R version 3.2.1 (R Foundation for Statistical Computing, Vienna, Austria).

Results

Each element assayed was consistently above detection limits, with the exception of Pb. Values of Pb below detection were not common (0.1%) within the region of interest and, when present, were replaced with the limits of detection (LOD; Table 2). Relative standard deviations (%RSD) were 3.7–5.1% for the NIST 612 standard and 8.8–10.1% for the MACS3 standard (Table 2).

Multi-element chemical signatures differed significantly among the three regions sampled in 2012 and 2013 ($F_{12,126} = 29.5$, $P < 0.001$) and between years ($F_{6,62} = 6.66$, $P < 0.001$), with a significant interaction between factors ($F_{12,126} = 2.43$, $P = 0.007$; Table 3). There was also a significant

Table 2. Estimated precision (relative standard deviation (%RSD)) and limits of detection (LOD) for elements analysed in blacktip shark vertebrae with laser ablation–inductively coupled plasma mass spectrometry

%RSD was computed for the glass bead standard produced by the National Institute of Standards and Technology (NIST 612, Charleston, SC, USA) and pressed carbonate standard produced by the United States Geological Survey (USGS MACS3, Denver, CO, USA), and LOD was computed using MACS3

Element	%RSD		LOD ($\mu\text{mol mol}^{-1}$)
	NIST 612	MACS3	
Li	5.0	8.9	0.039
Mg	4.5	8.8	0.084
Mn	3.7	9.5	0.027
Sr	4.4	8.5	0.016
Ba	5.1	9.3	0.006
Pb	4.8	10.1	0.008

Table 3. Two-way and one-way multivariate analysis of variance (MANOVA) results for tests of regional and interannual differences in multi-element signatures for juvenile blacktip sharks

Both multivariate analyses included Li : Ca, Mg : Ca, In(Mn : Ca), Sr : Ca, Ba : Ca and Pb : Ca. The two-way MANOVA includes age-0 sharks collected in the north-eastern Gulf of Mexico (NEG), north-central Gulf of Mexico (NCG) and western Gulf of Mexico (WG) in 2012 and 2013, whereas the one-way MANOVA includes sharks collected from the eastern Gulf of Mexico, NEG, NCG and WG in 2013 P -values in bold indicate significant differences. d.f., degrees of freedom

Test or factor	Pillai's trace	F	d.f.	P -value
Two-way MANOVA				
Region	1.47	29.5	12, 126	<0.001
Year	0.40	6.66	6, 62	<0.001
Region \times year	0.39	2.43	12, 126	0.007
One-way MANOVA				
Region	1.97	14.43	18, 135	<0.001

difference in multi-element signatures among regions sampled in 2013 ($F_{18,135} = 14.43$, $P < 0.001$; Table 3).

Results of the two-way ANOVA (Table 4) indicated all elemental ratios were significantly different among regions ($P < 0.001$), but only Mg : Ca exhibited a significant difference between year-classes ($P < 0.001$). There were significant interactions between region and year-class for Li : Ca and Mn : Ca ($P \leq 0.03$). Regional differences in mean element : Ca were also observed among the four regions sampled in 2013 (Table 5) and showed the same general trend observed in 2012 (Fig. 3). Sharks from the NCG tended to have the lowest mean Li : Ca, Mg : Ca and Sr : Ca and highest mean for Mn : Ca, whereas sharks from the WG had the highest mean Li : Ca, Ba : Ca and Pb : Ca. Sharks from the NEG had the highest mean vertebral Mg : Ca and Sr : Ca and lowest values for Ba : Ca and Pb : Ca. Both NEG and EG exhibited similar element : Ca ratios, with the exception of Mn : Ca ($P < 0.001$). In fact, Mn : Ca was the only elemental ratio that was unique for each region within year-class ($P < 0.05$).

Table 4. Two-way analysis of variance (ANOVA) results for tests of regional and interannual differences in element : Ca ratios for juvenile blacktip sharks

The two-way ANOVA includes age-0 sharks collected in the north-eastern Gulf of Mexico, north-central Gulf of Mexico and western Gulf of Mexico in 2012 and 2013. *P*-values in bold indicate significant differences. d.f., degrees of freedom; MS, mean sum of squares

Ratio	Factor	d.f.	MS	<i>F</i>	<i>P</i> -value
Li : Ca	Region	2	1.19×10^{-9}	42.62	<0.001
	Year	1	5.94×10^{-12}	0.42	0.51
	Region \times year	2	1.71×10^{-10}	6.11	0.004
	Residuals	67	9.39×10^{-10}		
Mg : Ca	Region	2	9.77×10^{-5}	11.47	<0.001
	Year	1	1.38×10^{-4}	32.39	<0.001
	Region \times year	2	8.97×10^{-6}	1.05	0.35
	Residuals	67	2.85×10^{-4}		
In(Mn : Ca)	Region	2	12.91	70.6	<0.001
	Year	1	0.08	0.92	0.34
	Region \times year	2	0.85	4.66	0.013
	Residuals	67	6.13		
Sr : Ca	Region	2	1.20×10^{-6}	50.41	<0.001
	Year	1	2.2×10^{-10}	0.02	0.890
	Region \times year	2	2.21×10^{-8}	0.93	0.400
	Residuals	67	7.94×10^{-7}		
Ba : Ca	Region	2	1.75×10^{-10}	21.47	<0.001
	Year	1	6.8×10^{-14}	0.02	0.9
	Region \times year	2	4.4×10^{-13}	0.05	0.95
	Residuals	67	2.73×10^{-10}		
Pb : Ca	Region	2	3.34×10^{-13}	42.73	<0.001
	Year	1	4.36×10^{-15}	1.11	0.29
	Region \times year	2	2.01×10^{-14}	2.67	0.08
	Residuals	67	2.62×10^{-13}		

Table 5. One-way analysis of variance (ANOVA) results for tests of regional differences in element : Ca ratios for juvenile blacktip sharks

The one-way ANOVA includes sharks collected from the eastern Gulf of Mexico, north-eastern Gulf of Mexico, north-central Gulf of Mexico and western Gulf of Mexico in 2013. *P*-values in bold indicate significant differences. d.f., degrees of freedom; MS, mean sum of squares

Ratio	Factor	d.f.	MS	<i>F</i>	<i>P</i> -value
Li : Ca	Region	3	1.22×10^{-9}	37.07	<0.001
	Residuals	48	5.26×10^{-10}		
Mg : Ca	Region	3	9.61×10^{-5}	8.01	<0.001
	Residuals	48	1.92×10^{-4}		
In(Mn : Ca)	Region	3	26.75	97.27	<0.001
	Residuals	48	4.4		
Sr : Ca	Region	3	1.97×10^{-6}	20.64	<0.001
	Residuals	48	1.52×10^{-6}		
Ba : Ca	Region	3	7.40×10^{-11}	6.23	0.001
	Residuals	48	1.89×10^{-10}		
Pb : Ca	Region	3	4.52×10^{-13}	51.23	<0.001
	Residuals	48	1.44×10^{-13}		

Mean cross-validated classification accuracies from LDFAs including all elemental ratios were 80, 90 and 88% for 2012, 2013 and both year-classes combined respectively (Fig. 4). Excluding Mn : Ca yielded a slightly higher mean classification accuracy of 81% for 2012. However, the exclusion of Mn : Ca




Fig. 3. Mean (\pm 95% confidence interval) region and year-class-specific element : Ca ratios of blacktip shark vertebrae analysed in 2012 (grey) and 2013 (white). WG, western Gulf of Mexico (Gulf); NCG, north-central Gulf; NEG, north-eastern Gulf; EG, eastern Gulf.

yielded lower mean classification accuracies of 84 and 85% for the 2013 year-class and both year-classes combined respectively (Fig. 4). The lowest classification accuracy was observed for WG samples of the 2012 year-class (54%), whereas the highest classification accuracy was for NCG samples of the 2013 year-class (100%).

Discussion

Results from the present study demonstrate the potential of using natural biogeochemical tags in juvenile blacktip shark vertebrae to distinguish Gulf nursery regions. Although water chemistry and other environmental factors that likely drive these biogeochemical signatures were not available for all regions, spatial differences in environmental conditions were clearly sufficient to produce distinct signatures among regions of interest (for a review, see *Elsdon et al. 2008*). Earlier attempts to distinguish shark nursery regions based on vertebral biogeochemical signatures had mixed success at similar spatial scales (e.g. hundreds to thousands of kilometres). *Smith (2013)* reported that vertebral signatures of juvenile scalloped hammerhead sharks (*Sphyrna lewini*) collected from three nursery areas along the Pacific coast of Mexico produced site-specific classification accuracies ranging from 26 to 80%. *Tillett et al. (2011)* reported significant differences among the vertebral chemistry of bull sharks (*Carcharhinus leucas*) among six




Fig. 4. Cross-validated classification accuracies (%) of linear discriminant function analyses using Li : Ca, Mg : Ca, In(Mn : Ca), Sr : Ca, Ba : Ca, Pb : Ca (top row) and Li : Ca, Mg : Ca, Sr : Ca, Ba : Ca, Pb : Ca (bottom row). Vertical panels represent year-class 2012, 2013 and both year-classes combined, as indicated. Circle diameter represents classification success. Black circles indicate sharks accurately classified to their natal region, with circles becoming progressively lighter with increased distance between the collection region and classification region. WG, western Gulf of Mexico (Gulf); NCG, north-central Gulf; NEG, north-eastern Gulf; EG, eastern Gulf.

nurseries in Australia, but noted that there was no difference in the vertebral chemistry of pigeye sharks (*Carcharhinus amboinensis*). This is not too surprising considering that neonate bull sharks frequent areas of lower salinity than the pigeye shark (Knip *et al.* 2011). Similar to neonate and juvenile pigeye sharks, blacktip and scalloped hammerhead sharks also prefer higher salinities than neonate bull sharks (Hueter and Tyminski 2007; Parsons and Hoffmayer 2007; Steiner *et al.* 2007). Thus, it appears that watersheds, basin geology or local biogeochemical cycling were more distinct among regions examined in the present study compared with the previous work by Tillett *et al.* (2011) and Smith (2013).

The high mobility of juvenile individuals can also create problems when applying natural biogeochemical tags because an assumption of this approach is that samples of a target group or region have not mixed with other groups or regions (Elsdon *et al.* 2008). Thus, the usefulness of these tags will be constrained by spatial scales over which minimal mixing occurs between adjacent areas of interest. Heupel *et al.* (2004) reported that young-of-year blacktip sharks in Terra Ceia Bay (FL, USA) had restricted core home ranges (mean monthly 50% kernel densities $\leq 0.5 \text{ km}^2$) throughout the summer, but the degree to which they ventured outside their core home range increased in

the month of July. Given the relatively large distance between regions in the present study and the high degree of site fidelity and limited home ranges reported by Heupel *et al.* (2004) for juvenile blacktips, the probability we sampled migrants from adjacent regions is extremely low. Because the activity space of juvenile blacktip sharks increases throughout their first summer of life, focussed sample collection during the early months of the birth season will increase the probability an individual is captured in its natal estuary. By increasing sampling effort in and around putative nurseries, it may be possible to determine the spatial resolution at which one can accurately assign natal origin.

Failing to account for intra-annual variability, especially in populations that exhibit a protracted spawning and birth season, can have a strong effect on classification accuracies and estimates of connectivity (Cook 2011; Smith 2013). In the present study, almost all sharks exhibited temporal changes in vertebral chemistry, which reflect temporal variability in abiotic (temperature and water chemistry) and biotic (expanding home range) factors. Our limited sample size precludes statistical analysis of how classification accuracies vary when sampling different regions along the vertebrae or whether sharks born at different times within an estuary have different natal signatures (Smith

2013). The duration of the birth season for blacktip sharks in the Gulf is ~2 months (Baremore and Passerotti 2013), which is much shorter than that of the scalloped hammerhead population along the Pacific coast of Mexico, which occurs between May and October (Smith 2013). Hamer *et al.* (2003) reported that separating juveniles into two intra-annual cohorts had a negligible effect on classification accuracies for age-0+ individuals of the snapper (*Pagrus auratus*) that settle in coastal nurseries of Victoria, Australia, between mid-December and March. Therefore, intra-annual variability may have little effect on classification accuracies of blacktip sharks in the Gulf given the short birth season.

The usefulness of natural biogeochemical tags derived from calcified structures stems from the relationship between water chemistry, other environmental factors and element : Ca ratios in the structures themselves. Under estuarine conditions, bony fish otolith Sr : Ca generally shows a positive correlation with salinity, whereas Ba : Ca typically shows a negative relationship with salinity (Limburg 1995; Elsdon and Gillanders 2005; Macdonald and Crook 2010). Because a large portion of otolith Sr and Ba is derived from the surrounding water (Walther and Thorrold 2006; Webb *et al.* 2012; Izzo *et al.* 2015), fluctuations in Sr : Ca and Ba : Ca can be used to infer the relative salinity of environments experienced by a fish during its lifetime (e.g. Limburg 1995). Similarly, Scharer *et al.* (2012) reported that the Sr : Ca in vertebral sections of the smalltooth sawfish (*Pristis pectinata*) had a positive correlation with salinity, and Smith *et al.* (2013) reported that the Ba : Ca in the vertebrae of round stingray (*Urolophus halleri*) was correlated with Ba in seawater. For juvenile blacktip sharks examined here, there were regional differences in Sr : Ca and Ba : Ca that were inversely related. However, it is difficult to attribute this pattern to regional differences in salinity without water chemistry data because the amount of Sr and Ba in freshwater end members will affect the amount of Sr and Ba available at a particular salinity (Wells *et al.* 2003; Kraus and Secor 2004). Moreover, several factors, including growth rate, ontogeny, temperature and diet, can affect the relationship between water chemistry and element : Ca deposition in calcified structures (for a review, see Sturrock *et al.* 2012). Growth rate does not appear to affect Li : Ca, Mg : Ca, Mn : Ca, Sr : Ca and Ba : Ca deposition in the round stingray; however, Mg : Ca, Mn : Ca and Ba : Ca were significantly affected by water temperature (Smith *et al.* 2013). The work of Smith *et al.* (2013) represents the only study to focus on the relationship between water chemistry and vertebral element : Ca in elasmobranchs; thus, we know little about the influences mentioned above and the extent of interspecies variability.

It is not necessary to fully elucidate the underlying abiotic and biotic factors driving regional variability in element : Ca ratios of calcified structures to use them as natural biogeochemical tags to examine population connectivity. However, other biologically relevant questions can be addressed if the primary factors influencing vertebral chemistry can be determined. Previously, regional differences in otolith Mn : Ca in gag grouper (*Mycteroperca microlepis*) and red snapper (*Lutjanus campechanus*) across the Gulf were attributed to latitudinal differences in soil chemistry (Hanson *et al.* 2004; Sluis *et al.* 2012). However, the majority of Mn in elasmobranch soft tissue appears to be derived from dietary sources (Mathews and Fisher

2009), which may explain observed enrichment in vertebral Mn : Ca relative to water Mn : Ca (Smith *et al.* 2013). An increase in Mn : Ca immediately following birth was present in the vertebrae of sharks from all regions in the present study, thus suggesting a common process influencing vertebral Mn : Ca. It is not clear to what degree the distribution of Mn is driven by the protein content of specific portions of vertebrae (Sturrock *et al.* 2012). Such transition metals do show an affinity for protein binding sites (Miller *et al.* 2006) and ~28% of otolith Mn can be bound to water soluble proteins (Izzo *et al.* 2016). However, there may also be a physiological process at play. The neonatal life stage is particularly stressful for elasmobranchs and can be associated with a decline in body mass (Duncan and Holland 2006) and high mortality (Heupel and Simpfendorfer 2002; Duncan and Holland 2006). Neonates rely on maternal nutrient reserves in the form of an enlarged liver to compensate for the lack of foraging experience (Hussey *et al.* 2010; Olin *et al.* 2011). Because the liver plays an important role in Mn homeostasis (Aschner and Aschner 2005; Madejczyk *et al.* 2009), metabolising liver tissue may result in excess Mn entering the bloodstream that becomes deposited in the vertebrae during the first weeks or months of life. Although Mn : Ca was the only elemental ratio that was significantly different among regions in both years in the present study, the possible link to food availability and maternal investment may result in intra-annual variability in vertebral Mn : Ca in a given area as less fit individuals perish. Thus, it is unknown whether vertebral Mn : Ca ratios of sharks collected early in the year would accurately reflect that of the surviving members of a cohort. Excluding Mn : Ca demonstrated that, although useful for discriminating groups, one can still achieve moderate to high classification accuracies using an array of other minor and trace elements.

Results from the present study suggest that examining connectivity between blacktip shark nurseries and offshore adult populations based on vertebral biogeochemical signatures holds some promise, but one factor that may compromise the effectiveness of such an approach is the unknown metabolic stability of vertebral tissue. Bone in higher vertebrates is constantly remodelled owing to its high healing potential (Kalfas 2001). The limited physiological response of elasmobranchs following natural (Officer *et al.* 1995) and experimentally induced (Ashhurst 2004) trauma suggests elasmobranchs may not be capable of extensive skeletal remodelling. However, unsuccessful attempts to age sharks based on radionuclide decay (^{226}Ra : ^{210}Pb) may indicate metabolic instability of the tissue, but another explanation would be exogenous uptake of ^{210}Pb during growth band formation (Welden *et al.* 1987; Fenton 2001). Furthermore, the success of bomb radiocarbon age validation (e.g. Campana *et al.* 2002; Passerotti *et al.* 2010, 2014), use of stable isotopes to detect ontogenetic shifts in diet (Estrada *et al.* 2006) and verification of annual bands using Sr : Ca patterns that correspond to known life history characteristics (Scharer *et al.* 2012) indicates the corpus calcareum likely remains relatively inert once formed.

Important questions remain surrounding the mechanisms governing mineral dynamics and sites of inclusion (e.g. substitution in calcium phosphate hydroxyapatite matrix, bound to proteins or incorporated interstitially), but the results of the

present study demonstrate that vertebral biogeochemical tags can be used to accurately distinguish nursery regions of blacktip sharks and may provide a means to estimate the relative contribution of different areas to the adult stock. By building an atlas of cohort-specific vertebral chemical signatures from different areas, it may be possible to infer the natal origin of adult sharks. This type of information would be particularly useful because some coastal sharks possess life history traits that make them vulnerable to overharvest. Whether protecting nursery areas should be the most important aspect of a management plan will depend on the biology of the species and life stage removed by the fishery (Kinney and Simpfendorfer 2009). However, estimates of connectivity between juvenile and adult populations can provide the insight necessary to evaluate the effectiveness of nursery conservation as a management tool.

Acknowledgements

Funding support for this project was provided by National Oceanic and Atmospheric Administration (NOAA) Fisheries, Panama City Laboratory, the Highly Migratory Species Division of NOAA Fisheries and the National Science Foundation (grant number DBI-1319188 to W. F. Patterson III). The authors thank D. M. Bethea and K. Smith (NOAA Fisheries), W. J. Bubley (Texas Parks and Wildlife Department), J. M. Drymon, M. N. Schrandt, A. M. Koretz and K. C. Gregalis (Dauphin Island Sea Lab) R. D. Grubbs, C. T. Peterson and J. L. Imhoff (Florida State University Coastal and Marine Laboratory), J. M. Hendon, J. M. Higgs and S. Ashworth (Gulf Coast Research Laboratory), R. J. D. Wells (Texas A&M University Galveston), N. M. Whitney and J. Morris (Mote Marine Laboratory) and the many staff members, students and volunteers who participated in field collection. The authors are also grateful for the guidance and advice provided by N. R. Miller (University of Texas Austin) during LA-ICP-MS analyses, and instrument time and technical support provide by C. J. P. O'Connor, J. Wilkins and the staff of the Scientific Division of Electro Scientific Industries (Bozeman, MT, USA).

References

- Aschner, J. L., and Aschner, M. (2005). Nutritional aspects of manganese homeostasis. *Molecular Aspects of Medicine* **26**, 353–362. doi:10.1016/J.MAM.2005.07.003
- Ashhurst, D. E. (2004). The cartilaginous skeleton of an elasmobranch fish does not heal. *Matrix Biology* **23**, 15–22. doi:10.1016/J.MATBIO.2004.02.001
- Baremore, I. E., and Passerotti, M. S. (2013). Reproduction of the blacktip shark in the Gulf of Mexico. *Marine and Coastal Fisheries* **5**, 127–138. doi:10.1080/19425120.2012.758204
- Beck, M. W., Heck, K. L., Jr, Able, K. W., Childers, D. L., Eggleston, D. B., Gillanders, B. M., Halpern, B., Hays, C. G., Hoshino, K., Minello, T. J., Orth, R. J., Sheridan, P. F., and Weinstein, M. P. (2001). The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *Bioscience* **51**, 633–641. doi:10.1641/0006-3568(2001)051[0633:TICAMO]2.0.CO;2
- Campana, S. E. (1999). Chemistry and composition of fish otoliths: pathways, mechanisms and applications. *Marine Ecology Progress Series* **188**, 263–297. doi:10.3354/MEPS188263
- Campana, S. E., Natanson, L. J., and Myklevoll, S. (2002). Bomb dating and age determination of large pelagic sharks. *Canadian Journal of Fisheries and Aquatic Sciences* **59**, 450–455. doi:10.1139/F02-027
- Carlson, J. K., Sulikowski, J. R., and Baremore, I. E. (2006). Do differences in life history traits exist for blacktip sharks, *Carcharhinus limbatus*, from the United States South Atlantic Bight and Eastern Gulf of Mexico? *Environmental Biology of Fishes* **77**, 279–292. doi:10.1007/S10641-006-9129-X
- Cook, G. S. (2011). Changes in otolith microchemistry over a protracted spawning season influence assignment of natal origin. *Marine Ecology Progress Series* **423**, 197–209. doi:10.3354/MEPS08942
- Duncan, K. M., and Holland, K. N. (2006). Habitat use, growth rates and dispersal patterns of juvenile scalloped hammerhead sharks *Sphyrna lewini* in a nursery habitat. *Marine Ecology Progress Series* **312**, 211–221. doi:10.3354/MEPS312211
- Elsdon, T. S., and Gillanders, B. M. (2005). Alternative life-history patterns of estuarine fish: barium in otoliths elucidates freshwater residency. *Canadian Journal of Fisheries and Aquatic Sciences* **62**, 1143–1152. doi:10.1139/F05-029
- Elsdon, T. S., Wells, B. K., Campana, S. E., Gillanders, B. M., Jones, C. M., Limburg, K. E., Secor, D. H., Thorrold, S. R., and Walther, B. D. (2008). Otolith chemistry to describe movements and life-history parameters of fishes: hypotheses, assumptions, limitations and inferences. *Oceanography and Marine Biology – an Annual Review* **46**, 297–330. doi:10.1201/9781420065756.CH7
- Estrada, J. A., Rice, A. N., Natanson, L. J., and Skomal, G. B. (2006). Use of isotopic analysis of vertebrae in reconstructing ontogenetic feeding ecology in white sharks. *Ecology* **87**, 829–834. doi:10.1890/0012-9658(2006)87[829:UOIAOV]2.0.CO;2
- Fenton, G. E. (2001). Radiometric Ageing of Sharks. FRDC Final Report 1994/021. Fisheries Research and Development Corporation, Canberra, Australia.
- Hamer, P. A., Jenkins, G. P., and Gillanders, B. M. (2003). Otolith chemistry of juvenile snapper *Pagrus auratus* in Victorian waters: natural chemical tags and their temporal variation. *Marine Ecology Progress Series* **263**, 261–273. doi:10.3354/MEPS263261
- Hanson, P. J., Koenig, C. C., and Zdanowicz, V. S. (2004). Elemental composition of otoliths used to trace estuarine habitats of juvenile gag *Mycteroperca microlepis* along the west coast of Florida. *Marine Ecology Progress Series* **267**, 253–265. doi:10.3354/MEPS267253
- Heupel, M. R., and Simpfendorfer, C. A. (2002). Estimation of mortality of juvenile blacktip sharks, *Carcharhinus limbatus*, within a nursery area using telemetry data. *Canadian Journal of Fisheries and Aquatic Sciences* **59**, 624–632. doi:10.1139/F02-036
- Heupel, M. R., Simpfendorfer, C. A., and Hueter, R. E. (2004). Estimation of shark home ranges using passive monitoring techniques. *Environmental Biology of Fishes* **71**, 135–142. doi:10.1023/B:EBFI.0000045710.18997.F7
- Heupel, M. R., Carlson, J. K., and Simpfendorfer, C. A. (2007). Shark nursery areas: concepts, definition, characterization and assumptions. *Marine Ecology Progress Series* **337**, 287–297. doi:10.3354/MEPS337287
- Hueter, R. E., and Tyminski, J. P. (2007). Species-specific distribution and habitat characteristics of shark nurseries in Gulf of Mexico waters off peninsular Florida and Texas. *American Fisheries Society Symposium* **50**, 193–223.
- Hueter, R. E., Heupel, M. R., Heist, E. J., and Keeney, D. B. (2005). Evidence of philopatry in sharks and implications for the management of shark fisheries. *Journal of Northwest Atlantic Fishery Science* **35**, 239–247. doi:10.2960/J.V35.M493
- Hussey, N. E., Wintner, S. P., Dudley, S. F., Cliff, G., Cocks, D. T., and Aaron MacNeil, M. (2010). Maternal investment and size-specific reproductive output in carcharhinid sharks. *Journal of Animal Ecology* **79**, 184–193. doi:10.1111/J.1365-2656.2009.01623.X
- Izzo, C., Doubleday, Z. A., Schultz, A. G., Woodcock, S. H., and Gillanders, B. M. (2015). Contribution of water chemistry and fish condition to otolith chemistry: comparisons across salinity environments. *Journal of Fish Biology* **86**, 1680–1698. doi:10.1111/JFB.12672
- Izzo, C., Doubleday, Z. A., and Gillanders, B. M. (2016). Where do elements bind within the otoliths of fish? *Marine and Freshwater Research* **67**, 1072–1076. doi:10.1071/MF15064
- Kalfas, I. H. (2001). Principles of bone healing. *Neurosurgical Focus* **10**, 1–4. doi:10.3171/FOC.2001.10.4.2

- Keeney, D. B., Heupel, M. R., Hueter, R. E., and Heist, E. J. (2005). Microsatellite and mitochondrial DNA analyses of the genetic structure of blacktip shark (*Carcharhinus limbatus*) nurseries in the northwestern Atlantic, Gulf of Mexico, and Caribbean Sea. *Molecular Ecology* **14**, 1911–1923. doi:10.1111/J.1365-294X.2005.02549.X
- Kinney, M. J., and Simpfendorfer, C. A. (2009). Reassessing the value of nursery areas to shark conservation and management. *Conservation Letters* **2**, 53–60. doi:10.1111/J.1755-263X.2008.00046.X
- Knip, D. M., Heupel, M. R., Simpfendorfer, C. A., Tobin, A. J., and Moloney, J. (2011). Wet-season effects on the distribution of juvenile pigeye sharks, *Carcharhinus amboinensis*, in tropical nearshore waters. *Marine and Freshwater Research* **62**, 658–667. doi:10.1071/MF10136
- Kohler, N. E., and Turner, P. A. (2001). Shark tagging: a review of conventional methods and studies. *Environmental Biology of Fishes* **60**, 191–224. doi:10.1023/A:1007679303082
- Kraus, R. T., and Secor, D. H. (2004). Incorporation of strontium into otoliths of an estuarine fish. *Journal of Experimental Marine Biology and Ecology* **302**, 85–106. doi:10.1016/J.JEMBE.2003.10.004
- Limburg, K. E. (1995). Otolith strontium traces environmental history of subyearling American shad *Alosa sapidissima*. *Marine Ecology Progress Series* **119**, 25–35. doi:10.3354/MEPS119025
- Macdonald, J. I., and Crook, D. A. (2010). Variability in Sr:Ca and Ba:Ca ratios in water and fish otoliths across an estuarine salinity gradient. *Marine Ecology Progress Series* **413**, 147–161. doi:10.3354/MEPS08703
- Madejczyk, M. S., Boyer, J. L., and Ballatori, N. (2009). Hepatic uptake and biliary excretion of manganese in the little skate, *Leucoraja erinacea*. *Comparative Biochemistry and Physiology Part C Toxicology & Pharmacology* **149**, 566–571. doi:10.1016/J.CBPC.2008.12.009
- Mathews, T., and Fisher, N. S. (2009). Dominance of dietary intake of metals in marine elasmobranch and teleost fish. *The Science of the Total Environment* **407**, 5156–5161. doi:10.1016/J.SCITOTENV.2009.06.003
- Miller, M. B., Clough, A. M., Batson, J. N., and Vachet, R. W. (2006). Transition metal binding to cod otolith proteins. *Journal of Experimental Marine Biology and Ecology* **329**, 135–143. doi:10.1016/J.JEMBE.2005.08.016
- Officer, R. A., Clement, J. G., and Rowler, D. K. (1995). Vertebral deformities in a school shark, *Galeorhinus galeus*: circumstantial evidence for endoskeletal resorption? *Journal of Fish Biology* **46**, 85–98. doi:10.1111/J.1095-8649.1995.TB05948.X
- Olin, J. A., Hussey, N. E., Fritts, M., Heupel, M. R., Simpfendorfer, C. A., Poulakis, G. R., and Frisk, A. T. (2011). Maternal meddling in neonatal sharks: implications for interpreting stable isotopes in young animals. *Rapid Communications in Mass Spectrometry* **25**, 1008–1016. doi:10.1002/RCM.4946
- Parsons, G. R., and Hoffmayer, E. R. (2007). Identification and characterization of shark nursery grounds along the Mississippi and Alabama gulf coasts. *American Fisheries Society Symposium* **50**, 301–316.
- Passerotti, M. S., Carlson, J. K., Piercy, A. N., and Campana, S. E. (2010). Age validation of great hammerhead shark (*Sphyrna mokarran*), determined by bomb radiocarbon analysis. *Fishery Bulletin* **108**, 346–351.
- Passerotti, M. S., Andrews, A. H., Carlson, J. K., Winter, S. P., Goldman, K. J., and Natanson, L. J. (2014). Maximum age and missing time in the vertebrae of sand tiger shark (*Carcharias taurus*): validated lifespan from bomb radiocarbon dating in the western North Atlantic and southwestern Indian Oceans. *Marine and Freshwater Research* **65**, 674–687. doi:10.1071/MF13214
- Pitts, L. C., and Wallace, G. T. (1994). Lead deposition in the shell of the bivalve, *Mya arenaria*: an indicator of dissolved lead in seawater. *Estuarine, Coastal and Shelf Science* **39**, 93–104. doi:10.1006/ECSS.1994.1051
- Scharer, R. M., Patterson, W. F., III, Carlson, J. K., and Poulakis, G. R. (2012). Age and growth of endangered smalltooth sawfish (*Pristis pectinata*) verified with LA-ICP-MS analysis of vertebrae. *PLoS One* **7**, e47850. doi:10.1371/JOURNAL.PONE.0047850
- Sluis, M. Z., Barnett, B. K., Patterson, W. F., III, Cowan, J. H., Jr, and Shiller, A. M. (2012). Discrimination of juvenile red snapper otolith chemical signatures from Gulf of Mexico nursery regions. *Marine and Coastal Fisheries* **4**, 587–598. doi:10.1080/19425120.2012.703163
- Smith, W. D. (2013). Vertebral elemental markers in elasmobranchs: potential for reconstructing environmental history and population structure. Ph.D. Thesis, Oregon State University, Corvallis.
- Smith, W. D., Miller, J. A., and Heppell, S. S. (2013). Elemental markers in elasmobranchs: effects of environmental history and growth on vertebral chemistry. *PLoS One* **8**, e62423. doi:10.1371/JOURNAL.PONE.0062423
- Southeast Data, Assessment, and Review (2012). Stock Assessment Report of SEDAR 29, HMS Gulf of Mexico Blacktip Shark. (Southeast Fisheries Science Center: Charleston, SC.) Available at http://sedarweb.org/docs/sar/S29_GOM%20blacktip%20report_SAR_final.pdf [Verified 8 November 2015].
- Steiner, P. A., Michel, M., and O'Donnell, P. M. (2007). Notes on the occurrence and distribution of elasmobranchs in the Ten Thousand Islands Estuary, Florida. *American Fisheries Society Symposium* **50**, 237–250.
- Sturrock, A. M., Trueman, C. N., Darnaude, A. M., and Hunter, E. (2012). Can otolith elemental chemistry retrospectively track migrations in fully marine fishes? *Journal of Fish Biology* **81**, 766–795. doi:10.1111/J.1095-8649.2012.03372.X
- Tillett, B. J., Meekan, M. G., Parry, D., Munksgaard, N., Field, I. C., Thorburn, D., and Bradshaw, C. J. (2011). Decoding fingerprints: elemental composition of vertebrae correlates to age-related habitat use in two morphologically similar sharks. *Marine Ecology Progress Series* **434**, 133–142. doi:10.3354/MEPS09222
- Vasconcelos, R. P., Reis-Santos, P., Costa, M. J., and Cabral, H. N. (2011). Connectivity between estuaries and marine environment: Integrating metrics to assess estuarine nursery function. *Ecological Indicators* **11**, 1123–1133. doi:10.1016/J.ECOLIND.2010.12.012
- Walther, B. D., and Thorrold, S. R. (2006). Water, not food, contributes the majority of strontium and barium deposited in the otoliths of a marine fish. *Marine Ecology Progress Series* **311**, 125–130. doi:10.3354/MEPS311125
- Webb, S. D., Woodcock, S. H., and Gillanders, B. M. (2012). Sources of otolith barium and strontium in estuarine fish and the influence of salinity and temperature. *Marine Ecology Progress Series* **453**, 189–199. doi:10.3354/MEPS09653
- Welden, B. A., Cailliet, G. M., and Flegal, A. R. (1987). Comparison of radiometric with vertebral band age estimates in four California elasmobranchs. In 'Age and Growth of fish'. (Eds R. C. Summerfelt and G. E. Hall.) pp. 301–315. (Iowa State University Press: Ames, IA.)
- Wells, B. K., Bath, G. E., Thorrold, S. R., and Jones, C. M. (2000). Incorporation of strontium, cadmium, and barium in juvenile spot (*Leiostomus xanthurus*) scales reflects water chemistry. *Canadian Journal of Fisheries and Aquatic Sciences* **57**, 2122–2129. doi:10.1139/F00-178
- Wells, B. K., Rieman, B. E., Clayton, J. L., Horan, D. L., and Jones, C. M. (2003). Relationships between water, otolith, and scale chemistries of westslope cutthroat trout from the Coeur d'Alene River, Idaho: the potential application of hard-part chemistry to describe movements in freshwater. *Transactions of the American Fisheries Society* **132**, 409–424. doi:10.1577/1548-8659(2003)132<0409:RBWOAS>2.0.CO;2
- Yeiser, B. G., Heupel, M. R., and Simpfendorfer, C. A. (2008). Occurrence, home range and movement patterns of juvenile bull (*Carcharhinus leucas*) and lemon (*Negaprion brevirostris*) sharks within a Florida estuary. *Marine and Freshwater Research* **59**, 489–501. doi:10.1071/MF07181