

Supplementary Material

Oestrogen-induced expression of decay accelerating factor is spatiotemporally antagonised by progesterone–progesterone receptor signalling in mouse uterus

Mi Ji Lee^A, Hyang Ah Lee^B, Mira Park^A, Hee Kyoung Park^A, Yeon Sun Kim^A, Seung Chel Yang^A, Hye-Ryun Kim^A, Jayeon Kim^{C,D} and Haengseok Song^{A,D}

^ADepartment of Biomedical Science, CHA University, Seongnam-si, Gyeonggi-do, 13488, Republic of Korea.

^BDepartment of Obstetrics and Gynaecology, Kangwon National University, Chuncheon-si, Gangwon-do, 24341, Republic of Korea.

^CSeoul CHA Fertility Centre, CHA University, Seoul, 04637, Republic of Korea.

^DCorresponding authors. Emails: hssong@cha.ac.kr; jayeon_kim@cha.ac.kr

Table S1. Gene-specific primer sequences used for RT-PCR and/or real-time RT-PCR

| Primer | | Sequence | Size (bp) |
|---------------|---------|--------------------------------|-----------|
| <i>DAF</i> | Forward | 5'- ACCTCCACTCCCAGGAAAAG -3' | 197 |
| | Reverse | 5'- TAGAGGAGACACCGACTAGCC -3' | |
| <i>Egr1</i> | Forward | 5'- ACCCTATGAGCACCTGACCAC -3' | 116 |
| | Reverse | 5'- TATAGGTGATGGGAGGCAACC -3' | |
| <i>LF</i> | Forward | 5'- CCGCTCAGTTGTGTCAAGAAA -3' | 276 |
| | Reverse | 5'- CAGGTCGCAGTTTGTAGGGG -3' | |
| <i>Hoxa10</i> | Forward | 5'- CCCCTTGGCAGTGATTCTG -3' | 189 |
| | Reverse | 5'- CTACGGCTGATCTCTAGGCG -3' | |
| <i>Cox1</i> | Forward | 5'- ATGAGTCGAAGGAGTCTCTCG -3' | 129 |
| | Reverse | 5'- GCACGGATAGTAACAACAGGGA -3' | |
| <i>rPL7</i> | Forward | 5'- TCAATGGAGTAAGCCCAAAG -3' | 246 |
| | Reverse | 5'- CAAGAGACCGAGCAATCAAG -3' | |