

[10.1071/RD22123](https://doi.org/10.1071/RD22123)

Reproduction, Fertility and Development

Supplementary Material

Redox profile and mediators of the unfolded protein response (UPR) in the placenta of rats during pregnancy

Jeane Martinha dos Anjos Cordeiro^A, Luciano Cardoso Santos^A, Bianca Reis Santos^A, Emilly Oliveira Santos^A, Acácia Eduarda de Jesus Nascimento^A, Gustavo José Cota de Freitas^B, Junya de Lacorte Singulani^B, Daniel de Assis Santos^B, Mário Sérgio Lima de Lavor^C, and Juneo Freitas Silva^{A,}*

^ADepartamento de Ciencias Biologicas, Centro de Microscopia Eletronica, Universidade Estadual de Santa Cruz, Ilheus, Brazil.

^BDepartamento de Microbiologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

^CDepartamento de Ciencias Agrarias e Ambientais, Hospital Veterinario, Universidade Estadual de Santa Cruz, Ilheus, Brazil.

*Correspondence to: Juneo Freitas Silva Departamento de Ciencias Biologicas, Centro de Microscopia Eletronica, Universidade Estadual de Santa Cruz, Campus Soane Nazare de Andrade, Rodovia Jorge Amado, km 16, Salobrinho, Ilheus, Bahia 45662-900, Brazil Email: jfsilva@uesc.br

Supplementary Figure S1. (A) Photomicrographs of HIF1 α immunostaining in the placental disk showing random fields from the junctional zone (JZ) and placental labyrinth (PL) chosen for immunostaining evaluation. Images of five random fields in each region were obtained to quantitative evaluation of the immunostaining area. B-C) Photomicrographs of the negative control showing the JZ (B) and PL (C) (streptavidin-biotin-peroxidase, counterstained with Harris hematoxylin, Bar= 50 μ m). MT, metrial triangle; DB, decidua basalis.

