## Supplementary material

## Prescribed fire as a tool to regenerate live and dead serotinous jack pine (Pinus banksiana) stands

Maria Sharpe<sup>A,B,C</sup>, Hyejin Hwang<sup>A</sup>, David Schroeder<sup>B</sup>, Soung Ryoul Ryu<sup>A</sup> and Victor J. Lieffers<sup>A</sup>

<sup>A</sup>Department of Renewable Resources, University of Alberta, Edmonton, AB, T6G 2H1, Canada.

<sup>B</sup>Ministry of Agriculture and Forestry, Government of Alberta, Edmonton, AB, T5K 2M4, Canada.

<sup>C</sup>Corresponding author. Email: maria.sharpe@gov.ab.ca



Fig. S1. Mean seedling density in relation surface, intermittent crown and continuous crown fires. Assessments made three years after fire in stands that were live or dead prior to burning (n = 3, error bars are s.e.).



**Fig. S2.** Mean depth of residual duff in relation to types of fire (surface, intermittent crown and continuous crown). Assessments made three years after fire in stands that were living or dead prior to burning (n = 3, error bars are s.e.).



**Fig. S3.** Mean seedling stocking in relation to fire type (surface, intermittent crown and continuous crown). Assessment made three years after fire in stands that were living or dead prior to burning (n = 3).