

[10.1071/PC24016](https://doi.org/10.1071/PC24016)

Pacific Conservation Biology

Supplementary Material

Annual response patterns in activity and demographic parameters of reptile species occupying a predictable Mediterranean climate

Richard A. How^{A,B,*}

^ASchool of Human Sciences, The University of Western Australia, Perth, WA 6009, Australia.

^BCollections and Research, Western Australian Museum, Welshpool DC, WA 6986, Australia.

*Correspondence to: Richard A. How School of Human Sciences, The University of Western Australia, Perth, WA, Australia Emails: ric.how@uwa.edu.au; richow1944@gmail.com

Supplementary Material

Body mass and SVL for adult females and males in different reproductive states for 18 species recorded regularly during Regime III of the study.

SPECIES	Mass [gm ± SD(n)]				SVL [mm ± SD(n)]			
	Females		Males		Females		Males	
	Gravid	Other Adults	Seminal Plugs	Other Adults	Gravid	Other Adults	Seminal Plugs	Other Adults
<i>C. fallens</i>	15.7±2.2(95)	12.6±1.9(276)	12.6±1.8(89)	12.5±2.0(488)	90.6±3.5(95)	88.4±2.9(273)	85.3±4.0(89)	84.8±4.1(488)
<i>S. spinigerus</i>	6.6±0.9(245)	5.3±1.0(1646)		4.0±0.6(416)	68.4±3.1(244)	65.5±3.2(1620)		60.9±2.5(416)
<i>H. quadrilineatus</i>	1.61±0.42(195)	1.22±0.28(1092)	1.52±0.11(4)	1.32±0.30(540)	52.6±5.8(195)	49.0±3.8(1093)	48.8±1.5(4)	48.5±2.8(543)
<i>L. elegans</i>	0.71±0.13(98)	0.56±0.11(415)	0.55±0.08(113)	0.53±0.08(544)	40.3±1.8(98)	38.8±1.9(418)	37.2±1.5(114)	36.9±1.6(545)
<i>L. lineopunctulata</i>	6.1±1.3(148)	4.4±1.3 (308)	4.3±0.8(278)	4.1±0.8(349)	93.9±6.6(148)	86.4±8.1(308)	85.0±4.9(278)	83.3±5.3(351)
<i>S. bertholdi</i>	11.3±2.3(24)	8.8±2.1(139)	6.4±1.0(40)	6.4±1.2(253)	238.0±16.0(24)	226.5±16.6(139)	199.3±11.3(40)	197.6±11.0(253)
<i>M. greyii</i>	0.42±0.09(145)	0.32±0.06(216)		0.36±0.05(2)	30.8±2.6(145)	29.1±2.6(222)		29.5±0.5(2)
<i>A. australis</i>	7.6±4.5(51)	4.5±2.5(140)	2.5±0.6(3)	4.1±1.3(154)	237.0±46.5(51)	206.2±33.6(140)	164.7±7.1(3)	197.3±21.8(155)
<i>P. minor</i>	34.2±7.4(33)	26.6±8.8(110)	27.6±3.7(4)	26.6±6.7(61)	106.0±7.5(29)	99.6±9.8(109)	101.0±3.1(4)	98.4±7.3(60)
<i>C. buechananii</i>	1.18±0.22(48)	0.98±0.24(59)	0.94±0.07(7)	0.91±0.17(43)	42.4±1.9(49)	41.6±2.7(60)	39.9±0.7(7)	40.1±2.7(45)
<i>L. burtonis</i>	15.9±3.6(60)	10.8±3.9(52)	8.5±2.3(23)	7.4±1.4(28)	187.1±18.5(60)	168.9±23.7(52)	146.4±13.4(23)	140.5±9.1(28)
<i>L. praepedita</i>	0.75±0.11(23)	0.58±0.10(56)	0.56±0.07(55)	0.56±0.08(80)	61.0±2.2(23)	57.6±2.5(56)	56.8±2.9(55)	56.2±1.9(80)
<i>B. semifasciatus</i>	15.1±3.6(23)	13.7±2.5(24)	11.3±1.7(11)	10.7±2.3(61)	287.7±14.1(23)	290.1±22.4(24)	262.7±17.2(11)	260.7±19.5(61)
<i>T. rugosa</i>		456.3±56.7(24)	462.2±49.4(23)	449.8±68.5(79)		284.0±9.2(28)	286.6±9.5(23)	285.1±9.49(78)
<i>A. repens</i>	0.83±0.20(9)	0.65±0.10(47)	0.60±0.09(23)	0.55±0.09(19)	97.7±6.0(9)	90.6±5.4(47)	86.6±4.9(23)	87.0±4.9(19)
<i>N. bimaculatus</i>	10.6±5.2(11)	7.2±4.3(46)	6.5±2.3(8)	5.1±1.7(38)	332.5±48.0(11)	306.9±50.3(46)	296.6±26.6(8)	273.3±26.4(38)
<i>C. chapmani</i>	4.05(1)	4.14±0.63(12)	2.86(1)	2.89±0.47(22)	46.0±0(1)	49.5±2.5(11)	42.0±0(1)	43.8±1.7(19)
<i>C. australis</i>	16.3(1)	13.3±1.9(9)	15.6±0.7(2)	15.6±1.5(16)	96.5(1)	93.2±2.0(9)	93.5±0.5(2)	93.2±2.5(16)