Supplementary Material

Grazing management of Australian native woody regeneration as an effective nature-based climate-change solution

C. M. Waters^{A,B,*}, R. B. Hacker^C, A. Sekaran^B and A. R. Grant^D

^AGreenCollar, 3 Hickson Road, The Rocks Sydney, NSW 2000, Australia

^BHawkesbury Institute for the Environment, Western Sydney University, Penrith, NSW 2751, Australia

^CPresent address: Ron Hacker Rangeland Consulting Services, 29 Edward Street, Tenambit, NSW, Australia

^DPresent address: PO Box 1591, Young, NSW 2594, Australia

*Correspondence to: Email: cathleen.waters@greencollargroup.com.au

Supplementary Fig. S1. The relative proportion of feed in diets of major domestic livestock for a poplar box-mulga woodlands of western NSW. *Adapted from Pahl 2019 (Table 2)*.

								Relative proportion in diet					
										Small () Mediur	m 🔘	Large
		Wet conditions						Dry conditions					
		Annual grasses	Ephemeral forbs	Perennial grasses	Chenopod perennial forbs	Non- chenopod perennial forbs	Browse from large shrubs and trees	Annual grasses	Ephemeral forbs	Perennial grasses	Chenopod perennial forbs	Non- chenopod perennial forbs	Browse from large shrubs and trees
Sheep	Autumn					•	•				•		•
	Winter				•	•					•		•
	Spring	•				•					•		•
	Summer	•		•	•	•					•		•
Cattle	Autumn		•				•						
	Winter	•	•								•	•	•
	Spring		•								•	•	
	Summer		•				•			•			
Goat	Autumn									•			
	Winter			•							•		
	Spring						•					•	
	Summer			•									