

[10.1071/WR23149](https://doi.org/10.1071/WR23149)

Wildlife Research

Supplementary Material


Identifying black rat (*Rattus rattus*) movement patterns aids the development of management programs in citrus orchards

Roger A. Baldwin^{A,}, Ryan Meinerz^A, and Justine A. Smith^A*


^ADepartment of Wildlife, Fish, and Conservation Biology, University of California, Davis, CA 95616, USA.

*Correspondence to: Roger A. Baldwin Department of Wildlife, Fish, and Conservation Biology, One Shields Avenue, University of California, Davis, CA 95616, USA Email: rabaldwin@ucdavis.edu

Site 1



Site 2



Supplementary Fig. S1. The square outline of our study site as represented by the outer row of detection nodes, as well as individual black rat home ranges found within our two study sites in the southern San Joaquin Valley, California.

Supplementary Table S1. The weight and sex of captured black rats from two citrus orchards in the southern San Joaquin Valley, California, USA, during summer 2020.

Site	Sex	Weight (g)
1	Male	138
1	Female	132
1	Male	212
1	Male	128
1	Male	222
1	Male	244
1	Female	104
1	Female	155
1	Female	139
1	Female	190
1	Male	184
1	Male	128
1	Male	199
1	Male	139
1	Male	134
1	Male	227
2	Male	168
2	Female	133
2	Male	275
2	Female	188
2	Female	163
2	Male	199
2	Female	216
2	Male	204
2	Male	221
2	Male	289
2	Female	171
2	Female	193
2	Male	224