## 10.1071/BT23110

Australian Journal of Botany

## Supplementary Material

## Pollination by multiple species of nectar foraging Hymenoptera in *Prasophyllum innubum*, a critically endangered orchid of the Australian Alps

Tobias Hayashi<sup>A,B,C</sup>, Bronwyn M. Ayre<sup>A,B</sup>, Björn Bohman<sup>D,E</sup>, Graham R. Brown<sup>F,G,H</sup>, Noushka Reiter<sup>A,B,C</sup>, and Ryan D. Phillips<sup>A,B,\*</sup>

<sup>A</sup>Department of Environment and Genetics and the Research Centre for Future Landscapes, La Trobe University, Melbourne, Vic, Australia.

<sup>B</sup>Royal Botanic Gardens Victoria, Science Division, Corner of Ballarto Road and Botanic Drive,

Cranbourne, Vic 3977, Australia.

<sup>c</sup>Ecology and Evolution, Research School of Biology, The Australian National University, Canberra, ACT 2600, Australia.

<sup>D</sup>School of Molecular Sciences, The University of Western Australia, Crawley, WA, Australia.

<sup>E</sup>Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Lomma, Sweden.

<sup>F</sup>Natural Sciences, Museum and Art Gallery of the Northern Territory, GPO Box 4646, Darwin, NT0801, Australia.

<sup>G</sup>CSIRO Land and Water, Private Mail Bag 44, Winnellie, NT0822, Australia.

<sup>H</sup>Research Institute for Environment and Livelihoods, Charles Darwin University, Darwin, NT0909, Australia.

\*Correspondence to: Ryan D. Phillips Department of Environment and Genetics and the Research Centre for Future Landscapes, La Trobe University, Melbourne, Vic, Australia Email: R.Phillips@latrobe.edu.au

**Supplementary Table S1**: Number and proportion of co-flowering plants present at MOP from three 120 x 2 m transects (10 meters apart) through the middle of the main *Prasophyllum innumbum* population. The count was conducted on the 8/2/2021 as an indication of which plant species were most floriferous during peak flowering of *P. innubum*. *Prasophyllum innubum* is listed first, with the remaining species ranked based on the proportion of flowers they represent in the community at the time of measurement. Note, for members of the Asteraceae, a capitulum was scored as a flower rather than individual florets.

Species	Transect 1		Transect 2		Transect 3		Overall		Proportion	
	Plants	Flowers	Plants	Flowers	Plants	Flowers	Plants	Flowers	Plants	Flowers
Prasophyllum										
innubum	25	186	7	35	10	106	42	327	0.043	0.182
Leptorhynchos										
squamatus	435	435	126	126	115	115	676	676	0.686	0.377
Spiranthes										
australis*	0	0	18	270	0	0	18	270	0.018	0.151
Stylidium										
montanum	15	93	10	44	10	58	35	195	0.035	0.109
Veronica										
subtilis	53	53	12	52	45	45	110	150	0.112	0.084
Microtis rara	0	0	0	0	7	33	7	33	0.007	0.018
Leucochrysum										
albicans	2	4	6	6	22	22	30	32	0.030	0.018
Xerochrysum										
subundulatum	0	0	0	0	32	32	32	32	0.032	0.018
Leptospermum										
sp.	1	22	0	0	0	0	1	22	0.001	0.012
Hypochaeris										
radicata	5	6	4	4	3	5	12	15	0.012	0.008
Prasophyllum										
sp. <i>aff</i>										
sphacelatum	0	0	0	0	1	12	1	12	0.001	0.007
<i>Epilobium</i> sp.	0	0	4	6	1	1	5	7	0.005	0.004
Brachyscome										
scapigera	1	1	4	5	0	0	5	6	0.005	0.003

Arthropodium										
milleflorum	1	1	2	3	0	0	3	4	0.003	0.002
Bulbine										
bulbosa	2	3	0	0	0	0	2	3	0.002	0.002
Trifolium										
repens	0	0	0	0	3	3	3	3	0.003	0.002
Ranunculus										
graniticola	0	0	1	3	0	0	1	3	0.001	0.002
Wahlenbergia										
sp.	0	0	1	2	0	0	1	2	0.001	0.001
Arthropodium										
milleflorum	2	2	0	0	0	0	2	2	0.002	0.001

\*Note: Number of Spiranthes flowers an estimate, based off an average of 15 flowers per

inflorescence of ten plants.