

[10.1071/BT24042](https://doi.org/10.1071/BT24042)

*Australian Journal of Botany*

### **Supplementary Material**

#### **Half a century of change in floristics and structure in an urban grassy woodland: implications for conservation management**

*Jamie B. Kirkpatrick<sup>A</sup>, Ian Jenkins<sup>A</sup>, and Kerry L. Bridle<sup>A,\*</sup>*

<sup>A</sup> School of Geography, Planning and Spatial Sciences, University of Tasmania, Private Bag 78, Hobart, Tas 7001, Australia.

\*Correspondence to: Kerry L. Bridle School of Geography, Planning and Spatial Sciences, University of Tasmania, Private Bag 78, Hobart, Tas 7001, Australia Email: [Kerry.Bridle@utas.edu.au](mailto:Kerry.Bridle@utas.edu.au)

Table S1. Constants and coefficients for year in the linear models for richness (rich.) variables. ns = P > 0.05, \* = P =/ < 0.05, \*\* = P = < 0.01, \*\*\* = P < 0.001

Variable	Constant	1974	1984	1994	2001
Species richness	47.23***	-3.16***	2.38***	0.91ns	0.62ns
Native species rich.	30.21***	-2.03***	1.34**	0.61ns	0.11ns
Native richness (%)	65.04***	-	-	-	-
Non-native rich.	14.44***	-1.04***	1.13***	0.22ns	-0.19ns
Annual herb richness	1.32***	-0.67***	0.42***	-0.16ns	-0.03ns
Perennial herb rich.	3.42***	0.27ns	1.48***	-0.39ns	-0.47*
Annual grass richness	7.53***	-	-	-	-
Perennial grass rich.	4.41***	-0.92***	0.36*	0.46**	0.26ns
Sedge and rush rich.	2.31***	-0.71***	0.26ns	0.37**	0.39**
Shrub richness	20.39***	-	-	-	-
Tree richness	1.02***	-0.80***	-0.66***	0.19ns	0.19ns
Geophyte richness	-0.37*	-0.06ns	0.31***	0.08ns	0.06ns

**Table S2. Percentage frequency of taxa found at least 5% of plots surveyed from 1974 – 2022, showing the significance of the variation between years.**

Taxa are ordered from greatest to smallest change in frequent from 1974 to 2022 within each life form class. '#' indicates change from 2022/1984 instead of 2022/1974 as taxon was not recorded in 1974. A = non-native taxon, B = both non-native and native species in taxon. Bold values are the highest frequencies for a taxon. Underlined values exhibit a continuous increase or decrease across the years. Significance of change in time was determined by Chi-squared analysis with \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ , ns = not significant.

Taxon	1974	1984	1994	2000	2022	Significance of change	Average frequency	Change 2022/1974
<i>Annual herbs</i>								
<sup>A</sup> <i>Cirsium vulgare</i>	0.0	5.8	2.9	1.4	<b>21.7</b>	***	6.4	3.8#
<sup>A</sup> <i>Petrorhagia nanteuilii</i>	2.9	26.1	26.1	<b>47.8</b>	29.0	***	26.4	10
<sup>A</sup> <i>Sonchus</i> spp.	5.8	<b>37.7</b>	4.3	4.3	18.8	***	14.2	3.3
<sup>A</sup> <i>Centaurium erythraea</i>	7.2	18.8	33.3	17.4	<b>46.4</b>	***	24.6	6.4
<sup>A</sup> <i>Vicia</i> spp.	11.6	<b>17.4</b>	13.0	4.3	10.1	ns	11.3	0.9
<i>Senecio</i> spp.	37.7	53.6	33.3	42.0	<b>58.0</b>	*	44.9	1.5
<i>Perennial herbs</i>								
<sup>A</sup> <i>Lysimachia arvensis</i>	2.9	<b>13.0</b>	1.4	1.4	11.6	**	6.1	4
<sup>A</sup> <i>Medicago</i> spp.	2.9	<b>21.7</b>	2.9	0.0	0.0	***	5.5	0
<sup>A</sup> <i>Trifolium</i> spp.	4.3	2.9	13.0	<b>14.5</b>	11.6	ns	9.3	2.7
<i>Leptorhynchos nitidulus</i>	7.2	10.1	4.3	<b>17.4</b>	13.0	ns	10.4	1.8
<sup>B</sup> <i>Linum</i> spp.	13.0	<b>36.2</b>	17.4	11.6	23.2	**	20.3	1.8
<i>Ranunculus lappaceus</i>	<b>11.6</b>	10.1	2.9	10.1	8.7	ns	8.7	0.8
<i>Oxalis perennans</i>	44.9	40.6	17.4	14.5	<b>47.8</b>	***	33.0	1.1
<i>Wahlenbergia</i> spp.	11.6	<b>15.9</b>	4.3	11.6	11.6	ns	11.0	1
<sup>A</sup> <i>Hypochaeris radicata</i>	27.5	<b>60.9</b>	31.9	15.9	23.2	***	31.9	0.8
<i>Hackelia suaveolens</i>	0.0	17.4	13.0	11.6	<b>20.3</b>	***	12.5	1.2#
<i>Chrysocephalum apiculatum</i>	7.2	11.6	<b>20.3</b>	11.6	2.9	*	10.7	0.4
<sup>A</sup> <i>Urospermum dalechampii</i>	53.6	<b>60.9</b>	59.4	53.6	47.8	ns	55.1	0.9
<i>Gonocarpus tetragynus</i>	18.8	18.8	14.5	<b>23.2</b>	11.6	ns	17.4	0.6
<sup>A</sup> <i>Sanguisorba minor</i>	11.6	<b>23.2</b>	20.3	13.0	4.3	*	14.5	0.4
<sup>A</sup> <i>Plantago lanceolata</i>	78.3	79.7	<b>87.0</b>	85.5	63.8	**	78.8	0.8
<i>Plantago varia</i>	<b>58.0</b>	33.3	20.3	30.4	27.5	***	33.9	0.5
<i>Goodenia lanata</i>	<b>11.6</b>	<b>11.6</b>	4.3	2.9	4.3	ns	7.0	0.4
<i>Geranium potentilloides</i>	24.6	<b>33.3</b>	14.5	15.9	10.1	**	19.7	0.4
<i>Acaena echinata</i>	<b>73.9</b>	<b>63.8</b>	<b>59.4</b>	<b>58.0</b>	<b>40.6</b>	**	59.1	0.5

<i>Leptorhynchos squamatus</i>	<b>29.0</b>	23.2	23.2	18.8	5.8	*	20.0	0.2
<i>Annual grasses</i>								
<sup>A</sup> <i>Aira</i> spp.	31.9	<b>55.1</b>	43.5	53.6	52.2	*	47.2	1.6
<sup>A</sup> <i>Briza</i> spp.	30.4	<b>34.8</b>	14.5	29.0	33.3	ns	28.4	1.1
<sup>A</sup> <i>Vulpia</i> spp.	14.5	<b>20.3</b>	18.8	11.6	15.9	ns	16.2	1.1
<i>Perennial grasses</i>								
<i>Microlaena stipoides</i>	1.4	27.5	5.8	18.8	<b>39.1</b>	***	18.6	27
<sup>A</sup> <i>Dactylis glomerata</i>	13.0	14.5	<b>31.9</b>	23.2	27.5	*	22.0	2.1
<sup>A</sup> <i>Holcus lanatus</i>	<b>8.7</b>	2.9	5.8	5.8	<b>8.7</b>	ns	6.4	1
<i>Anthosachne scabra</i>	53.6	<b>89.9</b>	82.6	85.5	71.0	***	76.5	1.3
<i>Austrostipa</i> spp.	73.9	94.2	94.2	<b>95.7</b>	91.3	***	89.9	1.2
<i>Deyeuxia quadriseta</i>	0.0	21.7	<b>31.9</b>	17.4	18.8	***	18.0	0.9 <sup>#</sup>
<i>Poa rodwayi</i>	88.4	85.5	85.5	89.9	<b>94.2</b>	ns	88.7	1.1
<i>Rytidosperma</i> spp.	94.2	84.1	94.2	<b>98.6</b>	81.2	**	90.4	0.9
<i>Themeda triandra</i>	91.3	94.2	92.8	<b>95.7</b>	78.3	**	90.4	0.9
<i>Dichelachne</i> spp.	<b>76.8</b>	71.0	75.4	63.8	63.8	***	70.1	0.8
<i>Lachnagrostis aemula</i>	58.0	62.3	<b>71.0</b>	42.0	33.3	***	53.3	0.6
<i>Pentapogon quadrifidus</i>	14.5	<b>21.7</b>	10.1	10.1	1.4	**	11.6	0.1
<sup>A</sup> <i>Lolium perenne</i>	<b>17.4</b>	<u>10.1</u>	<u>10.1</u>	<u>7.2</u>	<u>4.3</u>	ns	9.9	0.3
<i>Sedges and rushes</i>								
<i>Schoenus apogon</i>	1.4	<b>47.8</b>	44.9	31.9	29.0	***	31.0	20
<i>Lomandra longifolia</i>	18.8	24.6	30.4	27.5	<b>31.9</b>	ns	26.7	1.7
<i>Dianella</i> spp.	23.2	34.8	44.9	<b>52.2</b>	33.3	**	37.7	1.4
<i>Lepidosperma laterale</i>	21.7	26.1	34.8	<b>37.7</b>	24.6	ns	29.0	1.1
<i>Lepidosperma gunnii</i>	24.6	29.0	<b>44.9</b>	34.8	24.6	*	31.6	1
<i>Carex breviculmis</i>	68.1	71.0	62.3	<b>76.8</b>	33.3	***	62.3	0.5
<i>Shrubs</i>								
<i>Styphelia humifusa</i>	10.1	14.5	23.2	20.3	<b>33.3</b>	***	20.3	3.3
<i>Olearia ramulosa</i>	8.7	14.5	<b>23.2</b>	<b>23.2</b>	17.4	ns	17.4	2
<i>Lissanthe strigosa</i>	8.7	14.5	<b>18.8</b>	15.9	15.9	ns	14.8	1.8
<i>Bossiaea prostrata</i>	53.6	46.4	59.4	<b>72.5</b>	52.2	*	56.8	1
<sup>A</sup> <i>Rosa rubiginosa</i>	10.1	13.0	<b>15.9</b>	<b>15.9</b>	7.2	ns	12.5	0.7
<sup>A</sup> <i>Acacia pycnantha</i>	5.8	8.7	8.7	<b>10.1</b>	1.4	ns	7.0	0.3
<i>Pimelea humilis</i>	<b>44.9</b>	<b>40.6</b>	<b>39.1</b>	<b>34.8</b>	<b>34.8</b>	ns	38.8	0.8
<i>Hibbertia hirsuta</i>	<b>11.6</b>	<u>10.1</u>	<u>7.2</u>	<u>7.2</u>	<u>7.2</u>	ns	8.7	0.6

<sup>A</sup> <i>Chrysanthemoides monilifera</i>	<b>15.9</b>	13.0	7.2	4.3	2.9	*	8.7	0.2
<b>Trees</b>								
<i>Eucalyptus pulchella</i>	2.9	1.4	7.2	5.8	<b>14.5</b>	*	6.4	5
<i>Allocasuarina verticillata</i>	<b>8.7</b>	<b>26.1</b>	<b>47.8</b>	<b>47.8</b>	<b>72.5</b>	***	40.6	8.3
<i>Eucalyptus globulus</i>	<b>1.4</b>	<b>4.3</b>	<b>7.2</b>	<b>7.2</b>	<b>10.1</b>	ns	6.1	7
<i>Acacia mearnsii</i>	11.6	8.7	14.5	11.6	<b>39.1</b>	***	17.1	3.4
<i>Eucalyptus viminalis</i>	14.5	13.0	39.1	30.4	<b>39.1</b>	***	27.2	2.7
<i>Dodonaea viscosa</i>	7.2	5.8	5.8	7.2	<b>17.4</b>	ns	8.7	2.4
<i>Bursaria spinosa</i>	37.7	31.9	47.8	<b>60.9</b>	50.7	**	45.8	1.3
<b>Geophytes</b>								
<i>Thelymitra</i> spp.	1.4	<b>17.4</b>	15.9	11.6	2.9	**	9.9	2
<i>Arthropodium milleflorum</i>	30.4	24.6	30.4	<b>39.1</b>	26.1	ns	30.1	0.9
<sup>A</sup> <i>Romulea longifolia</i>	36.2	<b>59.4</b>	30.4	14.5	10.1	***	30.1	0.3

**Table S3. Significance of relationships between disturbance, climatic and landscape variables, and the most frequent taxa found on Queens Domain.**

The significance of the relationship was determined by; ANOVA or Chi-square and is represented by \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ , ns = not significant (**Error! Reference source not found.**). The impact of the relationship on taxon frequency is represented by + = positive effect on taxon occurrence related to an increase in the factor; - = negative effect on taxon occurrence related to an increase in the factor. A = non-native taxon, B = both non-native and native species in taxon

Taxon	Management condition <sup>1</sup>	Mown status <sup>2</sup>	No. of fires in the decade previous to surveying <sup>1</sup>	Time since fire <sup>1</sup>	Burnt status (2 years prior to surveying) <sup>2</sup>	Summer radiation <sup>3</sup>	Equinox radiation <sup>3</sup>	Winter radiation <sup>3</sup>	Slope (°) <sup>3</sup>	Aspect <sup>3</sup>	Altitude (m) <sup>3</sup>
<i>Annual herbs</i>											
<sup>A</sup> <i>Centaureum erythraea</i>	ns	ns	ns	ns	ns	***-	***-	***-	ns	***+	***+
<sup>A</sup> <i>Cirsium vulgare</i>	ns	**-	***+	ns	*+	*+	**+	**+	ns	*-	**-
<sup>A</sup> <i>Petrorhagia nanteuilii</i>	ns	***-	***+	***-	***+	*+	***+	***+	ns	***-	ns
<i>Senecio</i> spp.	ns	***-	**+	***-	**+	*-	**-	**-	ns	ns	**+
<sup>A</sup> <i>Sonchus</i> spp.	ns	***-	***+	*-	***+	ns	ns	ns	ns	***-	ns
<sup>A</sup> <i>Vicia</i> spp.	ns	**+	*-	**+	ns	ns	*-	ns	**-	***+	ns
<i>Perennial herbs</i>											
<i>Acaena echinata</i>	ns	ns	ns	*-	ns	ns	ns	ns	ns	***+	**+
<i>Chrysocephalum apiculatum</i>	ns	ns	*-	ns	*-	ns	**+	**+	ns	ns	***-
<i>Geranium potentilloides</i>	ns	ns	ns	ns	ns	ns	ns	ns	ns	*+	*+
<i>Gonocarpus tetragynus</i>	ns	ns	ns	ns	ns	***-	***-	***-	**+	***+	**+
<i>Goodenia lanata</i>	ns	ns	ns	**-	*+	ns	ns	ns	ns	ns	ns
<i>Hackelia suaveolens</i>	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<sup>A</sup> <i>Hypochaeris radicata</i>	ns	***+	**-	**+	ns	ns	ns	ns	ns	**+	ns

<i>Leptorhynchos nitidulus</i>	ns	ns	ns	*-	ns	ns	ns	ns	ns	ns	ns	ns
<i>Leptorhynchos squamatus</i>	ns	***+	***-	ns	*-	ns	*-	***-	**-	***+	ns	ns
<sup>B</sup> <i>Linum</i> spp.	ns	*-	*+	ns	**+	ns	*-	*-	ns	***-	ns	ns
<sup>A</sup> <i>Lysimachia arvensis</i>	*3	ns	***+	ns	*+	ns						
<sup>A</sup> <i>Medicago</i> spp.	ns	***+	**-	*+	ns	*+	ns	ns	**-	*+	ns	ns
<i>Oxalis perennans</i>	ns	ns	ns	ns	ns	*-	**-	*-	ns	ns	ns	ns
<sup>A</sup> <i>Plantago lanceolata</i>	ns	***+	***-	**+	ns	ns	*-	*-	*-	***+	ns	ns
<i>Plantago varia</i>	ns	ns	*-	ns	**-	ns	*-	*-	ns	***+	ns	ns
<i>Ranunculus lappaceus</i>	ns	ns	ns	ns	ns	***-	***-	***-	***+	***+	**+	ns
<sup>A</sup> <i>Sanguisorba minor</i>	ns	ns	ns	ns	ns	ns	ns	ns	ns	**-	ns	ns
<sup>A</sup> <i>Trifolium</i> spp.	*1	***+	ns	*+	ns	*+	ns	*-	***-	***+	ns	ns
<sup>A</sup> <i>Urospermum dalechampii</i>	ns	*+	*-	*+	ns	***+	***+	***+	ns	ns	***-	ns
<i>Wahlenbergia</i> spp.	ns	ns	ns	ns	ns	**-	*-	*-	ns	ns	**+	ns
<i>Annual grasses</i>												
<sup>A</sup> <i>Aira</i> spp.	ns	***+	ns	ns	ns	ns	ns	*-	***-	*+	ns	ns
<sup>A</sup> <i>Briza</i> spp.	ns	ns	*+	ns	ns	ns	***-	***-	ns	*+	***+	ns
<sup>A</sup> <i>Vulpia</i> spp.	ns	***+	ns	ns	*-	*+	*-	***-	***-	***+	ns	ns
<i>Perennial grasses</i>												
<i>Anthosachne scabra</i>	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
<i>Austrostipa</i> spp.	*3	ns	*+	ns	ns	*-	ns	ns	*+	ns	**+	ns
<sup>A</sup> <i>Dactylis glomerata</i>	***3	***+	ns	***+	ns	ns	*-	ns	**-	ns	ns	ns
<i>Deyeuxia quadriseta</i>	ns	**-	ns	ns	ns	***-	***-	***-	**+	ns	***+	ns
<i>Dichelachne</i> spp.	ns	*-	**+	***-	**+	ns	ns	ns	ns	ns	ns	***+
<sup>A</sup> <i>Holcus lanatus</i>	ns	ns	ns	***+	ns	**+	ns	ns	***-	**+	*	-
<i>Lachnagrostis aemula</i>	*1	ns	**-	*-	ns	ns	ns	ns	ns	***+	*+	ns
<sup>A</sup> <i>Lolium perenne</i>	ns	***+	***-	***+	**-	ns	*-	**-	**-	***+	*-	ns
<i>Microlaena stipoides</i>	ns	*-	***+	ns	**+	ns	**-	***-	ns	ns	*+	ns

<i>Pentapogon quadrifidus</i>	ns	**+	ns	ns								
<i>Poa rodwayi</i>	ns	ns	ns	ns	*-	ns	ns	ns	ns	ns	***+	ns
<i>Rytidosperma spp.</i>	ns	ns										
<i>Themeda triandra</i>	ns	*+	ns	ns								
<i>Sedges and rushes</i>												
<i>Carex breviculmis</i>	ns	ns	***-	ns	**+							
<i>Dianella spp.</i>	ns	***-	ns	ns	ns	***-	ns	ns	ns	***+	ns	
<i>Lepidosperma gunnii</i>	ns	ns	ns	*-	ns	**+						
<i>Lepidosperma laterale</i>	ns	***-	ns	**-	ns	**-	*+	***+	***+	***-	***-	*
<i>Lomandra longifolia</i>	ns	***-	ns	ns	ns	***-	***-	***-	***+	ns	ns	
<i>Schoenus apogon</i>	ns	ns	ns	*+	ns	ns	***-	***-	ns	***+	*+	
<i>Shrubs</i>												
<sup>A</sup> <i>Acacia pycnantha</i>	ns	ns	ns	*-	ns	ns	***+	**+	ns	*-	ns	
<i>Bossiaea prostrata</i>	ns	ns	ns	**-	ns	ns	ns	*-	ns	*+	***+	
<sup>A</sup> <i>Chrysanthemoides monilifera</i>	ns	*-	ns	ns	ns	***-	ns	ns	**+	ns	ns	
<i>Hibbertia hirsuta</i>	ns	ns	ns	ns	ns	ns	***-	***-	ns	***+	***+	
<i>Lissanthe strigosa</i>	ns	ns	ns	ns	ns	*-	***-	***-	ns	***+	**+	
<i>Olearia ramulosa</i>	ns	***-	ns	ns	ns	***-	***-	***-	***+	**+	***+	
<i>Pimelea humilis</i>	ns	ns	ns	ns	ns	***-	ns	***-	**+	ns	ns	
<sup>A</sup> <i>Rosa rubiginosa</i>	ns	ns	**-	*+	ns	*-	***-	***-	ns	ns	ns	
<i>Styphelia humifusa</i>	ns	ns	*+	ns	ns	***-	***-	***-	**+	ns	***+	
<i>Trees</i>												
<i>Acacia mearnsii</i>	ns	**-	**+	ns	ns							
<i>Allocasuarina verticillata</i>	***3	***-	***+	***-	***+	ns	**+	**+	***+	***-	*+	
<i>Bursaria spinosa</i>	ns	**-	ns	ns	ns	***-	ns	ns	***+	*+	ns	
<i>Dodonaea viscosa</i>	ns	**-	**+	*	ns	ns						
<i>Eucalyptus globulus</i>	ns	ns	ns	ns	ns	***-	***-	***-	ns	***+	ns	
<i>Eucalyptus pulchella</i>	ns	*	ns	ns	ns	**-	***-	***-	ns	***+	*+	
<i>Eucalyptus viminalis</i>	ns	**+	**+									

## *Geophytes*