

[10.1071/CP24156](https://doi.org/10.1071/CP24156)

Crop & Pasture Science

Supplementary Material

Microbial genes highlight different trends in short term for N cycling in historical alpine pastures

Salvatore Raniolo^{A,}, Laura Maretto^A, Maurizio Ramanzin^A, Piergiorgio Stevanato^A, Giuseppe Concheri^A, Andrea Squartini^A, and Enrico Sturaro^A*

^ADepartment of Agronomy, Food, Natural Resources, Animals and Environment, DAFNAE University of Padova, Viale dell'Università 16, Legnaro (PD) 35020, Italy.

*Correspondence to: Salvatore Raniolo Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE), University of Padova, Viale dell'Università 16, Legnaro (PD) 35020, Italy
Email: salvatore.raniolo@unipd.it

Supplementary Materials

Table S1. ANOVA and post-hoc test based on permutation test analyzing the abundance of 16S, nifH, nirK, nosZ, archaeal amoA (AOA) and eubacterial amoA (AOB) genes.

	16S			nifH			nirK					
	Df	Sum Sq	Pr(Prob)	Df	Sum Sq	Pr(Prob)	Df	Sum Sq	Pr(Prob)			
grazing	1	0.08	0.75	1	0.025	0.7451	1	0.21	0.51			
sampling time	3	0.67	0.09	.	3	8.33	<2e-16	***	3	4.48	0.030	*
grazing:sampling time	3	0.08	0.77	3	0.15	0.9433	3	0.15	0.99			
pasture area	2	0.80	0.10	2	0.90	0.4533	2	0.85	0.47			
grazing:pasture area	2	0.30	0.21	2	0.010	1	2	0.21	0.96			
Sampling time:pasture area	6	0.18	1.00	6	2.12	0.632	6	1.49	0.67			
Residuals	54	7.68		54	21.62		54	25.99				
	nosZ			amoA AOA			amoA AOB					
	Df	Sum Sq	Pr(Prob)	Df	Sum Sq	Pr(Prob)	Df	Sum Sq	Pr(Prob)			
grazing	1	5.27	0.027	*	1	0.79	0.56	1	0.93	0.58		
sampling time	3	5.83	0.14	3	4.94	0.60	3	8.84	0.58			
grazing:sampling time	3	0.69	1	3	3.07	0.95	3	2.83	0.73			
pasture area	2	9.18	0.0088	**	2	3.21	0.74	2	9.93	0.19		
grazing:pasture area	2	2.69	0.16	2	0.12	0.90	2	14.16	0.12			
Sampling time:pasture area	6	5.95	0.33	6	10.69	0.93	6	7.23	0.91			
Residuals	54	51.60		54	191.07		54	182.97				
	nosZ			nifH			nirK					
Pasture area	means	groups		Sampling time	means	groups	Sampling time	means	groups			
Juribello	4.72	ab		T1	4.71	a	T1	4.97	ab			
Vallazza	4.08	b		T2	5.02	a	T2	5.27	ab			
Venegiota	4.91	a		T3	4.15	b	T3	4.77	b			
				T4	4.95	a	T4	5.41	a			
Grazing	means	groups										
Yes		4.84	a									
No		4.30	b									




Fig. S1. Optimal number of clusters among sampling sites estimated with the Silhouette method.




Fig. S2. Corrplot of pedological and morphological variables based on Kendall rank's coefficient coefficient (* = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$).