

**Animal detections increase by using a wide-angle camera trap model but not by periodically repositioning camera traps within study sites**

Anke Seidlitz<sup>A,E</sup>, Kate A. Bryant<sup>A,B</sup>, Nicola J. Armstrong<sup>C</sup> and Adrian F. Wayne<sup>D</sup>

<sup>A</sup>Environmental and Conservation Sciences, Murdoch University, Murdoch, WA 6150, Australia.

<sup>B</sup>Centre for Climate-Impacted Terrestrial Ecosystems, Harry Butler Institute, Murdoch University, Murdoch, WA 6150, Australia.

<sup>C</sup>Mathematics and Statistics, Murdoch University, Murdoch, WA 6150, Australia.

<sup>D</sup>Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Locked Bag 2, Manjimup, WA 6258, Australia.

<sup>E</sup>Corresponding author. Email: [anke.seidlitz@gmx.net](mailto:anke.seidlitz@gmx.net)

## Supplementary Material

**Table S1.** List of bird species confidently identified

Common name	Scientific name
Yellow-rumped thornbill	<i>Acanthiza chrysorrhoa</i>
Australian owlet-nightjar	<i>Aegotheles cristatus</i>
Australian ringneck parrot	<i>Barnardius zonarius</i>
Rufous treecreeper	<i>Climacteris rufus</i>
Grey shrike-thrush	<i>Colluricincla harmonica</i>
Australian raven	<i>Corvus coronoides</i>
Laughing kookaburra	<i>Dacelo novaeguineae</i>
Western yellow robin	<i>Eopsaltria griseogularis</i>
Australian magpie	<i>Gymnorhina tibicen</i>
Fairy wren	<i>Malurus sp.</i>
Golden whistler	<i>Pachycephala pectoralis</i>

Scarlet robin	<i>Petroica boodang</i>
Common bronzewing	<i>Phaps chalcoptera</i>
Western rosella	<i>Platycercus icterotis</i>
Tawny frogmouth	<i>Podargus strigoides</i>
Grey fantail	<i>Rhipidura albiscapa</i>
Grey currawong	<i>Strepera versicolor</i>

---