

Supplementary Material

Evaluation of hepatic and renal effects in rat dams and their offspring after exposure to paracetamol during gestation and lactation

Andréa Morgato de Mello Miyasaki^{A,B,}, Camila Rigobello^{B,*}, Rodrigo Moreno Klein^B, Jefferson Crespigio^C, Karina Keller Flaiban^D, Ana Paula Bracarense^D, Bárbara Cristina Mazzucatto^E, Décio Sabbatini Barbosa^B and Estefânia Gastaldello Moreira^{B,F,G}*

^ADepartment of Pediatrics and Pediatrics Surgery, Universidade Estadual de Londrina, 86051-980, Londrina, PR, Brazil.

^BGraduate Program in Health Sciences, Universidade Estadual de Londrina, 86051-980, Londrina, PR, Brazil.

^CDepartment of Pathology, Universidade Estadual de Londrina, 86057-970, Londrina, PR, Brazil.

^DDepartment of Preventive Veterinary Medicine, Universidade Estadual de Londrina, 86057-970, Londrina, PR, Brazil.

^EDepartment of Veterinary Medicine, State University of Maringá, 87507-190, Umuarama, PR, Brazil.

^FDepartment of Physiological Sciences, Universidade Estadual de Londrina, 86057-970, Londrina, PR, Brazil.

^GCorresponding author. Email: egmoreira@uel.br






Fig. S1. Representative sections of dams' livers stained with HE (A-C). A: normal hepatic architecture (200x magnification). Independently from the treatment group it was observed portal tract with scarce inflammatory cells including eosinophils (B, arrow, 400x magnification) as well as sinusoidal eosinophil (C, circle, 400x magnification). Scale= 25 μ m.

A



B



C




Fig. S2. Representative sections (magnification 400x) of dams' kidney stained with HE. Independently of treatment, loss of brush border (B, arrow) and cytoplasmic vacuolization (C, arrow) were observed in the kidney of the animals. Scale= 25 μ m.