

Supplementary material

Only recent sexual partners contribute to oropharyngeal gonorrhoea positivity: the number of sexual partners over different time periods as an indicator of gonorrhoea and chlamydia infection duration among men who have sex with men

David Priest^A, Tim R. H. Read^{A,B}, Marcus Y. Chen^{A,B}, Catriona S. Bradshaw^{A,B}, Christopher K. Fairley^{A,B,*} and Eric P. F. Chow^{A,B,C,*}

^AMelbourne Sexual Health Centre, Alfred Health, 580 Swanston Street, Carlton, Vic. 3053, Australia.

^BCentral Clinical School, Faculty of Medicine, Nursing and Health Sciences, Monash University, 99 Commercial Road, Melbourne, Vic. 3004, Australia.

^CCorresponding author. Email: Echow@mshc.org.au

Table S1. Oropharyngeal gonorrhoea positivity and associations in MSM attending MSHC, excluding detection methods and condom use for anal sex in multivariable logistic model

Factors	Adjusted OR (95% CI)	P value
Age		
≤24	6.16 (3.79-10.02)	<0.001
25-34	4.91 (3.03-7.97)	<0.001
35-44	2.88 (1.74-4.74)	<0.001
45-54	2.43 (1.43-4.13)	0.001
≥55	1	Ref
Partners (3 months)		
0-1	1	Ref
2-5	1.47 (1.27-1.72)	<0.001
≥6	2.03 (1.69-2.42)	<0.001
Partners (4-12 months)		
0-1	1	Ref
2-5	1.11 (0.97-1.27)	0.142
≥6	1.02 (0.88-1.17)	0.837