

[10.1071/SH24041](https://doi.org/10.1071/SH24041)

Sexual Health

Supplementary Material

Sexual risk and testing for sexually transmissible infections in Aboriginal and Torres Strait Islander peoples and non-Indigenous young South Australians: results of an online survey

Stephen Harfield^{A,B,C,}, Salenna Elliott^A, Federica Barzi^A, Kiara Minto^A, Judith A. Dean^{A,B}, and James Ward^A*

^APoche Centre for Indigenous Health, Faculty of Health and Behavioural Sciences, The University of Queensland, Brisbane, Qld, Australia.

^BSchool of Public Health, Faculty of Medicine, The University of Queensland, Brisbane, Qld, Australia.

^CAboriginal Health Equity, South Australian Health and Medical Research Institute, Adelaide, SA, Australia.

*Correspondence to: Stephen Harfield UQ Poche Centre for Indigenous Health, The University of Queensland, Brisbane, Qld, Australia Email: s.harfield@uq.edu.au

Supplementary materials

Table S1 Missing responses for survey participant characteristics by Indigenous status, South Australia, 'Let's Talk About It 2019'.

	Aboriginal and/or Torres Strait Islander	Non-Indigenous	Overall
	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
Total respondents (N)	231	2062	2293
Gender	2 (1%)	6 (<1%)	8 (<1%)
Age group	4 (2%)	18 (1%)	22 (1%)
Remoteness	9 (4%)	58 (3%)	67 (3%)
Sexuality	2 (1%)	6 (<1%)	8 (<1%)
Marital status	3 (1%)	15 (1%)	18 (1%)
Currently studying	2 (1%)	7 (<1%)	9 (<1%)
Highest level of education completed	3 (1%)	14 (1%)	17 (1%)
Employed	2 (1%)	5 (<1%)	7 (<1%)

Table S2. Missing responses for STI and HIV testing and health care access, by Indigenous status, South Australia, 'Let's Talk About It 2019'.

	Aboriginal and/or Torres Strait Islander	Non- Indigenous	Overall
	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
Sexually active respondents (N)	186	1618	1804
Had a health check in the past 12 months	4 (2%)	62 (4%)	66 (4%)
Ever tested for an STI	4 (2%)	54 (3%)	58 (3%)
Ever tested for HIV	4 (2%)	54 (3%)	58 (3%)
Health check and sexually active (N)	95	605	700
Offered a STI test	0 (0%)	(0%)	0 (0%)
Location of Health check	0 (0%)	0 (0%)	10 (1%)
Ever tested STI (N)	128	746	874
Tested positive for any STI	0 (0%)	1 (<1%)	1 (<1%)
Location of STI test	0 (0%)	0 (0%)	0 (0%)

Location of HIV check	0 (0%)	0 (0%)	0 (0%)
-----------------------	--------	--------	--------

Table S3. Missing responses for sexual activity and factors influencing risk of STIs and HIV, by Indigenous Status, South Australia, ‘Let’s Talk About It 2019’.

	Aboriginal and/or Torres Strait Islander	Non- Indigenous	Overall
	<i>n (%)</i>	<i>n (%)</i>	<i>n (%)</i>
Total respondents (N)	231	2062	2293
Sexually active (oral, vaginal, or anal)	28 (12%)	183 (9%)	211 (9%)
Ever had sexual intercourse (vaginal or anal)	28 (12%)	186 (9%)	214 (9%)
Ever had oral sex	29 (13%)	182 (9%)	211 (9%)
Sexually active respondents (N)	186	1618	1804
Regular sexual partner	28 (12%)	186 (9%)	4 (<1%)
Sexually active people with a regular partner (N)	120	1156	1276

Only have sex with your partner	0 (0%)	2 (<1%)	2 (<1%)
Partner only has sex with you	0 (0%)	2 (<1%)	2 (<1%)
Number of sexual partners in the last year	6 (3%)	103 (6%)	109 (6%)
Relationship with last sexual partner	0 (0%)	12 (1%)	12 (<1%)
Sexual intercourse (N)	180	1516	1696
Condom used at last sexual encounter	0 (0%)	0 (0%)	0 (0%)
Last sex wanted?	1 (1%)	22 (1%)	23 (1%)
Consumed alcohol in past 12 months	3 (2%)	28 (2%)	31 (2%)
Consumed alcohol (N)	166	1480	1646
How many alcohol drinks consumed per session	0 (0%)	2 (<1%)	2 (<1%)
'Drunk' or 'high' during their last sexual encounter	0 (0%)	18 (1%)	18 (1%)
Drug score	0 (0%)	0 (0%)	0 (0%)