

Supplementary material for:

Organic based integrated nutrient management scheme enhances soil carbon storage in rainfed rice (*Oryza sativa*) cultivation

Parijat Saikia^{A,C}, Kushal Kumar Baruah^{A,C}, Satya Sundar Bhattacharya^{B,C} and Chandrima Choudhury^A

^ADepartment of Environmental Science, Tezpur University, Assam 784 028, India.

^BSoil and Agro Bio-engineering Laboratory, Department of Environmental Science, Tezpur University, Assam 784 028, India.

^CCorresponding authors. Email: kkbaruah14@gmail.com; parijatsaikia07@gmail.com; satya72@tezu.ernet.in

Table S0. Two year total C input from crop residue (CR) and from farmyard manure (FYM)

Treatment	Organic based nutrients	Inherent C (0-15 cm) (Kg ha ⁻¹)	Total C input (after two year) (Kg ha ⁻¹)	Inherent + input (Kg ha ⁻¹)	SOC loss during one crop growth period Kg ha ⁻¹ / 18 week (126 days)	C seq (Kg ha ⁻¹) after 2 year (0-15 cm)
T1	-----	24768	-----	24768	32410.20	-1338.0
T2	FYM	24768	490	25258	29135.05	2042.0
T3	FYM	24768	490	25258	32032.99	1622.0
T4	CR	24768	200	24968	34498.03	1052.0
T5	CR	24768	200	24968	31835.40	1972.0
T6	FYM+CR	24768	690	25458	28488.41	4492.0
T7	FYM+CR	24768	690	25458	25249.18	3452.0

Table S1. Values of soil moisture content (SMC, %) (two year values)

Values are mean and similar letters are not significantly different in the same column at $P < 0.05$ according to Duncan's multiple range test. T1 (100% NPK), T2 (100% NPK + FYM, 10 ton ha $^{-1}$), T3 (80% NPK + FYM, 10 ton ha $^{-1}$), T4 (100% NPK + CR 5 ton ha $^{-1}$), T5 (80% NPK + CR 5 ton ha $^{-1}$), T6 (100% NPK + CR 5 ton ha $^{-1}$ + FYM 10 ton ha $^{-1}$), T7 (80% NPK + CR 5 ton ha $^{-1}$ + FYM 10 ton ha $^{-1}$)

	Year 1: SMC (%) (0-15 cm)				Year 1: SMC (%) (15-30 cm)			
T1	33.29ab	30.70ab	28.50a	26.52a	31.72a	32.40ab	30.32a	29.05a
T2	35.97b	32.85c	30.62a	27.85a	34.50ab	33.55ab	31.45ab	29.82ab
T3	34.40ab	32.65c	31.12a	27.72a	36.02b	33.27ab	32.27ab	30.25ab
T4	34.65ab	32.50bc	30.15a	26.15a	33.45ab	32.62ab	31.275ab	29.70ab
T5	32.40a	30.55a	29.00a	26.75a	32.71ab	31.80a	30.75ab	29.37a
T6	36.07b	34.77d	31.07a	28.65a	35.92b	34.30b	32.90b	31.02b
T7	36.40b	32.50bc	31.20a	28.80a	35.15b	34.05ab	32.90b	30.42ab
	Year 2: SMC (%) (0-15 cm)				Year 2: SMC (%) (15-30 cm)			
T1	31.07a	31.55a	28.23a	23.86a	28.63abc	25.89a	26.58ab	23.54ab
T2	34.92ab	33.72a	30.80b	27.26bc	28.35bc	26.64ab	26.12ab	22.78a
T3	37.67b	34.17a	31.17b	28.50c	28.38bc	27.51bc	27.95c	23.85ab
T4	37.60b	32.85a	29.60ab	24.63a	27.15bc	26.99b	25.42a	23.34ab
T5	38.55b	32.31a	28.20b	25.27ab	29.56d	26.82b	26.21ab	23.82ab
T6	40.54b	32.62s	31.03b	28.41c	28.19b	28.30c	27.02b	23.97ab
T7	38.82b	33.22a	30.66ab	27.21bc	29.43cd	29.63d	26.73abc	23.42b
P-values year	0.011	0.242	0.005	0.015	0.000	0.000	0.000	0.000
P-values treatment	0.017	0.036	0.55	0.000	0.035	0.000	0.003	0.07
l.s.d.	1.61	0.886	0.868	0.77	0.781	0.562	0.558	0.425

Table S2. Values of soil temperature (ST, °C) (two year values)

Values are mean and similar letters are not significantly different in the same column at $P < 0.05$ according to Duncan's multiple range test. T1 (100% NPK), T2 (100% NPK + FYM, 10 ton ha $^{-1}$), T3 (80% NPK + FYM, 10 ton ha $^{-1}$), T4 (100% NPK + CR 5 ton ha $^{-1}$), T5 (80%NPK + CR 5 ton ha $^{-1}$), T6 (100%NPK + CR 5 ton ha $^{-1}$ + FYM 10 ton ha $^{-1}$), T7 (80%NPK + CR 5 ton ha $^{-1}$ + FYM 10 ton ha $^{-1}$)

Treatment	Year 1: ST (°C) (0-15 cm)				Year 1: ST (°C) (15-30 cm)			
	MTL	PI	FL	MT	MTL	PI	FL	MT
T1	30.27a	29.18a	29.81a	27.62bc	29.30a	28.23a	29.15a	27.10a
T2	29.90a	29.58a	28.10a	24.85a	28.94a	28.63a	27.55a	24.23a
T3	29.91a	29.48a	28.22a	25.23ab	28.95a	28.57a	27.67a	24.71a
T4	30.92a	29.77a	29.18a	28.28c	30.14a	28.827a	28.59a	27.76a
T5	31.08a	29.44a	29.69a	28.15c	30.12a	28.49a	29.14a	27.62a
T6	29.87a	29.27a	29.05a	25.86abc	28.91a	28.32a	28.55a	25.34a
T7	31.04a	29.49a	28.70a	26.63abc	30.08a	28.36a	28.15a	26.11a
Year 2: ST(°C) (0-15 cm)					Year 2: ST (°C) (15-30 cm)			
T1	31.31a	27.69a	28.44a	27.74b	30.75a	27.06a	27.93a	27.04b
T2	31.32a	28.34a	26.87a	23.85a	30.73a	27.73a	26.20a	23.15a
T3	31.12a	28.24a	26.71a	28.02b	30.56a	27.63a	26.10a	27.22b
T4	32.13a	28.53a	28.39a	27.00b	31.51a	27.79a	27.78a	26.30b
T5	32.03a	28.20a	28.46a	27.15b	31.4a	27.59a	27.73a	26.35b
T6	31.08a	28.31a	27.64a	24.86ab	30.52a	27.52a	27.03a	24.16ab
T7	32.25a	28.25a	27.47a	25.08ab	31.64a	27.64a	26.86a	24.38ab
P-values year	0.001	0.000	0.001	0.375	0.000	0.006	0.001	0.195
P-values treatment	0.154	0.942	0.051	0.001	0.140	0.957	0.060	0.001
l.s.d.	0.585	0.574	0.648	0.871	0.582	0.600	0.655	0.865

Table S3. Values of soil pH (two year values)

Values are mean and similar letters are not significantly different in the same column at $P < 0.05$ according to Duncan's multiple range test. T1 (100% NPK), T2 (100% NPK + FYM, 10 ton ha $^{-1}$), T3 (80% NPK + FYM, 10 ton ha $^{-1}$), T4 (100% NPK + CR 5 ton ha $^{-1}$), T5 (80%NPK + CR 5 ton ha $^{-1}$), T6 (100%NPK + CR 5 ton ha $^{-1}$ + FYM 10 ton ha $^{-1}$), T7 (80%NPK + CR 5 ton ha $^{-1}$ + FYM 10 ton ha $^{-1}$)

Treatment	Year 1: pH (0-15 cm)				Year 1: pH (15-30 cm)			
	MTL	PI	FL	MT	MTL	PI	FL	MT
T1	5.08b	5.19ab	5.44b	5.39abc	5.06a	5.41a	5.73ab	5.52b
T2	4.90a	5.06a	5.32ab	5.24a	5.14ab	5.34a	5.44a	5.34a
T3	4.94a	5.17ab	5.23a	5.35ab	5.18ab	5.46a	5.47ab	5.42ab
T4	5.08b	5.32ab	5.41ab	5.38abc	5.24ab	5.49a	5.66abc	5.51b
T5	4.90a	5.28ab	5.38ab	5.39abc	5.16ab	5.55a	5.67abc	5.50ab
T6	5.03ab	5.41b	5.44b	5.47bc	5.23ab	5.50a	5.72bc	5.54b
T7	4.98ab	5.24ab	5.43ab	5.58c	5.30b	5.44a	5.76c	5.58b
Year 2: pH (0-15 cm)				Year 2: pH (15-30 cm)				
T1	5.07b	5.23ab	5.32ab	5.34ab	5.14a	5.38a	5.25a	5.48bc
T2	4.86a	4.98a	5.31ab	5.41bc	5.20a	5.34a	5.36ab	5.41ab
T3	4.88a	4.99ab	5.28a	5.34ab	5.17a	5.33a	5.37ab	5.34a
T4	5.05b	5.17ab	5.36abc	5.35ab	5.25a	5.51b	5.45b	5.47bc
T5	4.95ab	5.11ab	5.43bc	5.31a	5.21a	5.37a	5.41b	5.47bc
T6	5.07b	5.10ab	5.48c	5.50d	5.28a	5.44ab	5.46b	5.54c
T7	5.05b	5.30ab	5.50c	5.47cd	5.30a	5.36a	5.47b	5.55c
P values year	0.917	0.015	0.893	0.717	0.352	0.018	0.000	0.369
P values								
treatment	0.000	0.036	0.004	0.001	0.032	0.04	0.011	0.000
LSD	0.044	0.083	0.054	0.051	0.058	0.05	0.0647	0.044

Table S4. Correlation of soil and plant parameters

*= correlation is significant at 0.05 level of significance; **= Significant at 0.01 level of significance; NS= Non significant; SOC= soil organic carbon, LB= leaf biomass,

RB=root biomass, N= mineralisable nitrogen, DOC=dissolved organic carbon, TC=total carbon, SMC= soil moisture content

	SOC (0-15 cm)	SOC (15-30 cm)	Storage (0-15 cm)	Storage (15-30 cm)	LB	RB	N (0-15 cm)	N (15-30 cm)	DOC (0- 15cm)	DOC (15-30 cm)	TC (0-15 cm)	TC (15-30 cm)	Photosyn thesis	SMC (0-15 cm)	SMC (15-30 cm)
SOC (15-30 cm)	0.23**	1													
Storage (0-15 cm)		0.78**	NS	1											
Storage (15-30 cm)		0.18*	0.98**	NS	1										
LB	0.49**	NS	0.53**	NS	1										
RB	0.33**	0.14*	0.31**	NS	0.4**	1									
N (0-15 cm)	0.31**	0.26**	0.34**	0.23*	0.4**	0.24**	1								
N (15-30 cm)	0.43**	0.23**	0.4**	0.17*	0.4**	NS	0.42**	1							
DOC (0-15cm)	0.46**	0.20**	0.47**	0.18**	0.4**	0.51**	0.24**	0.19**	1						
DOC (15-30 cm)	0.52**	NS	0.55**	NS	0.5**	0.49**	0.27**	0.31**	0.74**	1					
TC (0-15 cm)	0.46**	0.5**	0.45**	0.47**	0.33**	0.29**	0.35**	0.39*	0.42**	0.37**	1				
TC (15-30 cm)	NS	0.68**	NS	0.66**	NS	0.15*	0.28**	0.22**	0.19**	NS	0.41*	1			
Photosynthesis	0.25**	0.3**	0.25**	0.29**	NS	0.25**	0.26**	0.25*	0.27**	0.25**	0.33*	0.33**	1		
SMC (0-15 cm)	0.23**	0.29**	0.19**	0.27**	0.14*	-0.24**	0.39**	0.43**	NS	NS	0.21**	0.31**	0.13*	1	
SMC (15-30 cm)	NS	0.75**	NS	0.73**	NS	NS	0.24**	0.27**	NS	NS	0.36**	0.74**	0.19**	0.42**	1