

[10.1071/WR24175](https://doi.org/10.1071/WR24175)

Wildlife Research

Supplementary Material

Invasive wild pig carcass availability does not affect coyote abundance in South Carolina, USA

Heather E. Gaya^{A,}, Gino J. D'Angelo^A, Mark Vukovich^{B,D}, Jordan L. Youngmann^A, Stacey L. Lance^C, and John C. Kilgo^B*

^A Warnell School of Forestry and Natural Resources, University of Georgia, 180 E. Green Street, Athens, GA 30605, USA.

^B United States Department of Agriculture Forest Service Southern Research Station, PO Box 700, New Ellenton, SC 29809, USA.

^C University of Georgia, Savannah River Ecology Laboratory, Drawer E, Aiken, SC 29802, USA.

^DPresent address: United States Department of Agriculture Forest Service, Shawnee National Forest, 602 North First Street, Vienna, IL 62995, USA.

*Correspondence to: Heather E. Gaya Warnell School of Forestry and Natural Resources, University of Georgia, 180 E. Green Street, Athens, GA 30605, USA Email: heather.e.gaya@gmail.com


Supplemental Figures

Figure S1. Heat map of scat collected at the Savannah River Site, South Carolina, USA from July 2014 to July 2017. Points represent the number of scats collected within each 500-meter hex.

Areas in light blue (no hex shapes) represent hexes where 1 or fewer scats were collected.




Figure S2. Distance to the nearest carcass pile for all scat collected at the Savannah River Site, South Carolina, USA from July 2014 to July 2017. Carcasses were not present in piles on the landscape prior to July 2015.